



COMPARISON OF LUNG CAPACITY CHANGE IN ELDERLY WOMEN

Faten A .Hamza , ph.D.* and Reham E. Elden, ph.D.**

Department of Physical Therapy for Cardiovascular-Respiratory Disorders and Geriatrics, Faculty of Physical Therapy ,Delta University for science and Technology, Cairo, Egypt* and Physical Therapy Department of Moneira Hospital, Ministry of health Cairo, Egypt**

Abstract

Background: changes in the thoracic structure, breathing muscle weakness and VC decreases in the elderly, if exercise programs for the elderly can be developed in accordance with age, they will help to prevent VC decreases. **Aim of the study:** To examine the change of lung capacity of elderly women. **Subjects and Methods:** The subjects of this study were 55 healthy elderly women over 60 years of age. Subjects were classified in the following age categories: 60s, 70s and 80s. For the pulmonary function test in Damietta chest Hospital a spirometer (Pony FX, COSMED Inc., Italy) was used. The item for measurement of pulmonary function in elderly women was maximum-effort expiratory spirometer (MES). **Results:** Among items of maximum-effort expiratory spirometer, a significant difference according to age was demonstrated in forced vital capacity, forced expiratory volume in 1 second, peak expiratory flow, maximum expiratory flow 75%, maximum expiratory flow 50%, and inspiratory capacity. **Conclusion:** - as age increases lung capacity decreases remarkably. To prevent decrease in lung capacity as age progresses, Exercise program is essential to beginning at an early age

Keywords: Lung capacity, Elderly women.

Introduction

A respiratory disturbance is strongly associated with older adults', and pulmonary function decreases as aging progresses.⁽²⁷⁾The population increased by accelerating the Improved standard of living and Improved medical technology.⁽¹⁾The data collected by the Central Agency for Public Mobilization and Statistics of Egypt, the ratio of the elderly population (60 years and above) for both males and females during the period of appreciation (2006 -2013) Shows high proportion of older males from 5.9% in 2006 to 11.1% in 2013, as well as higher percentage of females from 6.1% in 2006 to 12.6% in 2013, also notes the high rate of older females than males during the period (2006-2013)increasing the preparation of the role of the elderly in addition to multiple types of pensions. ⁽²⁾The increase of the aged population has produced a positive effect—the increase of the average life expectancy. On the other hand, the negative effect is that there is a large number of

elderly who are suffering from the increase of chronic disease, which results in a lower quality of life.⁽³⁾ Therefore there has been increasing interest in maintaining good health and improving the quality of life⁽¹⁾. This is particularly significant according to the research about mortality from respiratory disease. Among people aged 60 and over, 22.6 per 100,000 have chronic respiratory disease, which was the fifth major cause of death, and it has been on the rise.⁽⁴⁾ As a result, interest in chronic obstructive pulmonary disease (COPD) has been increasing recently. Research shows that for the elderly who have COPD, as breathing difficulty increases, fear of death and symptoms of depression increase, which causes a decline in the quality of life⁽⁵⁾. Breathing difficulty is the most serious symptom for patients with COPD, which is especially life-threatening.⁽⁶⁾ The decrease of functional capacity from breathing difficulties.⁽⁷⁾ and motor disturbance.⁽⁸⁾ lower the quality of life drastically. Additional research comparing healthy people to patients with COPD has also demonstrated a lower standard of life related to health among those suffering from impaired lung function.⁽⁹⁾ In addition, social and economic expenses caused by COPD are forecasted to rise steadily⁽¹⁰⁾, demonstrating the need to take additional measures to improve lung function. Most research focused on improving pulmonary function. Shin⁽¹¹⁾ Examined the effects of walking exercise programs on cardiopulmonary function in female seniors. As a result, maximum oxygen intake, heart rate, relaxed blood pressure, and maximum forced vital capacity have improved as they were stabilized. More recently, Kang et al. ⁽¹²⁾ Demonstrated how a 12 week complex exercise training program affects cardiopulmonary function and the risk factor of metabolic syndrome of elderly persons who live in the countryside. In their study, maximal voluntary ventilation (MVV) has increased significantly in the group who performed a resistant workout and walking program. VO₂max and MVV both increased meaningfully in the group that performed aerobic and yoga exercises. Accordingly, Kang said that complex exercise must include a resistant workout, walking, aerobics, and yoga in order to improve pulmonary function in the elderly. Based on respiratory rehabilitation exercise programs improving pulmonary function on patients who have COPD (such as chronic bronchial trouble and pulmonary emphysema), research has demonstrated how exercise can be productive. Patients with general lung diseases such as interstitial lung disease, fibrous cystoma, bronchial expansion syndrome, thorax malformation, lung transplantation, pneumonectomy, and neuromuscular disease, can all benefit from exercise treatments⁽¹³⁾. However, results of research that has developed and applied various types of respiration rehabilitation for COPD have not been consistent. One of the reasons for the discrepancies is that the age of the participants was not taken into account when exercise programs were developed and applied. Therefore most research programs have overlooked the fact that exercise intensity depends on age. Therefore, this study was designed to find out how much pulmonary function would vary as age increases. On the basis of the results of this study, an effort was made to provide basic information on developing an exercise program that takes into consideration the age of the individual.

Subjects and Methods

A Fifty-five female senior citizens who agreed conduct in this study, subject's women over the age of sixty able to understand the instruction , clear cognitive abilities on lung capacity measurement. Writing to take part listening to the purpose of the study. They were all non-smokers, no history of lung disease. This study took place in Damietta chest hospital, Extended from October 2016 to December 2016. Written informed consent was received from each participant. In the 60–69 years age group,

age on average was 64.6 ± 2.7 years; height on average was 155.8 ± 3.1 cm; and weight on average was 57.3 ± 7.2 kg. In the 70–79 years group, age on average was 74.2 ± 3.1 years; height on average was 153.5 ± 5.0 cm; and weight on average was 57.0 ± 6.8 kg. In the 80–89 years group, age on average was 81.8 ± 2.0 years; height on average was 147.2 ± 5.5 cm; and weight on average was 54.6 ± 6.1 kg. This measuring instrument can check the amount and the speed of air exiting from the lungs. The measuring instrument of lung capacity used in this study was a digital pulmonary function measurement (Pony FX, COSMED Inc., Italy). Maximum-effort Expiratory Spirogram (MES) was measured in this study. This MSE measurement was taken while the subject was in a straight sitting posture in a chair. The subject's waist and shoulders were straight. Legs were opened to the width of the subject's shoulders, and feet were placed vertically on the floor. Subsequently, the subject's nose was closed with her hand. And the measuring instrument was held with one hand and inserted between the teeth with the lips closed over the measuring instrument. After 3–4 normal breaths, the subject inhaled and exhaled quickly and deeply while the measuring instrument was in place. The subject then held her breath for six seconds. The subject was given a one-minute rest between measurements. Three measurements were taken for each subject⁽¹⁴⁾ Unless dizziness was experienced, in which case only two measurements were taken. An average of the measurements was taken for each subject for analysis. These measurements were taken at the senior citizens center. This study designed to assessment the effect on the variation of lung capacity in elderly women, divided into three groups: age 60 to 69, age 70 to 79, and age 80 to 89. Seven factors from the pulmonary function tests examine the difference among dependent variables based on age, one-way ANOVA was performed. Every significant level for data analysis was under 5%.

Results

There was a statistical difference among the groups as forced vital capacity (FVC) ($p < 0.05$), forced expiratory volume in 1 second (FEV_1) ($p < 0.05$) and peak expiratory flow 1/sec (PEF) ($p < 0.05$). In the result of post analysis, FVC, FEV_1 , PEF and MEF 50% for subjects in their 70s was reduced compared to those in their 60s. Also, above all items for a subject in their 80s was reduced compared to those in their 70s. There was a statistical difference among the groups as maximum expiratory flow 75% (MEF 75%) ($p < 0.05$) and MEF 50% ($p < 0.05$). In the result of post analysis, MEF 75% and MEF 50% for subjects in their 80s were reduced compared to those in their 60s. However, in case of MEF 75% and MEF 50%, there was no difference significantly between 60s and 70s, and between 70s and 80s. There was a statistical difference among the three groups regarding inspiratory capacity (IC) ($p < 0.05$). In the result of post analysis, IC for subjects in their 80s was reduced compared to those in their 60s and it of 70s. But, in case of IC, there was no difference significantly between 60s and 70s. The result of pulmonary function based on the age of normal elderly women is shown in (Table 1). A statistical difference was demonstrated among the three groups.

Table (1). The changes of lung capacity in elderly women

Variables	60–69 years	70–79 years	80–89 years
FVC (l)*	$2.43 \pm 0.32^{\dagger\ddagger}$	$2.11 \pm 0.38^{\#}$	1.57 ± 0.32
FEV_1 (l)*	$1.98 \pm 0.28^{\dagger\ddagger}$	$1.69 \pm 0.35^{\#}$	1.31 ± 0.27
PEF (l)*	$4.75 \pm 1.19^{\dagger\ddagger}$	4.05 ± 1.43	3.07 ± 0.85
MEF 75%*	$4.37 \pm 1.05^{\dagger\ddagger}$	3.68 ± 1.35	2.75 ± 0.74

MEF 50%*	2.59 ± 0.7 [‡]	2.23 ± 0.88	1.75 ± 0.49
MEF 25%	0.98 ± 0.47	0.76 ± 0.29	0.78 ± 0.37
IC (l)*	1.84 ± 0.31 [‡]	1.57 ± 0.42 [#]	1.14 ± 0.38

Mean ± SD. *p<0.05

(60–69 year) > (70–79 year), (60–69 year) > (80–89 year), (70–79 year) > (80–89 year)

FVC: forced vital capacity; FEV1: forced vital capacity in one second; PEF: peak expiratory flow; MEF: maximum expiratory flow; IC: inspiratory capacity




Figure (1). The changes of lung capacity in elderly women (60–69 year) > (70–79 year), (60–69 year) > (80–89 year), (70–79 year) > (80–89 year)
FVC: forced vital capacity; FEV1: forced vital capacity in one second; PEF: peak expiratory flow values (Mean ± SD). *p<0.05




Figure (2). The changes of lung capacity in elderly women (60–69 year) > (70–79 year), (60–69 year) > (80–89 year), (70–79 year) > (80–89 year)
MEF: maximum expiratory flow; IC: inspiratory capacity values (Mean ± SD). *p<0.05

Discussion

The external environment can cause lung problems that are respiratory infections, exacerbated by toxins, and smoking with aging. Additionally, physiological capacity decreases and breathing mechanism problems, Occur due to limitations in chest wall movement and decreases in lung elasticity.⁽¹⁵⁾ Furthermore, fat content and elastin in lungs decrease as age increases, breathing muscle strength weakens, and pleural elasticity and fibers decrease in length and diameter, further weakening the lungs elastic contractility.⁽¹⁶⁾ As a result, lung tissues become stretched. Hence, changes in age-related breathing mechanisms cause limitations in respiratory airflow, and those limitations are known to reduce the rates of FEV₁ and FEV₁/FVC in breathing function tests, which show the occurrence of air trapping and hyperventilation as well as increases in residual volume and functional residual volume.⁽¹⁷⁾ Evidences show, Vital capacity (VC) in accordance with age among participants between the ages of 20 to 70, as age increases, VC correlate negatively.⁽¹⁸⁾ In a follow-up of the measurement changes in lung function among participants divided into four groups according to age (60 to 64, 65 to 69, 70 to 75, and over 75 years of age), and FEV₁, FVC, and VC decrease as age increases in all participants, both male and female, in all four groups.⁽¹⁹⁾ VC, FEV₁ and physical function and mobility in 1,022 women aged ≥ 75 years assessed; older women exhibited inferior pulmonary function as well as reduced physical function and mobility, FEV₁, FVC, and FEV₁/FVC decrease as age increases in participants in their 60s, 70s, and 80s.⁽²⁰⁾ This result corresponds with the results of previous studies. While previous studies did not consider measurement variables for peripheral bronchi, this study tested the numbers of MEF 75%, MEF 50%, and MEF 25% in order to check the changes in peripheral bronchi as age increases. The study confirmed that peripheral bronchi also decrease as age increases. In these decreases in peripheral bronchi, “closing volume” speeds up, that is, the air flow closes particularly rapidly in small airways due to the decrease in lung-supporting elastic fibers. The increase in closing capacity causes a decrease in oxygen saturation because of the decrease in alveolar ventilation for lung perfusion, which is triggered by air flow closure.⁽²¹⁾ Likewise, potential to develop breathing problems increases as age increases because of VC decrease and closure of peripheral bronchi. VC decreases in the elderly are considered to be due to decreases in breathing mechanism and to limitations in chest-wall movement. Although the lung-protective structure of the thoracic cavity is essential for normal lung functioning, it more greatly affects normal lung function as age increases due to changes in the structure of the thoracic cavity (i.e., spines, muscles, and ribs).⁽²²⁾ Lombardi, et al.⁽²³⁾ confirmed VC changes in various angles of the thoracic kyphosis in 55 non-smoking women, and reported that FEV₁ and VC decreased as angles increased in pulmonary function test. Muscles also change as age increases, as does the structure. Aging is known to be highly related to decreases in inspiratory and expiratory muscle strength.⁽²⁴⁾ According to previous studies on breathing muscle correlation, maximal inspiratory and expiratory pressures (assessments of maximal inspiratory and expiratory muscle strength) decreased in the elderly. When the mean heights and weights were compared among participants in their 50s to 80s, the mean maximal inspiratory pressure of participants in their 50s was 111 cm H₂O, and that of participants in their 80s was 70 cm H₂O, confirming a significant decrease.⁽²⁵⁾ The prevalence of the age-related decreases in maximal inspiratory and expiratory pressures relate to problems in breathing mechanism and to elderly people’s sarcopenia.⁽²⁶⁾ The results of this study confirmed the overall VC decrease in elderly people as age increases. This age-related VC change is caused by changes in the thoracic structure and breathing muscle weakness in elderly people, as confirmed by previous studies. Therefore, if exercise

programs for the elderly can be developed in accordance with age, they will help to prevent VC decreases in the elderly, and it will lead to successful aging. Additional studies are needed to develop exercise programs that are fit for the elderly in each age group and to confirm VC changes after doing the exercises.

References

1. Yang SA: A study on the relationship among physical function, quality of sleep and quality of life for the elderly in a community. *J Digit Convergence*, 2013, 11: 335–345.
2. Mohamed H: Central Agency for Public Mobilization and Statistics, Cairo, Egypt mido82_mido82@yahoo.com
3. Lee HR: The factors of influence on the quality of life in the aged with chronic disease. Chung -Ang University, Master's Thesis, 2008.
4. Jang YS: Dyspnea in the elderly. *Korean J Clin Geri*, 2005, 6: 81–89.
5. Kim JE: Comparison among death anxiety, depression, powerlessness, and quality of life in each level of dyspnea for elderly patients with chronic obstructive lung disease. Kyunghee University, Master's Thesis, 2014.
6. Guyatt GH, Townsend M, Berman LB, et al.: Quality of life in patients with chronic airflow limitation. *Br J Dis Chest*, 1987, 81: 45–54.
7. Strijbos JH, Postma DS, van Altena R, et al. : A comparison between an outpatient hospital-based pulmonary rehabilitation program and a home-care pulmonary rehabilitation program in patients with COPD. A follow-up of 18 months. *Chest*, 1996, 109: 366–372. [
8. Casaburi R, Porszasz J, Burns MR, et al. : Physiologic benefits of exercise training in rehabilitation of patients with severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 1997, 155: 1541–1551.
9. Anderson KL: The effect of chronic obstructive pulmonary disease on quality of life. *Res Nurs Health*, 1995, 18: 547–556.
10. You EJ: The social economic burden and impact of health related quality of life of COPD. Seoul National University, Master's Thesis, 2014.
11. Shin YH: The effect of walking exercise program on physical function and emotional state in elderly women. Ewha Women's University, Dissertation of Doctorate degree, 1997.
12. Kang CK, Kim HC, Lee MG: Effects of 12 weeks of combined exercise training on cardiopulmonary function and metabolic syndrome risk factors in elderly farmers. *The Korean Journal of Physical Education: Natural Science*, 2008, 47: 377–387.
13. Ries AL, Bauldoff GS, Carlin BW, et al. : Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based clinical practice guidelines. *Chest*, 2007, 131: 4S–42S.
14. Han D, Yoon N, Jeong Y, et al. : Effects of cervical self-stretching on slow vital capacity. *J Phys Ther Sci*, 2015, 27: 2361–2363
15. Qaseem A, Wilt TJ, Weinberger SE, et al. American College of Physicians American College of Chest Physicians American Thoracic Society European Respiratory Society: Diagnosis and management of stable chronic obstructive pulmonary disease: a clinical practice guideline update from the American College of Physicians, American College of Chest Physicians, American Thoracic Society, and European Respiratory Society. *Ann Intern Med*, 2011, 155: 179–191.
16. Hong YS, Ham JE: The study on physiological process and effects of exercise in aging people. *J Korea Soc Aerobic Exerc*, 2001, 5: 133–145.
17. Stanojevic S, Wade A, Stocks J, et al. : Reference ranges for spirometry across all ages: a new approach. *Am J Respir Crit Care Med*, 2008, 177: 253–260.

18. Song SH, Kim HY, Ju JC, et al. : The effect of aging on the pulmonary function of the healthy adults. *Korean J Anesthesiol*, 1990, 78: 1021–1026.
19. Kim DS, Pyo SY, Kim BY: Changes in the pulmonary function according to the increase of healthy adults' age. *Bull Dongnam Health Coll*, 2002, 20: 469–476.
20. Abe T, Suzuki T, Yoshida H, et al. : The relationship between pulmonary function and physical function and mobility in community-dwelling elderly women aged 75 years or older. *J Phys Ther Sci*, 2011, 23: 443–449.
21. Janssens JP: Aging of the respiratory system: impact on pulmonary function tests and adaptation to exertion. *Clin Chest Med*, 2005, 26: 469–484, vi–vii.
22. Bartynski WS, Heller MT, Grahovac SZ, et al. : Severe thoracic kyphosis in the older patient in the absence of vertebral fracture: association of extreme curve with age. *AJNR Am J Neuroradiol*, 2005, 26: 2077–2085.
23. Lombardi I, Jr, Oliveira LM, Mayer AF, et al. : Evaluation of pulmonary function and quality of life in women with osteoporosis. *Osteoporos Int*, 2005, 16: 1247–1253.
24. Freitas FS, Ibiapina CC, Alvim CG, et al. : Relationship between cough strength and functional level in elderly. *Rev Bras Fisioter*, 2010, 14: 470–476.
25. Sachs MC, Enright PL, Hinckley Stukovsky KD, et al. Multi-Ethnic Study of Atherosclerosis Lung Study: Performance of maximum inspiratory pressure tests and maximum inspiratory pressure reference equations for 4 race/ethnic groups. *Respir Care*, 2009, 54: 1321–1328.
26. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al. European Working Group on Sarcopenia in Older People: Sarcopenia: European consensus on definition and diagnosis: report of the European working group on sarcopenia in older people. *Age Ageing*, 2010, 39: 412–423.
27. Lee J, Kang T, Yeo Y, and et al.: The change of lung capacity in elderly women caused by life span. *J Phys Ther Sci*, 29: 658–661, 2017.