

Table 3. Phytoplankton Abundance at Station 3

No.	Phylum	Abundance (ind/Liter)					
		1	2	3	4	5	6
1	Pyrrophyta	401.510	1.078.251	152.168	492.660	146.870	614.040
2	Euglenophyta	36	179	7	0	43	214
3	Charophyta	534.643	77.540	81.567	2.813	119.238	329.440
4	Chlorophyta	36	179	821	828	1.578	1.571
5	Ochrophyta	75.506	6.997	3.113	1.035	8.939	2.570
6	Xanthophyta	71	36	1.028	1.071	1.499	571
7	Cyanophyta	143	143	186	343	1.928	10.915
Abundance		1.011.944	1.163.324	238.890	498.750	280.095	959.321

Based on the table above it can be seen that the highest phytoplankton abundance at station 3 is the Pyrrophyta phylum, while the lowest abundance is in the Rhodophyta phylum. The highest abundance is found in the second observation that is equal to 1.163.324 ind/Liter.

Table 4. Phytoplankton Abundance at Station 4

No.	Phylum	Abundance (ind / Liter)					
		1	2	3	4	5	6
1	Pyrrophyta	619.038	131.604	925.344	485.520	37.128	128.520
2	Euglenophyta	36	0	36	0	800	428
3	Charophyta	205.632	57.577	158.972	217.991	164.934	428.471
4	Chlorophyta	36	29	214	57	64	571
5	Ochrophyta	80.575	1992	7.176	1.542	7.154	21.563
6	Xanthophyta	71	7	3.856	1.999	1.071	2.142
7	Cyanophyta	107	50	2.071	914	3.606	22.919
Abundance		905.495	191.259	1.097.668	708.024	214.757	604.615

Based on the table above, it can be seen that the highest phytoplankton abundance at station 4 is the Pyrrophyta phylum, while the lowest abundance is in the Euglenophyta phylum. The highest abundance is found in the third observation which is equal to 1.097.668 ind/Liter.

The four tables above show that the highest abundance of phytoplankton is at station 2, 6th observation. While the lowest abundance is at station 4, 2nd observation. Where the species with the highest abundance is *Ceratium sp.* amounting to 2.467.584 ind/Liter at the 3rd observation at station 1.

The composition of species between stations is different, where at station 1 there were 50 species, at station 2 found as many as 49 species, at station 3 found as many as 37 species, and at station 4 found as many as 34 species. There are large fluctuations in abundance and type composition at each sampling station.

3. CORRELATION OF PHYTOPLANKTON ABUNDANCE TO DISSOLVED OXYGEN IN JATIGEDE RESERVOIR

The correlation coefficient of linear regression between phytoplankton abundance to dissolved oxygen in the Jatigede Reservoir on stations 1,2,3 and 4 that was obtained based on the results of the analysis were 0.967; 0,867; 0,936; and 0.964. These values show a very strong correlation, because they are in the interval of 0.80-1.00 (Sugiyono 2005).

The correlation of phytoplankton abundance (orange) to dissolved oxygen (blue) at each stations can be seen in the following graphs:




Figure 2. Correlation of Phytoplankton Abundance to Dissolved Oxygen at Station 1




Figure 3. Correlation of Phytoplankton Abundance to Dissolved Oxygen at Station 2




Figure 4. Correlation of Phytoplankton Abundance to Dissolved Oxygen at Station 3




Figure 5. Correlation of Phytoplankton Abundance to Dissolved Oxygen at Station 4

Based on the graphs above, it can be seen that the changes are always in line between the abundance of phytoplankton and dissolved oxygen concentration. This proves that abundant phytoplankton in Jatigede Reservoir caused photosynthesis to produce dissolved oxygen into the waters. The highest abundance of phytoplankton is at station 2, 6th observation. While the lowest abundance is at station 4, 2nd observation. Where the species with the highest abundance are *Ceratium* sp. amounting to 2.467.584 ind/Liter at the 3rd observation at station 1.

The large difference in abundance between each observation is caused by differences in water conditions at different times at the time of sampling. According to Sachlan (1982) the abundance of phytoplankton is influenced by water conditions where the concentration of nutrients such as nitrate and phosphate affects the growth of phytoplankton. The intensity of the light that enters the waters also has a great influence in relation to the photothesis process carried out by phytoplankton. The large difference in abundance at each time of observation is caused by differences in water conditions at different times. At station 4 the

second observation of the large abundance of phytoplankton is caused by rain on the night before sampling in the morning, causing an increase in the volume of the reservoir water resulting in dilution.

While the large differences in abundance in each species are caused by the ability of each species to utilize existing nutrients to carry out photosynthesis. The species with the highest abundance at each station is *Peridinium sp.* and *Ceratium sp.* *Peridinium sp.* and *Ceratium sp.* is a phytoplankton species from phylrophyta phylum belonging to the class of Dinophyceae. Dinophyceae in fresh water are generally non-toxic and harmless as Dinophyceae in seawater is toxic and has a negative effect on aquatic systems. But if the amount is excessive, a bloom can occur that can cause toxic properties for other planktonic organisms (allelopathy). *Peridinium sp.* and *Ceratium sp.* has the ability to prevent other phytoplankton from growing with high biomass, thereby reducing nutritional competition. This causes *Peridinium sp.* and *Ceratium sp.* in the community freshwater phytoplankton is a common type and can dominate abundance in lakes with tropical climates (Rengefors and Legrand 2001).

Strong correlation is caused by the condition of the waters that fluctuate significantly. At station 1 nutrient input occurs continuously from Cimanuk river and surrounding settlement. The number of phytoplankton species obtained was as many as 50 species, which were the most compared to other stations. Strong correlation at station 2 which is the focus of capture fisheries activities in Jatigede Reservoir where this indicates that phytoplankton as abundant natural food and fluctuate with the cycle of fish feeding in the station.

3. CONCLUSION

Based on the results, the highest concentration of dissolved oxygen Jatigede Reservoir was 7.3 mg/L. The most abundant phytoplankton species is *Peridinium sp.* and *Ceratium sp.* at each station. The abundance of phytoplankton is strongly related to the dissolved oxygen concentration, where the increase in phytoplankton abundance is also followed by an increase in dissolved oxygen concentration in the waters. Correlation values ranged from 0.867 to 0.967 which means the correlation is very strong.

References

- [1] Central Hall of the Cimanuk-Cisanggarung River Region. 2018. Leaflet Development of Jatigede Reservoir. KSNVT Construction of Jatigede Reservoir, Sumedang.
- [2] Boyd CE 2015. Water Quality: An Introduction. 2nd ed . Springer, Alabama.
- [3] Effendi H. 2003. Review of Water Quality for Management of Water Resources and Environment. PT. Kanisius, Yogyakarta.
- [4] Fitriani, S. I. 2013. Field Lecture Report: Sumedang Jatigede Dam Project . ITB, Bandung.
- [5] Mulyani, R., Wisnu W. 2012. Spatial Distribution of Harmful Algal Blooms (HABs) Species in the Green Mussel (*Perna viridis*) Cultivation Location of Kamal Muara, North Jakarta. *Akuatika Journal*. 3 (1): 28-39. May 2011
- [6] Nontji, A. 2008. Plankton Laut. LIPI Press, Jakarta.
- [7] Nugroho, AS, Tanjung, SD, Hendarto, B. 2014. Distribution and the Content of Nitrate and Phosphate in the Waters of Rawa Pening Lake. *Bioma*, Vol. 3, No. 1. April 2014
- [8] Nybakken, J. W. 1992 . *Marine Biology : An Ecological Approach*. 3rd Ed . Harper Collins College Publishers, New York.
- [9] Paikia, K., Kalora, JD Distribution of Nitrates and Phosphates to Phytoplankton Abundance in East Yapen Coastal Waters. *Journal of Fisheries and Marine Science* Vol. 1 No. 2 (2017)
- [10] Republic of Indonesia Government Regulation Number 82 of 2001 concerning Water Quality Management and Water Pollution Control of the President of the Republic of Indonesia
- [11] Rengefors, K. and C. Legrand. 2001. Toxicity in *Peridinium aciculiferum* –an Adaptive Strategy to Outcompete Other Winter Phytoplankton. *Journal of Limnology Oceanography* Vol . 46 No. 8: 1990-1997. American Society of Limnology and Oceanography, Inc.
- [12] Sachlan, M. 1982. *Planktonology*. Semarang: Faculty of Animal Husbandry and Fisheries Diponegoro University.
- [13] Sugiyono. 2005. *Quantitative and Qualitative Research Methods for R & D*. Bandung, Alfabeta