



Impact of the African rice midge (*Orseolia oryzivora*) on the yield of rice varieties grown in Baguinéda (Mali)

Fatoumata MAÏGA^{1*}, Ousmane DIARRA^{2,3}, Cheick TÉKÉTE¹, Mah FANE³, Sory SISSOKO¹, Boubacar Madio dit Aladiogo MAIGA⁴, Salimatou SAMAKE^{2,3}, Mohamed Lamine TÉKÉTE⁵, Sognan DAO³, Mamadou Oumar DIAWARA¹, Bernadio SODIO¹.

¹Department of Biology, Faculty of Sciences and Technologies (FST), University of Technical Sciences and Technology of Bamako (USTTB), B.P. 3206 Bamako, Mali.

² Institute of Applied Sciences (ISA) / University campus of KABALA phase II Tel: 20 70 77 43/ 20 70 77 46 /Website: www.isamali.org, Mali phone: (223) 20 22 44 21 Email: contact@isamali.org.

³Research Laboratory in Microbiology and Microbial Biotechnology (LaboREM-Biotech), Faculty of Sciences and Techniques, University of Sciences, Techniques and Technologies of Bamako, Mali BPE: 3206.

⁴ Central Veterinary Laboratory (LCV) 8 Km Road of Koulikoro BP 2295 Bamako, Mali Phone: (223) 20-24-33-44/20-24-23-04/20-24-23-05 Fax: (223) 20- 24 -98-09 Email: mali@abovetmali.org

⁵Institute of Rural Economy (IER), BP: 258, Street Mohamed V, Bamako, Mali.

* Corresponding author: fatimhamadoun@gmail.com

ABSTRACT:

Rice growing occupies a key place in agriculture and the search for food security in Mali. However, in rice production, damage caused by the African rice midge (*Orseolia oryzivora*) is a major constraint and therefore an economically important insect. The African rice midge is a major pest in the South Sudanian zone where development conditions are favorable for it. This study aims to document the effect of the insect on the yield of IER irrigated rice varieties in Mali. The study was carried out in the IER irrigated area of Baguineda during the agricultural seasons of 2016, 2017 and 2018. It made it possible to harvest the rice yield from a square of (25 m²) in the 50 study plots and the counting the number of tillers, panicles, full spikelets, empty spikelets, galls, and grains per tuft, at the time of heading at the level of a yield square (1 m²) placed within 50 study plots. All varieties of rice were susceptible to attack by the insect. An analysis of variance carried out from the data revealed a significant difference in the yield of the different rice varieties (P < 0.05). Average yields of rice varieties were very low and varied between 2.8 t/ha for Seberang to 0.27 t/ha for BW, with a considerable yield loss of 44% in 2016 compared to the other two 2017 and 2018 agricultural seasons in the area.

Keywords: *Orseolia oryzivora*, midge, rice, yield, Baguineda

I.INTRODUCTION:

Rice is the world's leading cereal grown for human consumption. Its global production in 2017 amounted to 756.7 million tons and represents 20% of global food energy needs (FAO, 2017). It is the second cereal after corn in terms of tonnage produced (FAO, 2017). In Mali, the agricultural sector employs nearly 90% of the population. Potential land for rice cultivation is estimated at 606,000 ha for lowland and floodplain rice cultivation and over 1,000,000 ha for IER irrigated rice cultivation (Hamadoun, 2015). Rice growing occupies a key place in agriculture and the search for food security in Mali. National annual rice production reached 2,211,920 tons of paddies in 2014 (Minister of Rural Development, 2017). This production is expected to increase due to the growth in demand for food in Mali and the sub-region. This speculation plays an essential role in national poverty reduction strategies. Mali has been committed since 2008 to the implementation of the West Africa Agricultural Productivity Program (WAAPP) to address these constraints (Hamadoun, 2015). The notorious lack of rice production is partly explained by multiple abiotic (drought, decline in soil fertility) and biotic (arthropods, diseases, weeds) constraints. Among the biotic constraints, insect pests occupy an essential place. In rice production, damage caused by *Orseolia oryzivora* Harris and Gagne (1982) (Diptera: Cecidomyiidae) constitutes a major constraint and therefore an economically important insect. The insect is rife in several African countries Nwilene et al., (2011b). In Mali, it is very present in the region of Sikasso, Koulikoro and Segou in the lowlands and IER irrigated areas Hamadoun, (1996). The damage is caused by the larvae that feed inside the rice heights, which subsequently thicken and take the shape of an onion leaf or onion tube. Infestation levels vary from 30% in the lowlands to 80% in the Irrigated perimeters, particularly according Hamadoun *et al.*, (1998). In Mali, the distribution of midge indicates very contrasting ecological areas whose characteristics are not sufficiently known to explain the large variations recorded Hamadoun et al., (2008). The perimeters of San and Baguineda experienced upsurges of the pest with production losses of around 55% Hamadoun et al. (1999) and Baguineda, considered a midge niche in Mali.

II. METHODOLOGY:

Study site:

The studies were conducted at the level of the Office of Irrigated Perimeters of Baguineda (OPIB).




Fig 1: Presentation of the Baguineda Irrigated Perimeter Office (Diawara, 2010)

Type and period of study:

The study was longitudinal on 50 study plots, located from the start of sector 1 to the end of sector 4 of the Baguineda IER irrigated Perimeter Office (OPIB) for the assessment of damage, identification of the 'insect'. It spanned the period from January 2016 to December 2018.

Biological material:

Biological materials are rice galls, lavas, nymphs and adults of *O. oryzivora*.

Sampling of study plots:

The number of different operators was chosen randomly from an exhaustive list, made available to us by the OPIB management. It is from this list that the survey units were drawn. A sample of 50 farmers for the study was obtained from a previous study, which had the theme: "Development of a Spatial repository for monitoring rice pests in the Irrigated of Baguineda (PIB)" (Diawara, 2010).




Fig 2: Sampling map of the 50 study plots, distributed between the three sectors of the Bagueda Irrigated Perimeter.

To determine yields, on the one hand it was done by harvesting rice plants at the level of a yield square (25 m²). The harvested rice plants were threshed and weighed. On the other hand it was based on counting the number of tillers per tuft, panicle per tuft, full spikelet per tuft, empty spikelet per tuft, gall per tuft, and grain per tuft, at the time of heading, by harvesting at the level of a yield square (1 m²) placed on each of the four sides as well as in the middle of the 50 study plots.

STATISTICAL ANALYSIS:

Data were entered using Excel 2013 software. Statistix.8.0 statistical analysis software, and R x64 3.1.2., used for analyzes of variance in order to determine the significance of the averages recorded between parameters according to the different periods (day, month and year) per plot in the same study area. A statistical test (Tukey or Newman Keuls) was carried out to evaluate the correlations between the different results and with the cultural practices of each producer at the 5% significance level. The results were presented in the form of a table or a graph.

III. RESULTS:

Analysis of variance between varieties and agronomic yield parameters:

The results of the variation in the number of galls and the yield of the different rice varieties for the 2016 rainy season represented in (Table I), show a significant difference ($p < 0.05$) recorded on the average number of spikelets full/tuft/m²: the lowest average (8.662) is recorded on the BW variety and the highest (110.205) on the Seberang variety. The average number of grains/tuft/m² is highly significant, the lowest (50,480) BW and the highest (988,122) with SEBERANG. On the other hand, no significant difference for the other parameters was recorded.



TABLE I: Analysis of variance (ANOVA) between the varieties and agronomic parameters observed in the 2016 rainy season

Varieties	Agronomic parameters						Weight (t)/ha
	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/ m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	
Adny	18,682	3,273	10,081	2,562	20,421 b	216,960 b	1,188
Seberang	23,529	6,995	12,782	5,724	110,205 a	988,122 a	2,800
IER wassa	17,257	3,035	6,855	2,901	10,303 b	95,848 b	1,159
BG	17,876	2,995	4,807	2,473	14,908 b	173,239 b	0,877
BW	14,241	3,025	7,604	2,500	8,662 b	50,480 b	0,268
CV	34,2	49,9	42,6	65,8	112,2	81,1	101,3
Probability	0.859	0.330	0.187	0.597	0.029	0.007	0.626
ES	6,25	1,677	3,659	1,832	24,14	175,4	1,184
Significance	NS	NS	NS	NS	S	HS	NS

CV= Coefficient of variation, ES= Standard Error NS= not significant S= significant, HS= highly significant a,b, etc. : mean values followed by the same letters in a column are not significantly different at the threshold ≤ 0.050 .

In the 2017 off-season, the analysis of variance (ANOVA) did not reveal any significant difference between the varieties with the agronomic parameters observed. On the other hand, we noted arithmetic differences between the values (Table II).

TABLE II: Analysis of variance (ANOVA) between varieties and some parameters observed among producers in the 2017 off-season.

Varieties	Agronomic parameters						Weight (t)/ha
	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/ m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/ m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	
Adny	13,646	12,802	0,000	120,853	5,320	1495,972	5,000
IER wassa	11,942	11,137	0,017	95,582	3,357	1186,314	4,416
BG	9,496	8,273	0,000	79,063	6,197	880,394	4,640
BW	10,802	10,065	0,000	74,674	3,292	895,276	4,200
<i>CV</i>	18 ,1	17,6	54 ,9	56,5	54,9	53,5	20,4
<i>Probability</i>	0,415	0,292	0,757	0 ,839	0,436	0,767	0,822
<i>ES</i>	2,14	1,94	2,09	53,8	2,09	625,6	2,29
<i>Signifiance</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>

CV= Coefficient of variation, *ES*= Standard Error *NS*= not significant *S*= significant, *HS*= highly significant *a,b, etc.* : mean values followed by the same letters in a column are not significantly different at the threshold ≤ 0.050 .

In 2017 in the rainy season, as in the 2017 off-season, the varieties behaved in the same way for all the parameters observed; the analysis of variance did not reveal any significant difference. On the other hand, we noted arithmetic differences between the values (Table III).

TABLE III: Analysis of variance (ANOVA) between varieties and some parameters observed among producers in the 2017 rainy season

Varieties	Agronomic parameters						Weight (t)/ha
	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/ m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	
Adny	18,228	14,753	0,626	3,533	108,201	1554,107	4,162
Seberang	30,362	17,142	0,946	3,680	143,922	2166,364	3,400
IER wassa	15,367	10,779	1,745	0,972	11,830	248,115	4,800
BG	18,536	16,872	0,616	3,835	156,745	1909,816	3,400
BW	14,642	13,599	0,079	5,499	142,895	1858,163	4,000
Kogoni	20,322	15,441	0,867	2,135	149,958	1556,103	4,000
CV	34,9	19,4	94,7	58,6	52,6	53,1	14,7
Probability	0,267	0,504	0,531	0,532	0,599	0,575	0,257
ES	6,63	2,89	0,641	66,5	1,84	843,3	0,596
Significance	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>

CV= Coefficient of variation, *ES*= Standard Error *NS*= not significant *S*= significant, *HS*= highly significant *a,b, etc.* : mean values followed by the same letters in a column are not significantly different at the threshold ≤ 0.050 .

In the 2018 off-season, the analysis of variance (ANOVA) reveals a significant difference ($p > 0.05$) between the varieties for yield per hectare. The Adny variety has the highest value (5.732T/ha). On the other hand, no significant difference was observed between the varieties for the other parameters (Table IV).

TABLE IV: Analysis of variance (ANOVA) between varieties and some parameters observed among producers in the 2018 off-season.

Varieties	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/ m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	Weight (t)/ha
Adny	16,66	13,67	0,00	167,67	2,33	1882	5,732 a
IER wassa	12,14	11	0,00	133,57	0,43	1597	4,688 b
BG	12	11	0,00	138	0,00	1660	4,800 b
Nenekala	12	11	0,00	142	1,60	1682	5,200 ab
CV	17,26	13,09	-	10,91	180,17	22,42	5,61
Probability	0,0951	0,1556	-	0,0797	0,1927	0,8447	0,0048
ES	0,818	0,484	-	5,874	0,417	99,132	0,358
Significance	NS	NS	NS	NS	NS	NS	S

CV= Coefficient of Variation, ES= Error Standard NS= Not Significant, S= Significant

In the 2018 rainy season, a statistical difference of $p > 0.05$ was observed between the varieties for the average number of spikes/tuft/m². On the other hand, no significant difference was observed between the varieties for the other parameters (Table V).

TABLE V: Analysis of variance (ANOVA) between varieties and some parameters observed among producers in the 2018 rainy season

Varieties	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	Weight (t)/ha
Adny	19,96	16,6	0,28	143,54 a	1,60	1930,3	8,820
Seberang	17,35	12,96	0,41	116,11 a	1,80	1568	8,812
IER wassa	16,25	13,01	0,37	75,08 b	1,10	1310,8	10,616
BG	19,56	13,51	0,062	80,70 b	1,00	117,5	9,468
CV	31,45	22,53	141,32	35,74	77,82	38,57	20,29
Probability	0,6056	0,1919	0,9630	0,0117	0,6893	0,1968	0,3348
ES	5,31	2,91	0,55	48,5	0,99	597,2	4,43
Significance	NS	NS	NS	S	NS	NS	NS

CV= Coefficient of variation, ES= Standard Error NS= Not Significant S= Significant, HS= Highly Significant a,b, etc. : mean values followed by the same letters in a column are not significantly different at the threshold ≤ 0.050 .

During the three rainy seasons of the study, no significant difference with $p < 0.05$ on the agronomic parameters was recorded (Table VI). The results showed that all varieties were susceptible to galling with a non-significant difference between them for the average number of galls/tuft/m².

TABLE VI: Analysis of variance (ANOVA) between varieties and some parameters observed in the rainy season during the three years (2016, 2017 and 2018)

Varieties	Agronomic parameters						Weight (t)/ha
	Average number of tiller/tufts/ m ²	Average number of panicles/tufts/m ²	Average number of galls/clump/ m ²	Average number of ears/tufts/ m ²	Average number of spikes/tuft/ m ²	Average number of grain/tuft/ m ²	
Seberang	22,461	14,040	2,143	3,040	152,261	1970,428	5,971
Adny	18,739	12,773	2,508	2,559	90,891	1285,479	5,451
Kogoni	20,322	15,441	0,867	2,135	149,958	1556,103	4,000
BG	19,047	10,951	2,052	2,364	96,354	1214,088	4,409
IER (wassa)	16,911	9,719	2,716	1,854	66,916	1001,156	6,331
BW	14,442	8,312	3,841	4,000	75,779	954,321	2,134
<i>CV</i>	32,6	45,3	163,5	70,9	72,7	69,4	61,5
<i>Probability</i>	6,109	5,520	4,039	1,782	68,31	898,8	3,330
<i>ES</i>	0,451	0,445	0,990	0,601	0,206	0,405	0,664
<i>Significance</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>

CV= Coefficient of variation, *ES*= Standard Error *NS*= Not Significant




Fig 3: Evolution of the yield of varieties in the fields in the rainy season 2016, 2017 and 2018. This figure illustrates the evolution of rice yield in the irrigated plots of OPIB in the commune of Baguineda. Indeed, the figure indicates a considerable increase in yields of different varieties and a drop in the number of galls. On average, the lowest yields of the varieties were observed during the 2016 rainy season which recorded the highest number of galls, while the best were recorded in 2018. Interestingly, the IER Wasssa variety had the best performance in all three environments. Its yield increases from 1159 Kg/ha (2016) to 10,033 Kg/ha in 2018. The low yields of rice varieties from the 2016 season can be explained by the strong midge attack during this year. We can therefore say that all the varieties studied are susceptible to midge.




Fig 4: Evolution of the yield of varieties in the fields in the 2017 and 2018 off-season.

This figure shows an evolution in the yield of the varieties during the 2017 and 2018 off-seasons. The analysis of variance does not show a significant difference between the varieties during the off-season. On average all varieties had yields greater than 4t/ha. These results could be explained by strong sunshine with a virtual absence of African rice midge in the study plots.




Fig 5: Genotype Main Effects and Genotype \times Environment Interaction Effects (GGE) for Yield for Yield (in the five environments) in the rainy season 2016, 2017 and 2018 and in the off-season 2017 and 2018.

The 2017 and 2018 rainy seasons form a mega-environment, indicating that these two conditions have similar environmental effects on the yield of varieties with a G \times E interaction (Figure 5).

Superiority performance calculates the cultivar superiority measure of Lin and Binns (1988). For each genotype, it is the sum of the squares of the difference between its mean in each environment and the mean of the best genotype, divided by twice the number of environments. The varieties with the smallest superiority values tend to be more stable and closer to the best genotype in each environment Adny followed by IER (Wassa) (Table VII). Wricke ecovalence produces the Wricke (1962) ecovalence stability coefficient. This is the contribution of each genotype to the sum of squares genotypes per environment, in an unweight analysis of genotypes per environment means. A low value indicates that the genotype responds consistently to changes in the environment. Thus the Nenekala variety followed by BG is the most favorable to environmental change while Seberang is little affected by the improvement of environmental conditions (T).

TABLE VII: Environmental genotype in rainy season (2016, 2017 and 2018) and off-season (2017 and 2018)

Genotype	Superiority performancy	Ecovalency stability coefficient
Adny	366019	720338
IER (wassa)	500267	961100
Nenekala	503333	248895
Kogoni	605392	2234817
BW	834264	1872386
BG	953820	453013
Seberang	4020574	19643303




Fig 6: Correlation between the 5 environments for the grain yield of the varieties

This matrix visualizes the correlations with different colors; red colors indicate a positive correlation between environments and blue colors indicate a negative correlation between environments. There is a strong correlation between the three rainy seasons $r > 0.7$, so the variety orders do not change much during the three rainy seasons. A weak negative correlation between the rainy seasons of 2017, 2018 and the off-seasons. So the performance of the varieties changes between the rainy seasons of 2017; 2018 and the off-seasons.

DISCUSSION:

The heavy onslaught of the 2016 rainy season greatly affected rice yield, with a significant difference recorded between varieties ($P < 0.05$). Average yields were very low and varied between 2.8 t/ha for Seberang to 0.27 t/ha for BW. The loss of yield was considerable in the same year 2016, i.e. 44% compared to the two other agricultural campaigns 2017 and 2018 in the area. These results are similar to those of Williams (1997) in Nigeria who demonstrated that the appearance of a gall at 5 heights 49 – 63 days after late transplanting causes 40% and 60% yield losses respectively. Similarly, Israel and Prakasa Rao (1968) stated that a severe infestation of rice can prolong the pruning stage, delay flowering, followed by uneven maturity and result in a bushy appearance of the plant. WARDA (2000) also reported yield losses in fields with 30% height infestation, suggesting that for every 1% additional infestation, a farmer can expect a 2-3% yield loss. . As consequences, there would be a drop in photosynthetic activity, therefore a drop in the synthesis of carbohydrates and also the formation of empty grains. A significant difference was obtained between the seasonal action and the population damage of the rice midge. Similarly, a significant difference is observed between different stages of rice and crop development and rice midge population damage. Nacro researchers and collaborators claimed in 1996 that every 1% increase in gall corresponds to 2% loss in yield.

The environmental genotypes of the varieties (Adny, IER wassa, Nènèkala, Kogoni, BW, and BG) in the rainy season show the calculated superiority performance of the Adny and IER (wassa) varieties the measure of superiority of the cultivar of Lin & Binns (1988) as the most stable and closest to the best genotype in each environment. In the off-season, it is the Nenekala variety which is the most stable and closest to the best genotype in each environment according to the ecovalency stability coefficient of Wricke (1962).

CONCLUSION:

The study revealed that the majority of rice farmers in OPIB use the Adny variety. The results show higher rates of galls on this variety during the rainy seasons, which affects yield with an almost absence of galls on rice in the dry season in the irrigated area of Baguineda. The infestation rate varied depending on the varieties and the growing season.

CONFLICT OF INTEREST STATEMENT:

Authors declare that they have no conflict of interest.

ACKNOWLEDGEMENT:

Fatoumata MAIGA is a doctor at the Faculty of Sciences and Techniques, University of Sciences, Techniques and Technologies of Bamako. His studies and research work are supported by a grant from the West African Agricultural Productivity Program (WAAPP) financed by the World Bank and managed by the National Research Committee (CNRA). We thank the World Bank and the CNRA for the IER financial and management support.

REFERENCES:

- ADRAO. 2000 : Une petite mouche à gros problèmes : la cécidomyie africaine des galles du riz. Rapport annuel ADRAO 2000. Points saillants des activités, ADRAO, Bouaké, Côte d'Ivoire, pp 20-26.
- Bonzi, S.M. 1979. Résultats des premières investigations sur les insectes ravageurs du riz en Haute-Volta. Communication au Séminaire de l'ADRAO sur les méthodes intégrées de lutte contre les maladies et les insectes du riz, tenu à Bobo-Dioulasso du 17 au 22 Septembre 1979, INERA, Station de Farako-Bâ. 9 p.
- Dakouo D., Nacro, S. and Sie M. 1988. Evolution saisonnière des infestations de la Cécidomyie du riz *Orseolia oryzivora* H. et G. (Diptera, Cecidomyiidae) dans le Sud-Ouest du Burkina Faso. *Insect Science and its application* 9, 469- 473.
- FAO. 2017. Statistique FAO. www.fao.org. Vu le 22 juin 2018.
- Hamadoun A. 2008. « *Gestion intégrée des nuisibles de riz dans les écosystèmes rizicoles du Mali* », IER- CRRRA Sikasso, 18p.
- Hamadoun, A. 2002. Résistance variétale à la cécidomyie africaine du riz. *In Compte rendu de la seconde revue régionale de la recherche rizicole*. ADRAO-Le centre du riz pour l'Afrique. Réseau Ouest et Centre Africain du Riz (ROCARIZ). Pages 229-235.
- Hamadoun, A. 2015. Les insectes et méthodes de lutte dans les écosystèmes rizicoles du Mali.

- Harris & Gagne.1982. Catalogue of Life : *Orseolia oryzivora*. Description of African rice gall midge, *Orseolia oryzivora* sp. with comparative notes on Asian rice gall midge, *Orseolia oryzae* (Wood Mason) (Diptera: Cecidomyiidae). *Bulletin of Entomological Research*. 72 (3) : 467-472.
- Israel P. and Prakasa Rao P.S. 1968. Influence of gall midge incidence in rice tilling and Johnson D. E. 1997. Les adventices en riziculture en Afrique de l'Ouest, Bouaké. Côte d'Ivoire, 312 p.
- Ministère de l'Agriculture. Plan de campagne agricole consolidé et harmonisé 2017/2018. République du Mali, 2017 ; 36 p.
- Nacro S., Dakouo D. and Heinrichs E. A.1995. Population dynamics, host plant damage and parasitism associated with the African rice gall midge in Southern Burkina Faso. *Insect Science and its Application*. 16 (3/4), 251-257.
- Nacro S. & Nenon J. P. 2009. Comparative study of the Morphology of the ovipositor of *Platysgaster diplosisae* (Hymenoptera: Platygasteridae) and *Aprostocetus procerae* (Hymenoptera: Eulophidae) two parasitoids associated with the African Rice Gall Midge, *Orseolia oryzivora* (Diptera: Cecidomyiidae). *Hindawi Publishing Corporation*, 7 p.
- Nwilene F.E. et al. 2011b. Identification and differentiation of gall midge species from West Africa. *Phytopathology*, **101**, 130-131.
- Nwillene, F. E., Nacro, S., Tamo M., Menozzi P., Heinrichs E. A., Hamadoun A., Dakouo D., Adda c., & Togola A. 2013. Managing insect pests of rice in Africa. In: Realizing Africa's rice promise. AfricaRice. Chap 18; H ISBN 9781845938123. pp 229-240.
- Ukwongwu, M.N. and Joshi R.C.1992. Distribution of African rice gall midge *Orseolia oryzivora* Harris and Gagne and its parasitoids in Nigeria. *Tropical Pest Management*, 38 (3) : 241-244.
- Umeh E., Joshi R.C., Ukwongwu M.N.1993. Les insectes nuisibles du riz au champ en Afrique : Biologie et méthode de lutte. 32 p.
- Williams, C.T., Harris, K.M. and Okhidievbie, O. 1997. Management of the African rice gall midge (*Orseolia oryzivora*) in West Africa. Completion Report for ODA Holdback Project R5619(H). CAB International, Wallingford, UK