



ON  $p$ -GROUPS OF ORDER  $2^{2n+e}$ ,  $e \in \{0, 1\}$  SATISFYING THE  
STRUCTURE GIVEN BY:  $k(G) = p^e + (p^2 - 1)n + (p^2 - 1)(n - 1)t$

ADEBISI , SUNDAY ADESINA

ABSTRACT. Let  $k(G)$  be the number of conjugacy classes of a group  $G$ . Then, there exists a non-negative integer  $t = t(G)$  such that:

$$k(G) = p^e + (p^2 - 1)n + (p^2 - 1)(p - 1)t \quad t \geq 0, \quad n \in \mathbb{Z}^+ \quad \text{and} \quad e \in \{0, 1\}$$

Define: even = positive (+) and odd = negative (-).

Then, the following hold

- (i) If  $n$  and  $t$  are of the same sign then,  $k(G)$  is negative.  
Otherwise,  $k(G)$  is positive.
- (ii) If  $n$  is 0,  $e$  and  $t$  are of the same sign, then  $k(G)$  is negative.  
Otherwise  $k(G)$  is positive.

And the result is generally in harmony with the Sylow's  $C$ ,  $D$  and  $E$  theorems.

### 1. INTRODUCTION

The three main Sylow theorems are the Sylow  $E$  theorem,  $D$  theorem and  $C$  theorem sometimes referring to the "existence", "development" and "conjugate" theorem respectively. Another theorem gives the number of Sylow  $p$ -subgroups of a group for fixed prime  $p$ ; this is sometimes referred to as the Sylow "counting theorem [14].

A Sylow  $p$ -subgroup (sometimes  $p$ -Sylow subgroup) of a group  $G$  is a maximal  $p$ -subgroup of  $G$ , i.e. a subgroup which is a  $p$ -group and which is not a proper subgroup of any other  $p$ -subgroup of  $G$ . The set of all Sylow  $p$ -subgroups for a given prime  $p$  can be denoted by  $Syl_p(G)$ . Here, all members are actually isomorphic to each other and have the largest possible order: if  $|G| = p^n m$ ,  $n > 0$  where  $p$  does not divide  $m$ , then any Sylow  $p$ -subgroup  $W$  has order  $|W| = p^n$ . That is,  $W$  is a  $p$ -group and  $(|G : W|, p) = 1$ . This properties can be exploited to further analyze the structure of  $G$  [17].

Suppose that  $G > \{1\}$  is a  $p$ -group and  $\chi \in Irr(G) =$  the set of irreducible complex characters of  $G$ .

If  $\chi \in Irr(G)$ , then  $\chi(1)^2 \leq |G : Z(G)|$ . Thus,  $\chi(1)^2$  is a divisor of  $|G : Z(G)|$ . Let  $k(G)$  be the class number of the group  $G$ . Then, consider  $Irr(G/Q)$  the subset  $\{\chi \in Irr(G) | Q \leq ker(\chi)\}$  of  $Irr(G)$ . Let  $Lin(G) = \{\chi \in Irr(G) | \chi(1) = 1\}$  be the set of linear characters of  $G$ ; then  $Irr_1(G) = Irr(G) \setminus Lin(G)$  is the set of nonlinear irreducible characters of  $G$ . Clearly,  $Irr_1(G) = Irr(G/G')$ .

**Proposition A** :[20] Let  $G$  be a group of order  $p^m$  and let  $D \leq Z(G) \ni |D| = p$ . Then,  $k(G) \geq p - 1 + k(G | D)$ . We have that  $|Irr(G | D)| \geq p - 1$

*Key words and phrases.*  $p$ -groups, Sylow's, Class number, irreducible character, conjugacy class.

2

ADEBISI , SUNDAY ADESINA

- (i) If  $k(G) = p - 1 + k(G | G)$  , then ,  $Z(G) = D$  and  $\chi(1)^2 = p^{m-1} \forall \chi \in Irr(G | D)$  .Hence ,m is odd .
- (ii) If  $m$  is even , then  $k(G) \geq p^2 - p + k(G | D)$  .

Let  $cl(G) = \{A_1, A_2, \dots, A_t\}$  be the set of all  $G$ -classes . i. e .  $t = k(G)$  ;  $a_i \in A_i$ . Then ,the number of commuting ordered pairs  $(x, y) \in G \times G$  is equal to

$$\sum_{i=1}^t |C_G(a_i)| \cdot |A_i| = |G| \cdot k(G).$$

$\therefore$  the number of non-commuting ordered pair  $(x, y) \in G \times G$  is equal to  $|G|^2 - |G| \cdot k(G)$ . Let  $\alpha_2(G)$  denote the number of ordered non-commuting pairs  $(x, y) \in G \times G \ni G = \langle x, y \rangle$  whence  $\alpha_2(G) = 0$  if  $G$  is either abelian or not generated by 2 elements . Then , Mann has shown that :

$$\sum_{Q \leq G} \alpha_2(Q) = |G|^2 - |G| \cdot k(G) \dots (1)$$

The following theorem is due to P.Hall .

**Proposition  $\alpha$ :**(see[20])

- (a) Suppose that  $G$  is a 2-generator  $p$ -group then ,  
 $\varphi_2(G) = (p^2 - 1)(p^2 - p) \cdot |\Phi(G)|^2$
- (b) If  $G = ES(m, p)$  is an extraspecial group of order  $p^{2m+1}$  , then , the number  $t = t(m, p)$  of nonabelian subgroup of order  $p^3$  in  $G$  is equal to  $\frac{p^{2m}-1}{p^2-1} \cdot p^{2m-2}$

**Proof :**(a)  $\varphi_2(G)$  is equal to the number of (ordered) minimal bases of  $G$  . Now, by assumption ,  $G/\Phi(G)$  is abelian of type  $(p, p)$  . If  $a \in G \setminus \Phi(G)$  ,  $G \setminus \langle a, \Phi(G) \rangle$  , then  $G = \langle a, b \rangle$  . This element  $a$  , can be chosen in  $|G| - |\Phi(G)| = |\Phi(G)|(p^2 - 1)$  ways and  $b$  , after the choice of  $a$  in  $|G| - p|\Phi(G)| = |\Phi(G)|(p^2 - p)$  ways . Applying the combinatorial product rule , the result follows.

(b) Every nonabelian 2-generator subgroup of  $G$  has order  $p^3$  .By using (1) and (a) , taking into cognisance the value  $k(G) = p^{2m} + p - 1$  , we have that  $p^3(p-1)(p^2-1)t = p^{2m+1}(p^{2m+1} - p^{2m} - p + 1) = p^{2m+1}(p^{2m} - 1)(p - 1)$  □

**Theorem  $\beta$ :** Let  $|G| = p^{2n+e}$  ,  $e \in \{0, 1\}$  . Then  $\exists$  a non-negative integer  $t = t(G) \ni k(G) = p^e + (p^2 - 1)[n + (p - 1)t] = p^e + (p^2 - 1)n + (p^2 - 1)(p - 1)t$  .

If  $Q \trianglelefteq G$  then  $t(G/Q) \leq t(G)$  .

**Proof :** (See [12]) By  $\alpha(a)$  , and (1) we get  $|G|^2 \equiv |G| \cdot k(G) \pmod{(p^2 - 1)(p^2 - p)}$  . And so,  $k(G) \equiv |G| \pmod{(p^2 - 1)(p - 1)}$ .

$$\begin{aligned} \text{Now , observe that: } & p^{2n+e} = p^e + (p^{2n} - 1)p^e \\ & = p^e + (p^2 - 1)(p^{2n-2} + p^{2n-4} + \dots + p^2 + 1)(p^e - 1 + 1) \\ & = p^e + (p^2 - 1)(p^{2n-2} + p^{2n-4} + \dots + p^2 + 1)(p^2 - 1) \\ & + (p^2 - 1)(p^{2n-2} + p^{2n-4} + \dots + p^2 + 1) \\ & \equiv p^e + (p^2 - 1)(p^{2n-2} + \dots + p^2 + 1) \equiv p^e + (p^2 - 1)n \pmod{(p^2 - 1)(p - 1)} \\ & \text{(since } p^{2n-2} + \dots + p^2 + 1 \equiv n \pmod{(p - 1)}) \text{ .} \end{aligned}$$

$\therefore k(G) = p^e + (p^2 - 1)n + (p^2 - 1)(p - 1)t = p^e + (p^2 - 1)[n + (p - 1)t]$ , for some  $t = t(G) \in \mathbb{Z}$  . Now, to prove that  $t(G)$  is non negative , use induction on  $|G|$  .By assuming that  $n > 1$  . Let  $Q \leq Z(G)$  be such that  $|Q| = p$  . Then , $k(G) > k(G/Q)$  since  $\bigcap_{\chi \in Irr(G)} Ker \chi(1) = Ker(\rho G) = \{1\}$  , where  $\rho G$  is the regular character of  $G$  . Now ,let  $e = 0$  , then  $|G| = p^{2n}$  and  $|G/Q| = p^{2(n-1)+1}$  , so , by the result above ,  $k(G) = 1 + (p^2 - 1)[n + (p - 1)t(G)]$  and  $k(G/Q) = p + (p^2 - 1)[n - 1 + (p - 1)t(G/Q)]$  . And so,  $0 < k(G) - k(G/Q) = 1 - p + (p^2 - 1)[1 + (t(G) - t(G/Q))(p - 1)]$  .  $\therefore t(G) \geq t(G/Q)$  is obtained , since  $t(G) - t(G/Q) \in \mathbb{Z}$  ,

and  $1 + [t(G) - t(G/Q)](p - 1) > \frac{p-1}{p^2-1} > 0$ .

And so,  $t(G) \geq t(G/Q) \geq 0$ , by induction.

Let  $e = 1$ , then,  $|G| = p^{2n+1}$ ,  $|G/Q| = p^{2n}$ . Then, by the result, we have that :  $k(G) = p + (p^2 - 1)[n + (p - 1)t(G)]$  and  $k(G/Q) = 1 + (p^2 - 1)[n + (p - 1)t(G/Q)]$ . So that  $0 < k(G) - k(G/Q) = p - 1 + (p^2 - 1)(p - 1)[t(G) - t(G/Q)]$ . This must be that  $t(G) - t(G/Q) \geq 0$ . Thus,  $t(G) \geq t(G/Q) \geq 0$   $\square$

**Proposition :** Suppose that  $Q \triangleleft G$  and  $|G : Q| = p$ . Let  $s$  be the number of  $G$ -invariant characters in  $Irr(Q)$ . Then ,

- (i)  $pk(G) = k(Q) + (p^2 - 1)s$ .
- (ii) If, in addition,  $G$  is a  $p$ -group, then  $p - 1$  divides  $s - 1$ .

**Proof :** (i) If  $\tau \in Irr(Q)$  is  $G$ -invariant, then,  $\tau^G = \chi^1 + \dots + \chi^p$  where  $Irr(\tau^G) = \{\tau^1, \dots, \tau^p\}$ ,  $\chi(1) = \tau(1)$ . (This was given by Clifford). Now, let  $\varphi \in Irr(Q)$  be not  $G$ -invariant, then,  $\varphi^G = \chi \in Irr(G)$  and  $\chi_Q = \varphi_1 + \dots + \varphi_p$  is called the Clifford decomposition, where  $\varphi = \varphi_1$ .

Now, since  $\bigcup_{\tau \in Irr(Q)} Irr(\tau^G) = Irr(G)$ ,

$$Irr((\sum_{\chi \in Irr(G)} \chi)Q) = Irr(Q)$$

we thus have that :  $k(G) = |Irr(G)| = sp + \frac{1}{p}(|Irr(Q)| - s) = sp + \frac{1}{p}(k(Q) - s)$ , and the result follows. By theorem ( $\beta$ ), we have the proof of (ii).  $\square$

**Proposition :** (See [12])The number  $s = s(G)$  of nonabelian subgroups of order  $p^3$  is a  $p$ -group  $G$ ,  $|G| = p^m$ ,  $m \geq 5$  is divisible by  $p^2$ .

**Proof :** (See [20])By proposition  $\alpha(a)$  and (1), we have that:  
 $sp^2(p - 1)(p^2 - 1) +$

$$\sum_{Q \leq G, |Q| > p^3} \alpha_2(Q) = p^m \cdot [p^m - k(G)] \dots (ii)$$

If  $Q \leq G$  is a 2-generator subgroup of order  $\geq p^4$ , then  $|\Phi(Q)| \geq p^2$ . Thus,  $\alpha_2(Q)$  is divisible by  $p^5$ , by  $\alpha(a)$ .  
 By (ii), we have the result.  $\square$

It was proved by Mann [in sec. 2 [20]] that

$$k(G) = p^e + (p^2 - 1)[n + (p - 1)t].$$

Now, to exhaust all possibilities,

$$\begin{aligned} k(G) = p^e + t(1 - p) - n(\text{mod } p) &= p^e + t - pt - n(\text{mod } p) \\ &= p^e + t - n(\text{mod } p) \end{aligned}$$

Now, when  $e = 0$ ,  $k(G) = 1 + t - n(\text{mod } p)$

$|G| = 2^{2n}$  implies  $n \geq 1$ ,  $p = 2$ .

If  $n$  is even, (then  $n = 2u$  say) implies that  $k(G) = 1 + t - 2u(\text{mod } 2) = 1 + t(\text{mod } 2)$ .

Further, if  $t$  is even,  $k(G) = 1(\text{mod } 2)$ . But if  $t$  is odd,  $t = 2d + 1$ , say, thus,  $k(G) = 0(\text{mod } 2)$  implies that  $k(G)$  is even.

If  $n$  is odd, (then,  $n = 2w + 1$  say), implies that  $k(G) = 1 + t - 2w - 1 = t(\text{mod } 2)$ .

So, if  $t$  is odd,  $k(G) = 1(\text{mod } 2)$  and even  $t$  implies  $k(G)$  is even.

By Sylow's, if  $B = \{B_i\}$  is a family of all subsets of  $G$ , each of order  $p^k$ .

Then,  $|B| =$  the number of conjugacy classes of order  $p^k$ . Hence, it implies that  $t$  is odd for  $p = 2$  in harmony with the result  $k(G) = 1(\text{mod } 2)$ .

Summarily, we have the generalized result given in tabular form as follows:

$e$	$n$	$t$	$k(G)$	
0	even	even	odd	
0	even	odd	even	
0	odd	odd	odd	
0	odd	even	even	
1	0	even	even	
1	0	odd	odd	
1	1	even	even	
1	1	odd	odd	
1	even	even	odd	
1	odd	odd	odd	

### Bibliography

- [1]: Burnside, W. (1913). On the outer automorphisms of a group. Proc. Lond. Math. Soc. (2) 11, (40-42).
- [2]: Govich, N. et.al. (1998) Groups of prime power order with many conjugacy (1913). On some properties of groups whose orders are powers of primes I.Proc. Lond. Math. Soc.(2)11 225-245,13(6-12).
- [3]: Hall, P. (1933). A contribution to the theory of groups of prime power order. Proc. Lond. Math. Soc. 36 (29-95).
- [4]: Hall, P. (1936). On a Theorem of Frobenius Proc. Lond. Math. Soc. 40 (468-5011).
- [5]: Hall, P. (1940). The classification of prime power groups. J. reine angew. Math. 182 (130-141).
- [6]: Isaacs, I.M. & Moreto, A. (2001). The character degrees and nilpotence class of a  $p$ -group. J. Algebra 238 (827-842).
- [7]: Issacs, I.M. (1972).The number of generators of a linear  $p$ -group,Can. J.Math.24(852-858).
- [8]: Isaacs, I.M. (1976). Character Theory of Finite Groups. Acad. Press, N.Y.
- [9]: Issacs, I.M. (1986). Sets of  $p$ -powers as irreducible character degrees. Proc. Amer. Math. Soc. 96(551-552).
- [10]: Issacs, I.M. (1994). Algebra: A Graduate Course Brooks/Cole.
- [11]: Knoche, H.G. (1951). Über den Frobeniusschen Klassenbegriff in nilpotenten Gruppen. I, II. Math. Z 55 (71-83) Ibid 59 (8-16).
- [12]: Mann, A. (1978). Conjugacy Classes in finite groups. Israel J. Math. 31 (78-84).
- [13]: Mann, A. (1990). On the power structure of some  $p$ -groups Circ. Mat. Palermo II, 23, (227-235).
- [14]: Mann, A. (1995). A transfer result for powerful Sylow Subgroups. J. Algebra 178 (299-301).

- [15]: Mann, A. (1999). Minimal Characters of  $p$ -groups. *J. Group Theory* 2 (225-250).
- [16]: Mann, A. (2007). The power structure of  $p$ -groups I. *J. Algebra* 42 (121-135), 318(953-956).
- [17]: Weir, A. (1955). Sylow  $p$ -subgroups of the classical groups over finite fields with characteristic prime to  $p$ . *Proc. Amer. Math. Soc.* 6 (529-533).
- [18]: Weir, A. (1955). The Sylow subgroups of the symmetric groups. *Proc. Amer. Math. Soc.* 6 (534-541).
- [19]: Wielandt, H. (1959). Ein Beweis für die Existenz der Sylowgruppen. *Arch. Math.* 10(401-402) (MR26#504).
- [20]: Yakov Berkovich (2008). *Groups of prime power order vol. 1*. Walter de Gruyter. GmbH & Co. KG, 10785 Berlin, Germany.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LAGOS, NIGERIA  
*E-mail address:* [adesinasunday@yahoo.com](mailto:adesinasunday@yahoo.com)