



PHYTOCHEMICAL SCREENING AND BRINE SHRIMP LETHALITY ASSAY OF BILLY GOAT WEED (*Ageratum conyzoides* Linn.) CRUDE EXTRACT

JOYREM J. ONDRADA*, MAURICE D. JIMENEA, ANA HAZEL D. BARISON, JERAH MAE B. GRANDE,
ANABELLE J. VILLAR, AIRENE J. FERNANDEZ, JASON A. IBAÑEZ, FRANCIS D. PABON

Bachelor of Secondary Education Major in Physical Science
College of Education

CARLOS HILADO MEMORIAL STATE COLLEGE
TALISAY CITY, NEGROS OCCIDENTAL, PHILIPPINES

Corresponding author's e-mail: ondradajoyrem125@gmail.com ; maumaujimenea@gmail.com

Abstract

*The study focused on the phytochemical screening and brine shrimp lethality assay of the Billy goat weed (*Ageratum conyzoides* Linn.). Dried leaves of the *A. conyzoides* were prepared as a sample for phytochemical composition analysis. The ethanolic crude extract of *A. conyzoides* was obtained for phytochemical screening using the test tube method. The same crude extracts were used for the detection of cytotoxic activity using brine shrimp lethality assay (BSLA) against brine shrimp eggs (nauplii). The phytochemicals detected were alkaloids, saponins, glycosides, flavonoids, phenols, terpenes, coumarins, steroids, tannins, cardiac glycosides, and resins. These phytochemicals were expected to possess pharmacological activities, anti-allergic, anti-oxidant, anti-inflammatory, and anti-bacterial properties. The toxicity level was found out to be active with LC_{50} of 56.23 μ g/mL. Different solvents are explored in order to optimize the phytochemical screening and cytotoxic level of Billy goat weed (*Ageratum conyzoides* Linn.).*

Key Words: *Ageratum Conyzoides, Brine Shrimp Lethality Assay, Crude extract, cytotoxic activity, Medicinal plant, Phytochemical Constituent, Phytochemical Screening*

INTRODUCTION

The medicinal plants above and beyond therapeutic agents are also a great foundation of information for a comprehensive variation of chemical constituents which could be developed as drugs with precise selectivity. These were the reservoirs of potentially useful chemical compounds which could serve as a newer leads and clues for modern drug design (Yadav et al., 2014). These statements were supported by Wadood et al. (2013) that the medicinal plants were useful for healing as well as for curing of human diseases because of the presence of phytochemical constituents.

Billy goat weed (*Ageratum conyzoides* Linn.) is an annual herb that grows about 60 cm high. It is found in countries in tropical and sub-tropical regions. *Ageratum* was widely utilized in traditional medicine systems where ever it grows (Taylor, 2013).

Medicinal plants had been discovered out of different species of plants. It has been used not only for source for survival but as well as to cure certain diseases and wounds. It is being utilized by most people, especially those who can't afford to buy synthetic drugs. In Philippines, medicinal plants were considered to be the main source of medicine for curing diseases.

One of the most commonly used terms to determine the chemical constituents is "Phytochemical screening". Generally, phytochemical screening refers to the extraction, screening and identification of the medicinally active substances found in plants. Indeed, synthetic drugs are more expensive than using medicinal plants. Synthetic drugs are unsuitable to those who bore these costs. Moreover, in order to determine the cytotoxic level of medicinal plants "Brine Shrimp Lethality Assay" had been performed. Conventionally, Brine shrimp lethality assay refers to a useful tool for preliminary assessment of toxicity. It has also been recommended for screening pharmacological undertakings in plant extracts.

The purpose of this study was to determine the chemical constituents present in Billy goat weed (*Ageratum conyzoides* Linn.) crude extract and its cytotoxic level using Brine shrimp Lethality Assay. Billy goat weed (*Ageratum conyzoides* Linn.) has active constituents which used to cure certain diseases. The bottom line of this study, medicinal plants such as Billy goat weed were beneficial to human.

This study aimed to chemically characterize the ethanolic crude extracts of Billy goat weed (*Ageratum conyzoides* Linn.) and its cytotoxic activity using Brine Shrimp Lethality Assay. Specifically, this study aimed the to determine the phytochemical constituents present in Billy goat weed (*Ageratum conyzoides* Linn) ethanolic crude extracts using test tube method; and to confirm its cytotoxic activity using Brine Shrimp Lethality Assay.

Materials and methods

Collection and Preparation of Plant Sample

The Billy goat weeds (*Ageratum conyzoides* Linn.) were collected from the Hda. Salome V, Brgy. San Pablo, Manapla Negros Occidental and was authenticated by the Bureau of Plants Industry of the Department of Agriculture located at Brgy. La Granja, Lacarlota City Negros Occidental. Only the leaves were chosen from the entire part of Billy goat weed (*Ageratum conyzoides* Linn.). The leaves undergo air drying. The samples were packed in a net bag so as to avoid the moisture. The collected samples have been analyzed at the Chemistry laboratory at the University of San Agustin, Iloilo City.

Dried Billy goat weed (*Ageratum conyzoides* Linn.) leaves were used in obtaining the extract. About 200 grams of sample was finely cut and grinded using mortar and pestle and immediately plunged into 95% ethyl alcohol to prevent enzymatic hydrolysis. The finely pounded leaves were then properly labeled and stored using Erlenmeyer flask in the refrigerator for about 0°C to 5°C. After 48 hours, the mixture was squeezed using gauze and filter in obtaining the filtrate. The filtrate was collected and subjected under rotary evaporation at the controlled 50° Celsius temperature of the water bath to achieve a syrupy concentrated extracts.

Phytochemical Screening of the Crude Extract

Phytochemical screening of the crude ethanolic extract of Billy goat weed (*Ageratum conyzoides* Linn.) leaves has been carried out using the test tube method described by Aguinaldo et al. (2004) and Tiwari (2011).

Detection of alkaloid - Extracts were dissolved individually in dilute hydrochloric acid and filtered. Filtrates were treated with Mayer's reagent (Potassium Mercuric Iodide). Formation of a yellow colored precipitate indicates the presence of alkaloid.

Detection of glycosides - Extracts have been hydrolyzed with dilute hydrochloric acid and then subjected to test for the presence of glycosides using modified Borntrager's test. Extracts were treated with ferric chloride solution and immerse in boiling water for about five minutes. The mixture was cooled and extracted with equal volumes of benzene. The benzene layer was separated and treated with ammonia solution. Formation of rose-pink color in the ammonical layer indicates the presence of anthranol glycosides.

Detection of saponins - Detection of saponins had been carried out using froth test. Extracts were diluted with distilled water to 20 ml and shaken in a graduated cylinder for 15 minutes. Formation of 1cm layer of

foam indicates the presence of saponins.

Detection of phytosterols - Detection of phytosterols had been carried using Salkowski's test. Extracts were treated with chloroform and filtered. The filtrates were treated with few drops of concentrated sulfuric acid, shaken and allowed to stand. Appearance of golden yellow color indicates the presence of triterpenes. Liebermann Buchard's test: extracts had been treated with chloroform and filtered. The filtrates were treated with few drops of acetic anhydride, boiled and allow cooling. Concentrated sulfuric acid was added. Formation of the brown ring at the junction indicates the presence of phytosterols.

Detection of phenols - Detection of phenols had been carried out using ferric chloride test. Extracts were treated with 3-4 drops of ferric chloride solution. Formation of bluish black color indicates the presence of phenols.

Detection of tannins - Detection of tannins had been carried out using the gelatin test. The extract was added with 1% gelatin solution containing sodium chloride. Formation of white precipitate indicates the presence of tannins.

Detection of flavonoids - Detection of flavonoids had been carried out using the lead acetate test. Extracts were treated with few drops of lead acetate solution. Formation of yellow color precipitate indicates the presence flavonoids.

Detection of diterpenes - Detection of diterpenes had been carried out using the copper acetate test. Extracts were dissolved in water and treated with 3-4 drops of copper acetate solution. Formation of emerald green color indicates the presence of Diterpenes.

Detection of terpenes - To 5 ml of the extract, add 2 ml of chloroform and 3 ml of Sulfuric acid (H_2SO_4) concentration. Formation of a reddish brown ring confirms the presence of terpenes.

Detection of coumarines - In a test tube 0.5 g of the moistened various extracts were taken. The mouth of the tube was covered with filter paper treated with 1 N NaOH solution. Test tube was placed for few minutes in boiling water and then the filter paper was removed and examined under the UV light for yellow fluorescence indicated the presence of coumarins.

Detection of steroids - A 0.5 g of the various solvent extract fraction of each plant was mixed with 2 ml of acetic anhydride followed by 2 ml of sulfuric acid. The color changed from violet to blue or green in some samples indicated the presence of steroids.

Detection of cardiac glycosides - A 5 ml of various solvent extract was mixed with 2 ml of glacial acetic acid containing one drop of ferric chloride ($FeCl_3$) solution, followed by the addition of 1 ml concentrated sul-

phuric acid. Brown ring was formed at the interface which indicated the presence of deoxysugar of cardenolides. A violet ring may appear beneath the brown ring, while in the acetic acid layer, a greenish ring may also form just gradually throughout the layer.

Detection of resins - Acetone-water Test: Extracts were treated with acetone. Small amount of water was added and shaken. Appearance of turbidity indicates the presence of resins.

Brine Shrimp Lethality Assay

Brine shrimp lethality assay had been carried out according to the principles and protocol described by Aguinaldo et al (2004) and Naidu (2013) with slight modification. Brine shrimp (*Artemia salina*) eggs hatched in artificial sea water prepared by dissolving 3.8 g rock salt in 100 ml distilled water. The container was covered with black cartolina, with aeration and illumination on the side of the chamber. After 48 hours of incubation at room temperature the larvae (active *nauplii*) were separated from the shells and are attracted to the brighter side of the hatching chamber. The *nauplii* were collected using micropipette. Ten active *nauplii* used for testing were placed in each vial containing 5 mL of artificial seawater solution. Tests have been done along with control and different concentrations.

Preparation of Artificial Seawater

Artificial seawater was prepared by dissolving 3.8 g of rock salt per 100 mL distilled water.

Preparation of Different Solutions

In a solution A, 50 mg of Billy Goat weed extract was dissolved in 5 ml of methanol. For solution B, 0.5 mL of solution A was diluted in 10 ml methanol. After the solutions were prepared, the researchers pipette 100µL of solution B, 50 µL of solution A and 500µL of solution A into separate vials respectively and were labeled from 1 to 3 with 5 replicates each concentration. Then, a control vial containing 1 mL of methanol was made. The solution in each vial of concentration had undergone air drying at room temperature.

Hatching the Shrimp

Shallow rectangular dish (22cm x 32cm) was filled with artificial sea water; the rectangular dish served as the hatching dish of the brine shrimp. Then, a plastic divider was placed, punched with several 2mm holes in the dish which divide into two unequal compartments. The minute brown shrimp eggs were sprinkled into a larger compartment. The larger compartment was covered with black cartolina to keep away from light; a smaller compartment was left open. The smaller open compartment was then illuminated. The hatched brownish orange *nauplii* were pipette after 48 hours from the illuminated compartment of the dish.

Counting the nauplii

The *nauplii* were pipette and count microscopically in the stem of the pipette, held against a well-lighted background.

Preparation of Food for the Shrimp

A suspension of 3mg of dry yeast (red star) in 3ml artificial seawater was prepared for the food for the *nauplii*.

Concentration of sample vials 1, 2 and 3 and control vial

Each sample vials diluted to 5ml with artificial sea water makes a final concentration of 10,000 and 1000 µg/ml respectively. With a 9 inch pipette, transfer ten nauplii into each sample vial labelled 1, 2 and 3 and control vial prepared. Add artificial sea water to each vial samples and control to make a total volume of 5ml. Add a drop of yeast suspension (3mg/5ml of sea water) as food in each vial. The vials were kept under illumination. The survivors were counted after 6 hours then after 24 hours. The record of the number of deaths after 24 hours was used in determination of the percent deaths for each dose level and for the control vials.

Lethality Concentration Determination

The surviving shrimps were counted using magnifying glass. Survivors were counted after 24 hours and the percentage mortality at each vial and control has been determined using the equation:

$$\% \text{ mortality} = \frac{\text{no. of dead nauplii}}{\text{initial no. of live nauplii}} \times 100$$

Statistical Tool

Probit analysis by Finney as discussed by Kim Vincent (2014) was used in determining the concentration at which lethality to brine shrimp represents 50% (LC50).

Results and discussion

Table 1 show the phytochemical constituents screening using the test tube method. It stated that Billy goat weed crude extract is positive for alkaloids, Saponins, glycosides, flavonoids, phenols, terpenes, coumarins, steroids, tannins, cardiac glycosides, resins. Phytosterols and Diterpenes are not present in Billy goat weed (*Ageratum conyzoides* Linn.). Different reagents were used in the determination of the phytochemical constituents in Billy goat weed (*Ageratum conyzoides* Linn.). They were Mayer's reagent distilled water, ferric chloride, lead acetate, sulfuric acid, acetic anhydride, copper acetate, sodium hydroxide, 1% gelatin solu-

tion, and acetone. This implies that those phytochemical constituents with positive remarks are present in Billy goat weed (*Ageratum conyzoides* Linn.) while those with negative remarks are not present in Billy goat weed (*Ageratum conyzoides* Linn.). The result in table 1 is anchored to the study of (Naz and Bano, 2013) where the value of medicinal plants lies in phytochemical constituents that cause definite pharmacological action on the human body. The plants are the vital source of innumerable number of antimicrobial compounds. Several phytochemical constituents like flavonoids, phenolics and polyphenols, tannins, terpenoids, sesquiterpenes etc., are effective antimicrobial substances against a wide range of microorganisms.

These phytochemical constituents present in Billy goat weed (*Ageratum conyzoides* Linn.) crude extracts have beneficial effects to human. The phytochemical constituents present in Billy goat weed (*Ageratum conyzoides* Linn.) may be utilized for medicinal purposes. In the study conducted by Doughari (2012), alkaloids are known for their pharmacological applications as anesthetics and Central Nervous System (CNS) stimulants, the antibacterial properties, the anticancer agent, the anti-arrhythmic, the pupil dilator, and the addictive stimulants; saponins has biological effects, inhibitory effects on inflammation, and possess antibacterial property according to Abioye et al. (2013); glycosides are neutral in reaction and can be readily hydrolyzed into its components with ferments or mineral acids. Glycosides are classified on the basis of type of sugar component, chemical nature of aglycone or pharmacological action according to Doughari (2012); flavonoids have long been recognized to possess anti-allergic, anti-inflammatory, antiviral, anti-proliferative and anti-carcinogenic activities as well as to affect some aspects of mammalian metabolism. The protective effects of flavonoids in biological systems are ascribed to their capacity to transfer electrons free radicals, chelate metal catalysts, activate antioxidant enzymes, reduce alpha-tocopherol radicals, and inhibit oxidases according to Ukachukwu et al. (2013); phenols possess antimicrobial, antioxidants and cytotoxicity activities according to Sowunmi and Afolayan (2015); terpenes possess anti-microbial, anti-carcinogen, antioxidant, analgesic (painkiller), anti-inflammatory, muscle relaxer, and anti-depressant according to Vogeler (2010); coumarins possess anti-tumor activity according to Lacy and Kennedy (2004); steroids possess therapeutic applications and potent anti-malarial effects according to Adeyemi et al. (2014); tannins has anti-inflammation, anti-cancer, and ethno-pharmacological uses according to Sowunmi and Afolayan (2015); cardiac glycosides possess anti-cancer and anti-tumor according to Montaña et al. (2014); and lastly, the resins according to Amabye (2015) has anti-inflammatory properties.

Table 1: Phytochemical Screening of Billy goat weed (*Ageratum conyzoides* Linn.)

Phytochemical Constituents	Reagents	Positive Results	Remarks
Alkaloids	Mayer's reagent	Yellow colored precipitate	+
Saponins	Distilled water	1 cm layer of foam	+
Glycosides	ferric chloride	Rose pink	+
Flavonoids	Lead acetate	Yellow Colored precipitate	+
Phytosterols	Sulfuric acid	Golden Yellow color	-
	Acetic anhydride	Brown ring	
Phenols	ferric chloride	Bluish black	+
Diterpenes	Copper acetate	Emerald green Color	-
Terpenes	Sulfuric acid	Reddish brown	+
Coumarins	Sodium hydroxide	Yellow Fluorescence	+
Steroids	Acetic anhydride	Violet to blue or green	+
Tannins	1% gelatin solution	White precipitate	+
Cardiac Glycosides	Sulfuric acid	Brown ring	+
Resins	acetone	Turbidity	+

(+) present; (-) absent

Table 2 shows the differences of death percentage of *nauplii* in different concentrations of ethanolic crude extract of Billy goat weed (*Ageratum conyzoides* Linn.). In the first concentration which is 10µg/ml, the death percentage of *nauplii* of 39.58%. The second concentration is 100 µg/ml; it has a death percentage of *nauplii* of 62.50%. Lastly, the third concentration is 1000 µg/ml; it has a death percentage of *nauplii* of 95.83%. Using the probit analysis, the LC₅₀ of ethanolic crude extract of the Billy goat weed (*Ageratum conyzoides* Linn.) was computed to be 56.23 ppm (µg/ml). Cytotoxicity test performed on the ethanolic crude extract of Billy goat weed (*Ageratum conyzoides* Linn.) was the so-called brine shrimp lethality assay (BSLA). This implies the result in table 2 shows that the more concentrated the solution with the ethanolic

crude extract of Billy goat weed (*Ageratum conyzoides* Linn.), the more potent it is.

Meanwhile, Peteros and Uy (2010) stated that the percentage mortality increased with an increase in concentration. It was supported by Del Socorro et al. (2014) that the maximum mortality observed at the highest treated-concentration whereas least mortality observed at the lowest treated-concentration. Based on the criteria set by Olowa and Nuñez (2013), the extract is toxic or active if it has a LC_{50} that is lesser than 1000 $\mu\text{g/ml}$ while it is nontoxic or inactive if the LC_{50} is greater than 1000 $\mu\text{g/ml}$. This goes to show that extract of Billy goat weed (*Ageratum conyzoides* Linn.) is indeed toxic. On the other hand another criterion was set on cytotoxicity using brine shrimp by Maridass (2008), which would require the LC_{50} value to be less than 250 $\mu\text{g/ml}$ in order to be considered toxic or active, so the cytotoxicity level of Billy goat weed (*Ageratum conyzoides* Linn.) based on Maridass criterion is active or toxic. It cannot be denied however that Billy goat weeds (*Ageratum conyzoides* Linn.) as a toxic material that can still be explored further. It should be noted that the computation of the LC_{50} was based on the percent mortality after 24 hours.

Table 1: Percent death and lethal concentration of Billy goat weed (*Ageratum conyzoides* Linn.) after 24 hours

Sample	Percent death after 24 hours			LC_{50} ($\mu\text{g/ml}$)
	10 $\mu\text{g/ml}$	100 $\mu\text{g/ml}$	1000 $\mu\text{g/ml}$	
Billy goat weed (<i>Ageratum conyzoides</i> Linn.) extracts	39.58%	62.50%	95.83%	56.23

Conclusion

It can be concluded that the crude extract contains alkaloids, saponins, glycosides, flavonoids, phenols, terpenes, coumarins, steroids, tannins, cardiac glycosides, resins. The toxicity level of Billy goat weed (*Ageratum conyzoides* Linn.) with LC_{50} of 56.23 $\mu\text{g/ml}$ is relatively active. In the light of the findings of the study, it is recommended that nutraceuticals and pharmacological analysis should likewise be performed. Phytochemical components identified should be isolated and further studied for structure elucidation. Other method such as Thin-layer Chromatography (TLC) should be explored as well in identifying phytochemical constituents. Different solvent should be explored in order to optimize the phytochemical screening and cytotoxic activity of Billy goat weed (*Ageratum conyzoides* Linn.).

Acknowledgment

The authors are thankful to the following: **Mr. Ricky Acanto**, College Professor in Carlos Hilado Memorial State College, who is always there to extend a helping hand with his untiring support and wisdom. **Personnel of College Library of Carlos Hilado Memorial State College**, for their genuine assistance on the books needed for the related literature and other data needed in this study.

References

- Abad, M. D.. (2014). CHEMICAL CONSTITUENTS, ANTIBACTERIAL PROPERTIES, AND CYTOTOXIC ACTIVITIES OF PYCNOPORUS COCCINEUS. *QSU Research Journal*, 3(1). Retrieved from <http://ejournals.ph/form/cite.php?id=11485>
- Abioye, E., Akinpelu, D., Aiyegoro, O., Adegboye, M., Oni, M., & Okoh, A. (2013). Preliminary Phytochemical Screening and Antibacterial Properties of Crude Stem Bark Extracts and Fractions of *Parkia biglobosa* (Jacq.). *Molecules*, 18(7), 8485–8499. <https://doi.org/10.3390/molecules18078485>
- Acanto, R. (2015). *PHYTOCHEMICAL SCREENING AND ANTIBACTERIAL ACTIVITY OF BROWN ALGAE (Sargassum oligocystum) OINTMENT*, 2(1). https://www.academia.edu/20808101/PHYTOCHEMICAL_SCREENING_AND_ANTIBACTERIAL_ACTIVITY_OF_BROWN_ALGAE_Sargassum_oligocystum_OINTMENT
- Acanto, R. B. (2016). *Phytochemical screening and cytotoxic activity of Nypa fruticans fruit crude extracts*. Unpublished researched. CHMSC. Philippines.
- Adeyemi, T. O. A., Ogboru, R. O., Idowu, O. D., Owoeye, E. A., & Isese, M. O. (2014). Phytochemical Screening and Health Potentials of *Morinda Lucida* Benth. *International Journal of Innovation and Scientific Research*, 11(2), 515–519. <http://www.ijisr.issr-journals.org/abstract.php?article=IJISR-14-232-03>
- Amabye, T. G. (2016). Evaluation of Phytochemical, Chemical Composition, Antioxidant and Antimicrobial Screening Parameters of *Rhamnus prinoides* (Gesho) Available in the Market of Mekelle, Tigray, Ethiopia. *Natural Products Chemistry & Research*, 04(01), 1–5. <https://doi.org/10.4172/2329-6836.1000198>
- Balangcod, T. D., & Balangcod, K. D. (2015). Ethnomedicinal Plants in Bayabas, Sablan, Benguet Province, Luzon, Philippines. *Electronic Journal of Biology*, 11(3). <https://ejbio.imedpub.com/ethnomedicinal-plants-in-bayabas-sablan-benguet-province-luzon-philippines.php?aid=7365>
- Del Socorro, M. M. L., Bendoy, C. P., & Dacayana, C. M. L. (2014). Cytotoxic Effects of Betel Vine, Piper betle Linn. Leaf Extracts Using *Artemia salina* Leach (Brine Shrimp Lethality Assay). *Journal of Multidisciplinary Studies*, 3(1), 100–111. <https://doi.org/10.7828/jmds.v3i1.629>
- Dhandapani, R., & Sabna, B. (2008). Phytochemical constituents of some Indian medicinal plants. *Ancient science of life*, 27(4), 1–8. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3330865/>
- Doctor, T. R., (2014). Phytochemical screening of selected indigenous medicinal plants of Tubay, Benguet Province, Cordillera Administrative Region, Philippines. *International Journal of Scientific and Research Publications*, 4(4). Retrieved from <http://www.ijisr.org/research-paper-0414/ijisr-p2826.pdf>
- James Hamuel Doughari (March 21st 2012). *Phytochemicals: Extraction Methods, Basic Structures and Mode of Action as Potential Chemotherapeutic Agents*, *Phytochemicals - A Global Perspective of Their Role in*

Nutrition and Health, Venketeshwer Rao, IntechOpen, DOI: 10.5772/26052. Available from: <https://www.intechopen.com/books/phytochemicals-a-global-perspective-of-their-role-in-nutrition-and-health/phytochemicals-extraction-methods-basic-structures-and-mode-of-action-as-potential-chemotherapeutic->

- Ekpenyong, C. E., Akpan, E. E., & Daniel, N. E. (2014). Phytochemical Constituents, Therapeutic Applications and Toxicological Profile of *Cymbopogon citratus* Stapf (DC) Leaf Extract. *Journal of Pharmacognosy and Phytochemistry*, 3(1), 133–141. https://www.phytojournal.com/vol3Issue1/Issue_may_2014/32.1.pdf
- R. Hamidi, M., Jovanova, B., & Kadifkova Panovska, T. (2014). Toxicological evaluation of the plant products using Brine Shrimp (*Artemia salina* L.) model. *Macedonian Pharmaceutical Bulletin*, 60(01), 9–18. <https://doi.org/10.33320/maced.pharm.bull.2014.60.01.002>
- AJOY, G., & PADMA, C. (2013). BRINE SHRIMP CYTOTOXIC ACTIVITY OF 50% ALCOHOLIC EXTRACT OF CROTON BONPLANDIANUM BAILL. *Asian Journal of Pharmaceutical and Clinical Research*, 6(7), 40-41. Retrieved from <https://innovareacademics.in/journals/index.php/ajpcr/article/view/82>
- Ijaiya, I. S., Arzika, S., & Abdulkadir, M. (2014). Extraction and Phytochemical Screening of the Root and Leave of *Annona Senegalesis* (Wild Custad Apple). *Academic Journal of Interdisciplinary Studies*, 3(7), 9–15. <https://doi.org/10.5901/ajis.2014.v3n7p9>
- Lacy, A., & O’Kennedy, R. (2004). Studies on Coumarins and Coumarin-Related Compounds to Determine their Therapeutic Role in the Treatment of Cancer. *Current Pharmaceutical Design*, 10(30), 3797–3811. <https://doi.org/10.2174/1381612043382693>
- Lanzotti V. (2013) Diterpenes for Therapeutic Use. In: Ramawat K., Mérillon JM. (eds) *Natural Products*. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-22144-6_192
- Maridass, M. (2008) "Evaluation of Brine Shrimp Lethality of *Cinnamomum* Species," *Ethnobotanical Leaflets: Vol. 2008 : Iss. 1 , Article 106*. Available at: <https://opensiuc.lib.siu.edu/ebf/vol2008/iss1/106>
- Calderón-Montaño, J. M., Burgos-Morón, E., Orta, M. L., Maldonado-Navas, D., García-Domínguez, I., & López-Lázaro, M. (2014). Evaluating the Cancer Therapeutic Potential of Cardiac Glycosides. *BioMed Research International*, 2014, 1–9. <https://doi.org/10.1155/2014/794930>
- Naidu, J. R., Ismail, R., & Sasidharan, S. (2014). Acute Oral Toxicity and Brine Shrimp Lethality of Methanol Extract of *Mentha Spicata* L (Lamiaceae). *Tropical Journal of Pharmaceutical Research*, 13(1), 101. <https://doi.org/10.4314/tjpr.v13i1.15>
- Nuhu, A. A. (2014). Bioactive Micronutrients in Coffee: Recent Analytical Approaches for Characterization and Quantification. *ISRN Nutrition*, 2014, 1–13. <https://doi.org/10.1155/2014/384230>
- Ongpoy, R. C. Jr., (2015). Phytochemical screening and antimicrobial study of the different leaf extracts of *Alocasia sanderiana* bull., an endemic philippine plant. *International journal of scientific & technology research volume 4, issue 12*. Retrieved from <http://www.ijstr.org/final-print/dec2015/Phytochemical-Screening-And-Antimicrobial-Study-Of-The-Different-Leaf-Extracts-Of-Alocasia-Sanderiana-Bull-An-Endemic-Philippine-Plant.pdf>
- Olowa, L. F. and Nuñez, O. M. (2013). Brine shrimp lethality assay of ethanolic extract of three selected species of medicinal plants from Iligan City, Philippines. *International Research Journal of Biological Sciences* 2(11), 74-77. Retrieved from <http://www.isca.in/IJBS/Archive/v2/i11/12.ISCA-IRJBS-2013-177.pdf>

- Peteros, N.P. and Uy, M.M. (2010). Antioxidant and cytotoxic activities and phytochemical screening of four Philippine medicinal plants. *Journal of Medicinal Plants Research*. Retrieved from [http://www.scirp.org/\(S\(lz5mqp453edsnp55rrgict55\)\)/reference/ReferencesPapers.aspx?ReferenceID=174863](http://www.scirp.org/(S(lz5mqp453edsnp55rrgict55))/reference/ReferencesPapers.aspx?ReferenceID=174863)
- Sowunmi, L. I., & Afolayan, A. J. (2015). Phytochemical constituents and antioxidant properties of acetone extract of *Cleome gynandra* (L.) growing in the Eastern Cape, South Africa. *African Journal of Traditional, Complementary and Alternative Medicines*, 12(3), 1. <https://doi.org/10.4314/ajtcam.v12i3.1>
- Wadood, A. (2013). Phytochemical Analysis of Medicinal Plants Occurring in Local Area of Mardan. *Biochemistry & Analytical Biochemistry*, 02(04), 1–4. <https://doi.org/10.4172/2161-1009.1000144>
- YADAV, M., CHATTERJI, S., GUPTA S.K., & WATAL G.. (2014). PRELIMINARY PHYTOCHEMICAL SCREENING OF SIX MEDICINAL PLANTS USED IN TRADITIONAL MEDICINE. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(5), 539–542. <https://innovareacademics.in/journal/ijpps/Vol6Issue5/9439.pdf>

