



















texture, and taste scores were 6,8; 7,1; 6,6 and 6,7; alternative value 7,00 and priority value 0,28. The test result of crude fiber content is 3,91%.

## References

- [1] Affandi, R. 2017. The Strategy of Resource Utilization Eel Fish (*Anguilla* spp.) in Indonesia. *Indonesian Journal of Ichthyology*, 5 (2): 77-81.
- [2] Agusman, Apriani, S. N. K., and Murdinah. 2014. The Use of Flour Seaweed *Euचेuma cottonii* In the Manufacture of Analog Rice from Flour *Modified Cassava Flour (Mocaf)*. *JPB Fisheries*, 9(1): 1-10.
- [3] Agustini, L. 1996. The standardization Formula Empek-empek Palembang from Fish Cork (*Ophiocephalus striatus Blkr*). *Thesis*. Department of Food Technology and Nutrition, Faculty of Agricultural Technology, Bogor Institute of Agriculture. Bogor.
- [4] Aliani, M. and L. J. Farmer. 2005. Precursors of Chicken Flavor. II. Identification of Key Flavor Precursors Using Sensory Methods. *Journal of Agricultural and Food Chemistry*, 53 (16): 6455-6462.
- [5] Altun, T., T. Nazim, N. Erdal, and S. Yusuf. 2005. Some Growth Parameters on European Eel (*Anguilla anguilla*) Feed with Different Feeds. *Journal of Fisheries & Aquatic Sciences*, 22 (1): 215 – 219.
- [6] Arai, T. 2014. *Do We Protect Freshwater Eels or Do We Drive Them To Extinction?*. SpringerPlus 3, 534.
- [7] DeMan, M. J. 1989. *Principle of Food Chemistry*. The Avi Pub Co. Inc. Ireland.
- [8] The Department of Nutrition. 1999. *Ministry of Health and Institute of Health Singapore*. Singapore.
- [9] Fahmi, M. R., D. D. Solihin, K. Soewardi, L. Pouyau, Z. Shao and P. Berrebi. 2012. A novel semimultiplex PCR assay for identification of tropical eels of the genus *Anguilla* in Indonesia water. *Fish Science*, 79: 185-191.
- [10] Fernando, D. 2007. Physical and Organoleptic Properties of Meat of Broiler Chickens Fed Feed with The Addition of Flour of Bitter Leaf. *Thesis*. Program the Study of Livestock Product Technology Faculty of Animal Husbandry. Institute of Agriculture Bogor. Bogor.
- [11] Hariadi, H. and Y. Rahimah. 2017. The Influence of The Addition of White Oyster Mushroom (*Pleurotus ostreatus*) Against the Nutrient content and the Organoleptic Properties of the Meatballs Tilapia (*Oreochromis niloticus*) Carrots (*Daucus carota* L). *Journal of Science and Food Technology*, 3 (1): 172-177.
- [12] Herdiani, F. 2003. Utilization Seaweed (*Euचेumma cottonii*) To Increase the Levels of Iodine and dietary Fiber on Jam and Dodol. *Thesis*. Institute Bogor Agricultural. Bogor.
- [13] Hsu, S. Y. and Chung, H. Y. 1998. Effects of processing factors on qualities of emulsified meatball. *Journal of Food Engineering*, 36: 337-347.
- [14] Larasanti, F., R. I. Pratama, K. Haetami, and I. Rostini. 2019. The addition of seaweed flour (*Kappaphycus alvarezii*) and carrageenan on the preference level of freshwater pomfret meatballs. *World Scientific News*, 134 (2): 52-62.
- [15] Lebert, J. 2003. Meat: Eating Quality. *Encyclopedia of Food Sciences and Nutrition (Second Edition)*: 3777-3785.
- [16] Liviawaty, E. and Afrianto. 1998. *The Maintenance of The Eel*. Canisius. Yogyakarta. 133 p.
- [17] Maina, J. 2018. Analysis of The Factors That Determine Food Acceptability. *The Pharma Innovation Journal*, 7(5): 253-257.
- [18] Manurung, H. 2009. The effect of Substitution of Fish With seaweed and A Prolonged Storage on The Quality of Fish Balls. *Journal of VISION*, 17 (3): 217-228.
- [19] Marimin. 2004. *Techniques and Application of Decision Making Criteria Compound*. PT Grasindo. Jakarta.
- [20] Moehyi, S. 1992. *Organizers Food and Catering Services*. Bharata. Jakarta.
- [21] Nafsiyah, I., M. Nurilmala and A. Abdullah. 2018. The Composition of Nutrients Eel Fish *Anguilla bicolor bicolor* and *Anguilla marmorata*. *Journal of Fishery Products Processing Indonesia*, 21 (3): 504-512.
- [22] National Standardization Body. 2014. *Fish Balls*. SNI 7266:2014. The National Standardization Agency. Jakarta.
- [23] Oktaviani, D. 2012. Its physico-Chemical characteristics of the Gel and Meatballs from Surimi Fish Sailfish (*Istiophorus* sp.) The Frequency of Laundering Two Times. *Thesis*. Faculty of Fisheries and Marine Science. Institute Bogor Agricultural. Bogor.
- [24] Pratama, R. I., I. Rostini and M. Y. Awaluddin. 2013. The Composition of The Content of Compound Flavor of Goldfish (*Cyprinus carpio*) Fresh and Steaming Results. *Journal Aquatika*, 4 (1): 55-67.
- [25] Pratama, R. I., I. Rostini and E. Rochima has. 2018. The Amino Acid Profile and Volatile Flavour Compounds of Raw and Steamed Patin Catfish (*Pangasius hypophthalmus*) and Narrow-barred Spanish Mackerel (*Scomberomorus commerson*). *IOP Conference Series: Earth and Environmental Science*. 116 012056 1-17.
- [26] Pratama, R. I., I. Rostini and E. Liviawaty. 2018. Volatile Components of Raw Pangasius Catfish (*Pangasius hypophthalmus*) and Nile Tilapia (*Oreochromis niloticus*). *IOP Conference Series: Earth and Environmental Science*. 176 012040 1-8.
- [27] Princestasari, L. D. and L. Amalia. 2015. Formulation of Seaweed (*Gracilaria* sp.) In the Manufacture of Beef Meatballs High Fiber and Iodine. *J. Nutritional Food*, 10 (3): 185-196.
- [28] Purba, M. 2014. The formation of the Flavor of Poultry Meat by a Process Heating and Oxidation of Lipids. *WARTAZOA*, 24(3): 109-118.
- [29] Purwanto, E. 2006. The Influence of The Addition of Seaweed (*Kappaphycus alvarezii*) On the Quality of The Mochi Cakes. *Thesis*. Faculty Fisheries and Marine Science. Bogor Institute of Agriculture. Bogor.
- [30] Rahmawati, L. A. 2013. Utilization of The Eel (*Monoptherus albus Zuiewu*) in the Manufacture of Meatballs. *Thesis*. The Faculty of Ecology Human. Bogor Institute of Agriculture. Bogor.

- [31] Ristanti. 2003. The Manufacture of Seaweed Flour (*Eucheuma cottonii*) As a Source of Iodine and *Dietary Fiber*. Thesis. Faculty of Agricultural Technology. Bogor Institute of Agriculture. Bogor.
- [32] Sahlan, S. E. Liviawaty, I. Rostini, and R. I. Pratama. 2018. The Difference in Types of Fish as Raw Material to The Level of Preference Kamaboko. *Journal of Fisheries and Marine*, 9 (1): 129-133.
- [33] Santosa, Andasuryani, and D. Kurniawan. 2016. The Characteristics of The Flour Seaweed (*Eucheuma cottonii*). *The National Conference of Applied Sciences, Engineering, Business and Information Technology*, 1 (1): 346-361.
- [34] Soekarto, S. T. 1985. *Assessment Organoleptic to the Industry of Food and Agricultural products*. Bhratara Karya Aksara. Jakarta.
- [35] Suitha, I. M. and Suhaeri, A. 2008. *The Cultivation of Eels*. PT. Agromedia Pustaka. Jakarta.
- [36] Supriadi, C. 2004. Supplementation of Seaweed Flour (*Eucheuma cottonii*) In the Manufacture of Bread and Cookies. Thesis. Faculty of Agricultural Technology. Bogor Institute of Agriculture. Bogor.
- [37] Wibowo, S. 2000. *Making Meatballs and Fish balls*. Penebar Swadaya. Jakarta.
- [38] Wulandari, M. and E. Handarsari. 2010. The Influence of Addition of Rice Bran to Protein Concentration and Organoleptic Characteristic. *Journal of Food and Nutrition*, 1 (2): 55-62.

