

Table 2. Dry Choux Pastry Yield

Red Tilapia Bone Flour (%)	Weight (g)		Dry Choux Pastry Yield (%)
	Dough	Dry Choux Pastry	
0	497	187	37,62
7,5	506	190	37,55
10	516	193	37,40
12,5	528	197	37,31

Based on the results above, the yield of dry choux pastry decreases with the increasing fortification of red tilapia bone flour. This is because a bone meal is hygroscopic which absorbs water. Red tilapia bone flour can absorb the water content in a dry choux pastry batter so that it will result in a decrease in water content. According to Nurfitriani et. al (2018), a decrease in water content that is high enough to cause gluten elasticity to decrease, the binding capacity or structure of the dough built by gluten is less integrated or less compact, causing dry bread to be a little hard and produce residual crumbs, thus making the yield less and less.

Appearance

Appearance is an important organoleptic parameter because it is a sensory trait that is first seen by consumers (Herliany et al. 2013). The appearance results are shown in Table 3.

Table 3. The Dry Choux Pastry Appearance Mean in Each Treatment

Red Tilapia Bone Flour (%)	Median	Appearance Mean
0	7	7,8 b
7,5	8	8 b
10	7	6,7 ab
12,5	5	6 a

Explanation: The mean followed by the same letter showed that there are no significant differences according to the multiple comparison test at 95% confidence interval.

Based on statistical tests, the fortification of red tilapia bone flour in dry choux pastry did not give a real difference in appearance at 0%, 7,5%, and 10% treatments, while 12,5% treatment was significantly different from the 0% and 7,5% treatments. where the color is slightly brown compared to other treatments. The brownish color of dry sus cakes is caused by a reaction non-enzymatic browning (Maillard). Reaction Maillard is a reaction between carbohydrates and proteins that are affected by temperature, resulting in a color change to brown called melanoidin. Melanoidin is formed when glucose and lysine amino acids react at high temperatures. The lysine amino acid comes from the breakdown of the helical structure and the collagen peptide bond due to gradual heating. A collagen is mostly a form of protein in a bone meal (Rachmansyah et al. 2018).

Aroma

The aroma is one of the factors that will determine consumers to choose products because the aroma can attract consumers' attention to these food products (Bakhtiar et al. 2019). The aroma results are shown in Table 4.

Table 4. The Dry Choux Pastry Aroma Mean in Each Treatment

Red Tilapia Bone Flour (%)	Median	Aroma Mean
0	7	6,7 a
7,5	7	7,1 a
10	8	7,6 a
12,5	7	6,6 a

Explanation: The mean followed by the same letter showed that there are no significant differences according to the multiple comparison test at 95% confidence interval.

Based on statistical tests, fortification of red tilapia bone flour did not significantly affect the aroma of dry choux pastry. The dry choux pastry has a distinctive aroma of dry choux pastry and the aroma of red tilapia bone which is not detected by the panelist senses because red tilapia bone flour has a fishy aroma that is not too fragrant, so there is no difference in aroma from the addition of red tilapia flour. According to Nurfitriani et. al. (2018), red tilapia bone flour has a neutral and fishy aroma that is not too fragrant.

The aroma of dry choux pastry is influenced by the components of the ingredients used in making dry choux pastry including salt, margarine, and eggs. The use of salt, margarine, and eggs in cake processing will be able to increase the distinctive aroma of

the product. According to Oktariani et al. (2017), salt and margarine can increase the aroma in the processing during combustion. While materials such as eggs will produce aroma compounds such as aldehydes that can produce aromas such as fruit and butter, as well as pyrazine compounds that can produce distinctive aromas on a sponge (Imami and Sutrisno 2018).

Texture

The texture is one of the parameters that play an important role in the acceptability of a food product (Agustina et al. 2016). The texture results are shown in Table 5.

Table 5. The Dry Choux Pastry Texture Mean in Each Treatment

Red Tilapia Bone Flour (%)	Median	Texture Mean
0	5	5,9 a
7,5	7	6,2 ab
10	7	6,9 ab
12,5	9	7,6 b

Explanation: The mean followed by the same letter showed that there are no significant differences according to the multiple comparison test at 95% confidence interval.

Based on statistical tests, the fortification of red tilapia bone flour in dry choux pastry did not give a significant texture difference in the treatment of 7,5%, 10%, and 12,5% while the 0% treatment was significantly different from the 12,5% treatment where the texture was less crisp compared to other treatments. The highest average texture value is in the treatment of adding 12,5% red tilapia bone flour which is 7,6 (likes) with the texture of a crisp dry choux pastry when bitten in the mouth. This is due to the addition of a greater amount of red tilapia bone flour so that it will add crispness to the dry choux pastry when bitten in the mouth. By the statement Darmawangsyah et al. (2016), that the greater addition of fishbone meal will add crispness to the resulting cookies. The crispness of dry choux pastry is caused by the hydrophilic groups in the protein contained in the fishbone meal much more than starch, thus making the texture more savory and crispy.

Taste

Taste is the most important factor in determining whether or not a food product is preferred compared to other factors because the taste is a determining factor for a product to be accepted or not (Rachmansyah et al. 2018). The taste results are shown in Table 6.

Table 6. The Dry Choux Pastry Taste Mean in Each Treatment

Red Tilapia Bone Flour (%)	Median	Taste Mean
0	7	6,8 a
7,5	7	7,1 a
10	8	7,8 a
12,5	7	6,7 a

Explanation: The mean followed by the same letter showed that there are no significant differences according to the multiple comparison test at 95% confidence interval.

Based on statistical tests, all treatments had no significant effect on the taste of dry choux pastry. That is because red tilapia bone flour has a tasteless taste. By the statement Nurfitriani et al. (2018), that red tilapia bone flour has a tasteless or tasteless taste. The lowest average value of the taste of dry choux pastry was found in the treatment of adding 12,5% red tilapia bone flour which was 6,7 (liked). This is due to the large amount of red tilapia bone flour that is added to the treatment of adding 12,5% red tilapia bone flour which results in a stronger fish bone flour flavor so that the distinctive taste of dry choux pastry will be reduced, as well as more chalky. According to Bunta et al. (2013), the main mineral component of fishbone flour is calcium which can give a chalky taste, so the more concentration of fishbone flour is added, the taste in fishbone flour is increasingly felt so that the level of panelists' preference for bagea cake decreases.

Decision Making by bayes method

The calculation results in determining the best treatment using the Bayes method by considering the appearance, aroma, texture and taste criteria of red tilapia bone flour dry choux pastry are shown in Table 7.

Table 7. Decision Matrix of Red Tilapia Bone Flour Dry Choux Pastry Assessment with Bayes Method

Treatment (%)	Criteria				Alternative Value
	Appearance	Aroma	Texture	Taste	
0	7	7	5	7	6,64
7,5	8	7	7	7	7,18
10	7	8	7	8	7,64
12,5	5	7	9	7	6,99
Criteria Value	0,18	0,14	0,18	0,50	

Based on the calculation of the weight criteria for appearance, aroma, texture, and taste of dry choux pastry obtained the highest number of criteria weight is the flavor parameter of 0,50 followed by the appearance, aroma, and texture. This shows that the taste criteria are the panelists' main consideration in choosing the product of dry choux pastry. Based on all parameters observed, the addition of 10% red tilapia bone flour was the panelist's most preferred treatment compared to other treatments.

Hardness Level

Hardness testing was carried out on the treatment without the addition of red tilapia bone flour and the most favored dry choux pastry treatment by the panelists based on the level of preference (10%). The results of the test for hardness of dry choux pastry are presented in Table 8.

Table 8. The hardness of Dry Choux Pastry

Red Tilapia Bone Flour (%)	Hardness of Dry Choux Pastry (g Force)
0	921,91
10	1161,81

Based on the above results, the greater the amount of addition of red tilapia bone flour to dry sus cakes, the hardness level of dry choux pastry increases. The hardness of the dry choux pastry is caused by the addition of red tilapia bone flour containing calcium and phosphorus which causes the water content of the dry choux pastry to be lower to cause the higher hardness value. By the statement Syadeto et al. (2017), the more addition of fish bone meal, the harder the product is related to the high calcium and phosphorus content in fishbone meal so that the hardness of the product will also change according to the amount of fishbone meal concentration.

Proximate

The proximate test is carried out to determine the nutritional content contained in dry choux pastry the most susceptible matter by panelists was based on the results of the assessment of the level of preference of the dry choux pastry, namely the treatment of adding 10% red tilapia bone flour and also testing the treatment without adding red tilapia bone flour. The results of the proximate analysis of the sample of dry choux pastry are presented in Table 9.

Table 9. Proximate Analysis Results of Dry Choux Pastry Samples(%)

Parameters (%)	Treatment(%)	
	0	10
Calcium Content	0,028	0,486
Water Content	3,45	2,46
Protein Content	9,16	10,04
AshContent	2,52	5,01
Fat Content	44,69	42,91

In general, the addition of red tilapia bone flour can increase calcium content, protein content, and ash, while water and fat content tend to decrease. Changes in the increase and decrease will be explained in the next section.

Calcium Content

The value of the calcium level of the treatment is 10% greater than the treatment of 0%. This shows that the addition of red tilapia flour to dry choux pastry can increase the calcium content in dry choux pastry. This is because red tilapia flour is the highest mineral source in the raw material for making dry choux pastry. The main minerals found in fish bones include calcium, phosphorus, and carbonate (Susanti et al. 2011). Fishbones contain a lot of calcium in the form of calcium phosphate as much as 14% of the total

bone structure. This form of calcium phosphate complex is found in bones and can be absorbed by the body well around 60-70% (Edam 2016).

Water Content

The value of water content shows a decrease in yield along with the increase of red tilapia bone flour. This is because the water contained in dry choux pastry will be bound by Ca^{++} found in fishbone flour, so the water content decreases. According to Darmawangyah et al. (2016), that with the addition of fishbone meal also means the addition of Ca^{++} which will find particles OH^- which are part of the water elements or H_2O so that the water content decreases with the addition of fishbone meal. Also, during the roasting process, a lot of water is evaporated from the dry choux pastry mixture, causing the water content to decrease. According to Haryani et al. (2017), that in the roasting process the water content will decrease due to the process of moving the mass of water from the middle of the product to the surface. While on the surface, the water content is very quickly evaporated so that by the time of roasting, the water content of the material has been lost a lot. Water content in pastries has fulfilled SNI 01-2973-1992 which is a maximum of 5%.

Protein Content

The value of the protein content of the resulting dry choux pastry rises together with the increasing amount of red tilapia flour. This is influenced by the high protein content in the red tilapia bone flour that is added, so the protein content of the dry choux pastry is increasing. Tilapia bone meal contains 40,8% protein (Petenuci et al. 2008). The use of raw materials that contain high protein will produce processed products that have high protein content, and vice versa (Hasniar et al. 2019). In addition to the use of added ingredients, the high and low values of protein are largely influenced by water content. According to Pratama et al. (2014), that the high or low measured protein value can be influenced by the amount of water content lost (dehydration) from the material. The measured protein value will be even greater if the amount of water lost is greater. The protein content of dry sus cake has fulfilled SNI 01-2973-1992 which is a minimum of 9%.

Ash Content

The value of ash content increases with an increasing amount of red tilapia bone flour. This is because red tilapia bone flour contains a lot of minerals, so the higher the amount of addition of red tilapia bone flour, the higher the ash content produced. By the statement Darmawangyah et al. (2016), the value of high ash content in pastries with the addition of fishbone flour treatment was caused by the main constituent components of fish bones containing living cells in the form of mineral salts. Mineral salts consisting of several elements include 80% calcium phosphate and the rest consists of calcium carbonate and magnesium phosphate. The ash content of dry choux pastry does not meet SNI 01-2973-1992 which is a maximum of 1.5%.

Fat content

The value of fat content shows a decrease with an increasing percentage of red tilapia bone flour. This can occur because the addition of a fishbone meal causes ash to increase resulting in decreased fat content. Ash content comes from mineral salt components consisting of calcium carbonate and magnesium phosphate. Decreased fat levels are caused by fat reacting with calcium and magnesium then forming deposits and eventually becoming crust, so that the fat read will decrease. According to Cholil et al. (2016), calcium and magnesium compounds react with a mixture of fat and base to form a precipitate. Because calcium and magnesium compounds are relatively difficult to dissolve in water, they tend to separate from the solution in the form of a precipitate or precipitate which eventually becomes a crust. The fat content value of dry choux pastry has fulfilled SNI 01-2973-1992 which is a minimum of 9.5%.

Conclusion

Based on the results of the study, fortification of red tilapia bone flour up to 12,5% is still preferred by panelists based on hedonic tests. The treatment of adding 10% red tilapia bone flour is most preferred compared to other treatments based on the test *Bayes* having a higher alternative value of 7,64 with the calcium content of 0,486%, water content of 2,46%, the protein content of 10,04%, the ash content of 5,01%, fat content 42,91% and hardness 1161,81 gf.

References

- [1] Afrianto, E., dan E. Liviawaty. 1989. *Pengawetan dan Pengolahan Ikan*. Penerbit Kanisius. Yogyakarta.
- [2] Agustina, L., Udiantoro., dan Suhandriyanto. 2016. Penentuan Formulasi Bahan Tambahan sebagai Bahan Baku Substitusi Produksi Tempe Menggunakan Uji Ambang Batas (*Threshold*) dan Uji Kesukaan (Hedonik). *Ziraa'ah*, 40 (2): 212-221.
- [3] Badan Standardisasi Nasional. 1992. *Syarat Mutu Kue Kering*. (SNI 01- 2973-1992). Jakarta.
- [4] Bakhtiar, S. Rohaya., dan H. M. Ayunda. 2019. Penambahan Tepung Tulang Ikan Bandeng (*Chanoschanos*) sebagai Sumber Kalsium dan Fosfor pada Pembuatan Donat Panggang. *Jurnal Teknologi dan Industri Pertanian Indonesia*, 11 (1): 38-45.
- [5] Bunta, D. I., A. S. Naiu., dan N. S. Yusuf. 2013. Pengaruh Penambahan Tepung Tulang Ikan Tuna terhadap Karakteristik Hedonik Kue Bagea Khas Gorontalo. *Jurnal Ilmiah Perikanan dan Kelautan*. 1 (2): 81-88.
- [6] Cholil, M., A. N. Anna., dan N. Setyaningsih. 2016. Analisis Kesadahan Air Tanah di Kecamatan Toroh Kabupaten Grobogan Propinsi Jawa Tengah. *ISSN 2407-9189*, 88-89.
- [7] Darmawangsyah., P. Jamaludin., dan Kadirman. 2016. Fortifikasi Tepung Tulang Ikan Bandeng (*Chanos chanos*) dalam Pembuatan Kue Kering. *Jurnal Pendidikan Teknologi Pertanian*, 2 (2): 149-156.
- [8] Direktorat Jenderal Perikanan Budidaya Kementerian Kelautan dan Perikanan. 2016. *Laporan Kinerja 2016*. Direktorat Jenderal Perikanan Budidaya Kementerian Kelautan dan Perikanan. Jakarta.
- [9] Edam, M. 2016. Fortifikasi Tepung Tulang Ikan terhadap Karakteristik Fisiko-Kimia Bakso Ikan. *Jurnal Penelitian Teknologi Industri*, 8 (2): 83-90.
- [10] Haryani, K., Hargono., N. A. Handayani., P. Ramadani., dan D. Rezekia. 2017. Substitusi Terigu dengan Pati Sorgum Terfermentasi pada Pembuatan Roti Tawar: Studi Suhu Pemanggangan. *Jurnal Aplikasi Teknologi Pangan*, 6 (2): 61-64.
- [11] Hasniar., M. Rais., dan R. Fadilah. 2019. Analisis Kandungan Gizi dan Uji Organoleptik pada Bakso Tempe dengan Penambahan Daun Kelor (*Moringa oleifera*). *Jurnal Pendidikan Teknologi Pertanian*, 5 (1): S189 – S200.
- [12] Herliany, N.E., J. Santoso., dan E. Salamah. 2013. Penggunaan Coating Karaginan terhadap Mutu Organoleptik Udang Kupas Rebus Selama Penyimpanan Dingin. *Jurnal Agroindustri*, 3 (2): 61 – 70.
- [13] Imami, R.H., dan A. Sutrisno. 2018. Pengaruh Proporsi Telur dan Gula serta Suhu Pengovenan terhadap Kualitas Fisik, Kimia dan Organoleptik pada Bolu Bebas Gluten dari Pasta Ubi Kayu (*Manihot esculenta*). *Jurnal Pangan dan Agroindustri*, 6 (3): 89-99.
- [14] Justicia, A., E. Liviawaty., dan H. Hamdani. 2012. Fortifikasi Tepung Tulang Nila Merah sebagai Sumber Kalsium terhadap Tingkat Kesukaan Roti Tawar. *Jurnal Perikanan dan Kelautan*, 3 (4): 17-27.
- [15] Marom, A., P. Astuti., dan M. F. Putri. 2014. Pengaruh Penggunaan Tepung Kulit Ari Biji Kedelai sebagai Bahan Substitusi terhadap Kualitas Choux Pastry Kering. *Food Science and Culinary Education Journal*, 3 (1): 56-57.
- [16] Nurfitriani, S.A., E. Liviawaty., I. Nurruhwati., and Junianto. 2018. Red Tilapia's Bone Flour Fortification as A Source of Calcium on Dried Bread's Acceptance Level. *Global Scientific Journals*, 6 (10): 18-26.
- [17] Oktariani, E., Elida., dan R. Holinesti. 2017. Pengaruh Substitusi Tepung Ubi Jalar terhadap Kualitas Sus Kering. *Journal Home Economic and Tourism*, 15 (2): 1-18.
- [18] Petenuci, M.E., F. B. Stevanato., J.E.L. Visentainer., M. Matsushita., E. E. Garcia., N.E. de Souza., and J.V. Visentainer. 2008. Fatty Acid Concentration, Proximate Composition, and Mineral Composition in Fishbone Flour of Nile Tilapia. *Archivos Latino Americanos de Nutricion*, 58 (1): 87-90.
- [19] Pratama, R.I., I. Rostini., dan E. Liviawaty. 2014. Karakteristik Biskuit dengan Penambahan Tepung Tulang Ikan Jangilus (*Istiophorus sp.*). *Jurnal Akuatika*, 5 (1): 30-39.
- [20] Putranto, H. F., A. N. Asikin., dan I. Kusumaningrum. 2015. Karakterisasi Tepung Tulang Ikan Belida (*Chitala sp.*) sebagai Sumber Kalsium dengan Metode Hidrolisis Protein. *Ziraa'ah*, 40 (1): 11-20.
- [21] Rachmansyah, F., E. Liviawaty., A. Rizal., dan N. Kurniawati. 2018. Fortifikasi Tepung Tulang Cakalang sebagai Sumber Kalsium terhadap Tingkat Kesukaan Kerupuk Gendar. *Jurnal Perikanan dan Kelautan*, 9 (1): 62-70.
- [22] Rohmah, D.Y.N., Y.S. Darmanto., dan U. Amalia. 2015. Karakteristik Lem dari Tulang Ikan dengan Habitat yang Berbeda (Payau, Tawar, Laut). *Jurnal Pengolahan dan Bioteknologi Hasil Perikanan*, 4 (2): 11-16.
- [23] Saranaung, A., M. S. Sangi., dan D. G. Katja. 2018. Pengaruh Ukuran Bahan terhadap Rendemen dan Kualitas Minyak Biji Pala (*Myristica Fragrans Houtt*) dengan Metode Soxhletasi. *Jurnal Mipa Unstrat Online*, 7 (1): 39-43.
- [24] Sumandiarsa, I. K., A. N. Siregar., dan R. O. Priadi. 2017. Mutu dan Perhitungan Biaya Pembekuan Fillet Ikan Nila (*Oreochromis niloticus*) menggunakan Contact plate freezer Skala Laboratorium. *Jurnal Akuatika Indonesia*, 2 (1): 79-86.
- [25] Susanti, L., M. Zuki., dan F. Syaputra. 2011. Pembuatan Mie Basah Berkalsium dengan Penambahan Tulang Ikan Tenggiri (*Somberomerus lineolatus*). *Jurnal Agroindustri*, 1 (1): 35-44.
- [26] Syadeto, H. S., Sumardiarto., dan L. Purnamayati. 2017. Fortifikasi Tepung Tulang Ikan Nila (*Oreochromis niloticus*) sebagai Sumber Kalsium dan Fosfor serta Mutu Cookies. *Jurnal Ilmiah Teknosains*, 3 (1): 17-21.