

- characterization of their antifungal activity,” *J. Med. Chem.*, vol.52, no.5, pp. 1334–1351,.
- [13] T. Zhu Dawal M.S More, Shanshan Shon and Qin Lu Xingpai yeet *al.*, (2017) “Design and synthesis of novel N(4)-substituted thiosemicarbazones bearing a pyrrole unit as potential anticancer agents,” *Oncol. Lett.*, vol. 13, no. 6, pp. 4493–4500,
- [14] Z. Piri, Z. Moradi-Shoeili, and A. Assoud, (2017) “New copper(II) complex with bioactive 2–acetylpyridine-4N-p-chlorophenylthiosemicarbazone ligand: Synthesis, X-ray structure, and evaluation of antioxidant and antibacterial activity,” *Inorg. Chem. Commun.*, vol. 84, pp. 122–126,
- [15] R. Ramachandran, G. Prakash, P. Vijayan, P. Viswanathamurthi, and J. Grzegorz, (2017) “Inorganica Chimica Acta Synthesis of heteroleptic copper (I) complexes with phosphine-functionalized thiosemicarbazones : An efficient catalyst for regioselective N -alkylation reactions,” *Inorganica Chim. Acta*, vol. 464, pp. 88–93.
- [16] O. Abdullahi, W. Handy, M. Shazwan, S. Jamil, and M. Shamsuddin, (2020) “Copper oxide derived from copper (I) complex of 2-acetylpyridine-N (4) - (methoxy phenyl) thiosemicarbazone as an efficient catalyst in the reduction of 4-nitrophenol,” vol. 16, no. 3, pp. 351–358,
- [17] T. S. Lobana, R. J. Butcher, T.W.Failes and P. Turner (2012) “Pyrrole-2-carbaldehyde thiosemicarbazones of Nickel(II) and palladium(II): Synthesis, structure, and spectroscopy,” *Zeitschrift fur Anorg. und Allg. Chemie*, vol. 638, no. 5, pp. 804–810,
- [18] T. S. Lobana, G. Bawa, A. Castineiras, and R. J. Butcher, (2007) “First example of pyrrole-2-carbaldehyde thiosemicarbazone as tridentate dianion in [Pd(η^3 -N₄,N₃,S-ptsc)(PPh₃)] complex,” *Inorg. Chem. Commun.*, vol. 10, no. 4, pp. 506–509,
- [19] D. R. Richardson (2013) “2-Acetylpyridine thiosemicarbazones are potent copper chelators and antiproliferative agents: *J. Med. Chem.*, vol. 52, No. 6, pp. 1218–1233,
- [20] P. F. Rapheal, E. Manoj, and M. R. Prathapachandra Kurup,(2007) “Copper(II) complexes of N(4)-substituted thiosemicarbazones derived from pyridine-2-carbaldehyde: Crystal structure of a binuclear complex,” *Polyhedron*, vol. 26, no. 4, pp. 818–828..
- [21] R. Manikandan, P. Viswanathamurthi, K. Velmurugan, R. Nandhakumar, T. Hashimoto, and A. Endo, (2014,) “Synthesis, characterization and crystal structure of cobalt(III) complexes containing 2-acetylpyridine thiosemicarbazones: DNA/protein interaction, radical scavenging and cytotoxic activities,” *J. Photochem. Photobiol. B Biol.*, vol. 130, pp. 205–216,
- [22] G. A. Bain, D. X. West, J. Krejci, J. Valdc-, S. Hernbndez-ortega, and R. A. Toscanob, (1997) “Synt & etic and spectroscopic iuve & gatims of N (4) -substituted isatin thiosemicarbazones and their copper complexes,” *Science (80)* vol. 16, No. 5.
- [23] M. Hakimi, H. Vahedi, M. Rezvaninezhad, E. Schuh, and F. Mohr,(2011) “Synthesis and characterization of copper(I) complexes from triphenylphosphine and isatin Schiff bases of semi- and thiosemicarbazide,” *J. Sulfur Chem.*, vol. 32, no. 1, pp. 55–61,.
- [24] N. Verma and N. Kumar, (2019) “Synthesis and Biomedical Applications of Copper Oxide Nanoparticles: An Expanding Horizon,” *ACS Biomater. Sci. Eng.*, vol. 5, no. 3, pp. 1170–1188,
- [25] F. K. Kadir, M. Shamsuddin, and M. M. Rosli, (2016) “Crystal structure of bis(acetophenone 4-benzoylthiosemicarbazonato- κ^2 N1,S) nickel(II),” *Acta Crystallogr. Sect. E Crystallogr. Commun.*, vol. 72, no. ii, pp. 760–763,
- [26] S. S. Chavan, S. K. Sawant, S. B. Pawal, and M. S. More, “(2016) Copper(I) complexes of

- 2-methoxy-(5-trifluoromethyl-phenyl)-pyridine-2-yl-methylene-amine: Impact of phosphine ancillary ligands on luminescence and catalytic properties of the copper(I) complexes,” *Polyhedron*, vol. 105, pp. 192–199,
- [27] R. Ramachandran, G. Prakash, P. Vijayan, P. Viswanathamurthi, and J. G. Malecki,(2017) “Synthesis of Heteroleptic Copper (I) Complexes with Phosphine- Functionalized Thiosemicarbazones : Efficient Catalyst for Regioselective N- Alkylation Reactions,” *Inorganica Chim. Acta*, No. I, pp.162 - 177
- [28] A. N. Jadhav and S. S. Chavan, (2014) “Alkynyl functionalized iminopyridine copper (I) phosphine complexes: Synthesis, spectroscopic characterization and photophysical properties,” *J. Lumin.*, vol. 148, pp. 296–302,
- [29] I. Ali, W. A. Wani, and K. Saleem, (2013) “Empirical formulae to molecular structures of metal complexes by molar conductance,” *Synth. React. Inorganic, Met. Nano-Metal Chem.*, vol. 43, No. 9, pp. 1162–1170,
- [30] M. Hakimi, K. Moeini, Z. Mardani, and R. Takjoo,(2014) “Synthesis and spectral study of a copper(I) complex, [CU(L)(PPH₃)₂], with ns-donor ligand,” *Phosphorus, Sulfur Silicon Relat. Elem.*, vol. 189, no. 5, pp. 596–605,

© GSJ