



Study of some heavy metals and the p^H assay in walnuts found in four different locations Port-Harcourt, Nigeria.

¹ Don-lawson C, and ²R. Okah

Department of Science Laboratory Technology, Captain Elechi Amadi Polytechnic, Rumuola, Rivers State.

Corresponding Author's mail; ogweru12345@gmail.com

Abstract

The analysis of lead (Pb), Iron (Fe), and P^H in walnuts were carried out in four different locations which includes Rumuokoro, Rumuokwuta, mile 3 and Aba road, the lead, iron and nickel concentrations were analysed using atomic absorption spectrophotometer. (AAS) while the P^H of the samples were analysed using a ph meter . the results obtained showed wide variations of concentrations of heavy metals present which includes lead (5.38mg), iron, (0.168mg) for rumuokoro, for rumuokwuta lead,(6.38mg). iron.(1.168mg) , for aba road, lead,(4.38). iron,(1.138mg) and mile 3 to borikiri axis,lead (6.30mg),iron (1.130mg) the level of lead in the four different locations were high compared to the World Health Organisation recommendation which is about 0.005mg this was actually attributed to the industrial activities taking place in these areas, and this may result to some health risks like cancer, asthma, hypertention,etc if the walnuts sold in these areas are excessively consumed, however, the P^H result obtained for the four different locations were 5.80,5.70,5.80,5.60 and 5.80 respectively, this revealed that the cooked walnuts are quite acidic and may be due to the sources of water used.

Keywords: walnuts, Iron, Lead, atomic absorption spectrophotometer etc.

Corresponding Author: ogweru12345@gmail.com

INTRODUCTION

The walnut is economically very important tree species cultivated throughout the world such as the west indies, japan, china etc. the determination of lead, iron, nickel and p^H in walnuts is also very imperative because of their biological and economical importance of walnut (Akinhanmi et al,2008). African walnut tetracarpidium belong to the family of euphirbiacea and is commonly found in south-east and south-west Nigeria and Cameroon.

Walnut(*Tetracarpidiumconophorum*) is an important crop that is cultivated throughout the world's temperate regions for its edible nuts (Edem etal 2009). It is known in the southern Nigeria as ukpa (Igbo), Western Nigeria as asala (Yoruba). It is known in the littoral and the Western Cameroon as kaso or ngak (Adeleye et al, 1991). They are usually planted under an indigenous tree that can provide strong support for the heavy weight of the climber when fully established on the crown of the tree, and in cases where they cannot be harvested manually; they are left for full maturation after which the pod falls off by itself and are picked, removed from the rotten pods, washed and sold in the markets(Hemery,2006). *T.conophorum*, like many other plants in Africa and in many other parts of the world has been proven to have decorative, nutritive, medicinal, agricultural and industrial values over the years. Conophor are cultivated principally for the nuts which are usually cooked and consumed as snacks(Enujiugha and Ayodele, 2003). *Tetracarpidiumconophorum* contained in a pod which may house; one shelled nut(single), two shelled (double) and three shelled nut(triple). The walnut shells could be black or brown from the plant. The nut is whitish upon cracking from the shell. The nut has a thin layer between two halves (when a nut is divided into two equal parts) of nut. The seed(subglobose) is about 2.5cm long and has wooly materials that attach the nut to the shell when cracked open. A bitter taste is usually observed upon drinking water immediately after eating nuts. This could be attributed to the presence of chemical substances such as alkaloids (Ayodele 2003) Walnut shells is a waste generated in the walnut (*Tetracarpidiumconophorum*) harvest, containing natural compounds with

antioxidants properties. The walnut shell has antioxidant compounds such as flavonoids which have been determined (Akbari et al, 2012). *Tetracarpidium conophorum* can be cooked, roasted or sun dried and toasted seeds could be ground like melon seeds and thickener in soup preparation. The plant is known in Africa especially in the Eastern and Western parts of Nigeria for its antibacterial efficacy (Copur et al, 2007). Decoction of leaves and seeds serve as beverage which relieves abdominal pains and fever (Malu et al, 2009). Dried walnut can be ground and turned into flour which can be used as composite flour during baking or in-place of milk in tea preparation (Stevens and Domelan, 2003). the determination of lead, iron, and p^H in walnuts became very imperative because it is significantly consumed and as food. (Ajaiyeoba and Fadare, 2006).

Materials and Method

The materials used include pH meter, ethanol, (BDH England) AAS, acetone (BDH England), glasswares like beaker, conical flask, flat bottom flask etc, distilled water.

The walnuts were collected from four different locations which includes rumuokoro, rumuokwuta-choba, mile 3-borokiri and Aba road.

Procedure for the determination of heavy metals in walnut

Digestion procedure for metal analysis of the walnuts

- Dry sample in room temperature for seven days or dry in an oven at temperature of 105°
- Grind sample to homogenize lumps
- Sieve sample using 2mm sieve or 250 m mesh to bring sample to the same size
- Take a quantity of sample and weigh in balance (5.0g)
- Add 2ml conc HNO_3 and 6ml conc of HCl a small quantity of water or 10ml HNO_3 / H_2O_2 mixture v/v (3:1) and 4ml of H_2O

- Heat the mixture/ solution in a fume cupboard until brownish foam stops.
- Make the funnel using filter paper and insert over 100ml measuring cylinder
- Transfer the digested content into the filter paper and make up the volume either 20ml or 5ml and not more
- Take the filtrate to the AAS and run the metal of interest.

RESULT AND DISCUSSION

4.1 Fe and Pb test for specimen A (Rumuokoro)

PARAMETER	Method	Result
Pb	AAS	5.38mg
Fe	AAS	0.168mg

4.2 Fe and Pb test for specimen B (Rumuokwuta-Choba)

PARAMETER	Method	Result
Pb	AAS	6.38mg
Fe	AAS	1.168mg

4.3 Fe and Pb test for specimen C (Aba Road)

PARAMETER	Method	Result
Pb	AAS	4.38mg
Fe	AAS	1.138mg

4.4 Fe and Pb test for specimen D (mile 3 Borokiri)

PARAMETER	Method	Result
Pb	AAS	6.30mg
Fe	AAS	1.160mg

4.5 Result of the analysis of P^H samples

Specimen	Result
Specimen A P ^H	5.80
Specimen B P ^H	5.70
Specimen C P ^H	5.60
Uncooked P ^H	6.80

DISCUSSION

From the results obtained in analysing the concentration or level of lead, iron and P^H of walnuts collected from the four different locations in Port Harcourt which are Rumuokoro axis, Rumuokwuta-Choba, Aba road and Mile 3-Borokiri respectively. The results shows that the concentration of lead (Pb) is very high in the different locations which may be attributed to industrial activities of both mining and non-mining activities, the presence of gas flaring, carbonmonoxide and non-oil activities can be sources of lead poisoning which are highly concentrated in these areas, however, the concentration of iron (Fe) is low when compared to that of lead in these areas. Consequently, excessive consumption of walnuts sold in these areas may have some health implications like diarrhoea, low sperm count or infertility, cancer, loss of appetite, etc. Moreover, the concentration of lead in these areas are higher to the World Health Organization reference standard for lead which is 0.005mg. The P^H of the cooked walnuts were also analysed but revealed that the P^H of the cooked walnuts were highly acidic and may be attributed to same industrial activities.

CONCLUSION

From the assay conducted and the results obtained, it is obviously shown that the concentration of lead(Pb) in Rumuokoro axis, Rumuokwuta-Choba , Aba road and mile 3 and Borokiri axis is high and this is likely attributed to both mining and non-mining industrial activities which are easily absorbed or exposed to the walnuts that are public sold or consumed in these areas, it has also shown that the acceptable limits of lead concentration in nuts by WHO is lower than the analysed, however, this may have some health implication due to lead poisoning. The P^H analysis also showed that the walnuts are generally acidic.

ACKNOWLEDGEMENT

I want to whole-heartedly give all glory and honour to the almighty God for his grace, strength and quotient to write, I am also very much grateful to an erudite scholar and researcher, Dr. Donubari-Lawson Chioma for her numerous contributions to research, publications and conferences. God richly bless you.

REFERENCE

Adeleye I.O (1991). Dried poultry droppings as a source of dietary nitrogen for sheep fed corn-cobs based diets.

West African Journal of Biological and Applied Chemistry 36(1):19-32

Ajaiyoba E.O and Fadare D.A (2006) Antimicrobial potential of extracts and fractions of the African walnut-

Tetracarpidium conophorum. African Journal of Biotechnology.

Akbari V. Heidari R. and Jamei R. (2012) Antiradical activity of different parts of walnuts (*Juglans regia* L.) fruit as a function of genotype.

Food Chemistry, 135(4): 2404-2410.

Akinhanmi T.F, Akintokun P.O and Atasi V.N (2008) :“chemical composition and physiochemical properties of cashew nut (*Anacardium occidentale*) oil and cashew nutshell liquid” *Journal of Agricultural, Food and Environmental sciences, 2(1): 1-10*

Ayankunbi M.A, Egele P. and Keshinro O.O (2007) “Effects of methods of preparations on the nutrients composition of some cassava products-Garri(eba), “Lafun” and “Fufu”,

Food Chemistry, 41 (3):349-254

Ayodele O.B (2003). Nutrition in Nigeria:Ibadan, Catoon Publishers, 14th edition
USA:267-270

Ayrulmis N. Avci E. Buyuksari U. and Koc E. (2009) Utilization of pine (*Pinus pinea* L.)

Cone in manufacture of wood based composite for Ecological Manage. Journal of Agriculture, 259:65-70.

Azeredo H.M.C, Faria J.A.F Aparecida M. and Da Silva A.P (2004). *Minimisation of peroxide formation rate in soybean by antioxidants combination. Food Research International, 37(7):689-694.*

Copur Y. Guler C. Akgul M. and Tascioglu C.(2007). Some chemical properties of hazelnut husk and its suitability for particleboard production.

Journal Build Environ, 42(25):68-72

Edem C.A, Dosunmu M. Basse I. and Francesca I. (2009). Determination of Proximate consumption, Ascorbic acid and Heavy metals Content of African Walnut

(Tetracarpidium conophorum). Pakistan Journal of Nutrition, 8(3): 225-

Enujiugha V.N and Ayodele O.O (2003). Evaluation of nutrients and antinutrients in lesser known under-utilized oil seeds.

International Journal of food science and technology, 38:525-528

Hemery G.E (2006). Growing walnut in mixed stands. *Quart Journal or Forestry, 95:31-92*