































<http://www.biomedcentral.com/1472-6963/14/311>

Tsilimingras, D. (2014). Postdischarge Adverse Events. *Austin J Clin Med*, 1(1), 1002.

Van Walraven, C., Mamdani, M., Fang, J., & Austin, P. C. (2004). Continuity of care and patient outcomes after hospital discharge. *Journal of General Internal Medicine*, 19(6), 624–631.

<https://doi.org/10.1111/j.1525-1497.2004.30082.x>

Wilkerson, L., & Blacketer, M. T. (2012). Reducing the Burden of Adverse Drug Events. *The Kentucky Pharmacist*, 13(9), 34–38.

## List of Tables

**Table 1: Adverse Events and Demographic Factors**

Age in years	Reported Adverse Events		Total
	Yes	No	
less than 20 years	4(1.9%)	26(12.6%)	30(14.6%)
21 to 30	17(8.3%)	38(18.4%)	55(26.7%)
31 to 40	7(3.4%)	19(9.2%)	26(12.6%)
41 to 50	7(3.4%)	16(7.8%)	23(11.2%)
51 to 60	13(6.3%)	20(9.7%)	33(16.0%)
61 and above	17(8.3%)	22(10.7%)	39(18.9%)
<b>Total</b>	<b>65(31.6%)</b>	<b>141(68.4%)</b>	<b>206(100%)</b>
<b>Marital status</b>			
Married	51(24.8%)	92(44.7%)	143(69.4%)
Single	6(2.9%)	41(19.9%)	47(22.8%)
Widow (er)	8(3.9%)	8(3.9%)	16(7.8%)
<b>Total</b>	<b>65(31.6%)</b>	<b>141(68.4%)</b>	<b>206(100%)</b>
<b>Sex</b>			
Male	23(11.2%)	45(21.8%)	68(33.0%)
Female	42(20.4%)	96(46.6%)	138(66.9)
<b>Total</b>	<b>65(20.4%)</b>	<b>141(68.4%)</b>	<b>206(100%)</b>
<b>Occupation of participants</b>			
Farmer	17(8.3%)	27(13.1%)	44(21.4)
House wife	2(0.9%)	6(2.9%)	8(3.9%)

-



Aged	11(5.3%)	5(2.4%)	16(7.8%)
Student	7(3.4%)	18(8.7%)	25(12.1%)
Trader	14(6.8%)	26(12.6%)	40(19.4%)
Artisans	3(1.5%)	18(8.7%)	21(10.2%)
Employees	8(3.9%)	22(10.7%)	30(14.6%)
Others	3(1.5%)	19(9.2%)	22(10.8%)
<b>Total</b>	<b>65(31.6%)</b>	<b>141(68.4%)</b>	<b>206(100%)</b>

Educational level of participants

Primary	6(2.9%)	21(10.2%)	27(13.1%)
JHS	8(3.9%)	11(5.3%)	19(9.2%)
SHS	1(0.5%)	18(8.7%)	19(9.2%)
Tertiary	11(5.3%)	25(12.1%)	36(17.5%)
None	39(18.9%)	66(32.0%)	105(50.9%)
<b>Total</b>	<b>65(31.6%)</b>	<b>141(68.4%)</b>	<b>206(100%)</b>

**Table 2: Effect of Demographic Factors on Adverse Events**

Model	Unstandardized Coefficients		Standardized	T	P-value
	B	Std. Error	Beta		
(Constant)	1.626	0.187		8.685	
Age in years	-0.039	0.021	-0.146	-1.875	0.062
Marital status	-0.008	0.054	-0.010	-0.139	0.889
Sex	0.032	0.069	0.032	0.458	0.648
Occupation	0.028	0.016	0.142	1.800	0.073
Educational level	0.006	0.026	0.020	0.241	0.810

Dependent variable: Reported Adverse Events

(Significant level= 0.05)



**Table 3: Effect of Demographic Factors on Severity of Adverse Events**

Model	Unstandardized		Standardized		
	Coefficients		Coefficients	T	P-value
	B	Std. Error	Beta		
(Constant)	2.217	0.371		5.984	
Age in years	-0.044	0.041	-0.085	-1.075	0.284
Marital status	0.044	0.107	0.030	0.413	0.680
Sex	0.145	0.137	0.075	1.059	0.291
Occupation	0.022	0.031	0.056	0.705	0.481
Educational level	-0.014	0.051	-0.022	-0.271	0.787

Dependent variable: Severity of Adverse Events

(Significant level= 0.05)



**Table 4: The influences of Demographic Factors on of the Types of Adverse Event**

Model	Unstandardized Coefficients		Standardized		
	B	Std. Error	Coefficients	T	P-value
(Constant)	7.361	0.967		7.614	
Age in years	-0.209	0.106	-0.153	-1.963	0.049
Marital status	-0.374	0.278	-0.097	-1.344	0.180
Sex	0.185	0.357	0.036	0.518	0.605
Occupation	0.098	0.080	0.096	1.227	0.221
Educational level	-0.073	0.134	-0.044	-0.546	0.586

(Significant level= 0.05)

Dependent variable: Types of Adverse Event



**Table 5: A Cross tabulation of Age and Severity of Adverse Events**

Age in Years	<u>Severity of Adverse Events</u>					Total
	One Day of Symptoms	Several Days of Symptoms	Non-permanent Disability	Death	No Adverse event Reported	
≤20 years	0(0.0%)	4(1.9%)	0(0.0%)	0(0.0%)	26(12.1%)	30(14.6%)
21 to 30	0(0.0%)	10(5.3%)	6(3.4%)	1(0.5%)	36(17.5%)	53(25.7%)
31 to 40	0(0.0%)	3(1.5%)	4(1.9%)	0(0.0%)	19(9.2%)	26(12.6%)
41 to 50	0(0.0%)	3(1.9%)	1(0.5%)	3(1.5%)	15(7.3%)	22(10.7%)
51 to 60	1(0.5%)	7(3.4%)	4(1.9%)	1(0.5%)	20(9.7%)	33(16.0%)
61 and above	2(0.9%)	7(3.4%)	5(2.4%)	3(1.5%)	21(10.2%)	38(18.4%)
<b>Total</b>	<b>3(1.45%)</b>	<b>34(16.5%)</b>	<b>20(9.7%)</b>	<b>8(3.9%)</b>	<b>137(66.5%)</b>	<b>206(100%)</b>

**Table 6: Types of adverse events and age in years Cross tabulation**

Types Of Adverse Events	Age in years						Total
	less than			61 and			
	20 years	21 to 30	31 to 40	41 to 50	51 to 60	above	
Adverse Drugs Events	1(0.5%)	5(2.4%)	2(1.0%)	3(1.5%)	4(1.9%)	7(3.4%)	22(10.7%)
Procedural Related Events	0(0.0%)	1(0.0%)	0(0.0%)	0(0.0%)	2(1.0%)	1(0.5%)	4(1.9%)
Nosocomial Infections	0(0.0%)	2(1.0%)	2(1.0%)	0(0.0%)	0(0.0%)	3(1.5%)	7(3.4%)
Wounds	2(1.0%)	0(0.0%)	1(0.5%)	0(0.0%)	0(0.0%)	2(1.0%)	5(2.4%)
Re-Admissions	0(0.0%)	3(1.5%)	1(0.5%)	2(1.0%)	1(0.5%)	1(0.5%)	8(3.9%)
Deaths	0(0.0%)	1(0.5%)	0(0.0%)	3(1.5%)	1(0.5%)	3(1.5%)	8(3.9%)
Others	2(1.0%)	8(3.9%)	1(0.5%)	0(0.0%)	5(2.4%)	1(0.5%)	17(8.3%)
<b>Total</b>	<b>5(2.4%)</b>	<b>20(9.7%)</b>	<b>7(4.4%)</b>	<b>8(3.9%)</b>	<b>13(6.3%)</b>	<b>18(8.7)</b>	