





result can be obtained within hours up to 2 days. The leukocyte count is variable in these patients. Leukopenia, leukocytosis and lymphopenia reported, but lymphopenia is reported more often. The chest radiography may reveal lung infiltrates. In a CT scan of the lungs, bilateral ground-glass infiltrates associated with dense areas may be seen. Elevated liver transaminases are also reported. <sup>(7,8,22,23,24,25,26,27)</sup>

The fatality rate of the COVID-19 varies in different countries but generally is 1-3%. At the beginning of the outbreak, the WHO reported that the severe cases in china have mostly occurred among people aged 40 years and beyond and among those with associated illnesses and mostly affected men, but this characteristic may be changing. According to the WHO, the recovery time of the disease is about 2 weeks in mild cases and 3-6 weeks in severe cases. <sup>(2,9)</sup>

Several studies have been conducted on COVID-19 in the world so far. A study by Zheng et al. titled evaluation of the clinical and epidemiologic characteristics of the COVID-19 cases in Wuhan city of China between 16 January to 4 February 2020 on 73 patients (40 male and 33 females) majority of the patients were aged 40-49 years. The main symptoms were fever and cough, respectively. In 30 % of the patient's leukopenia was reported, and in 45% of the patients, lymphopenia was present. <sup>(10)</sup>

Another study by Maria khan et al. titled epidemiologic and clinical characteristics of COVID 19 cases in a tertiary hospital of Peshawar, Pakistan between 15 March to 21 April on 121 patients whose PCR tests was positive; the mean age of the patients was 43 years, and 70% of them were males. Common symptoms of the patients were fever, cough, and shortness of breath, respectively. <sup>(11)</sup>

Another study by Tambe et al. between 31 March to 24 April 2020 in a tertiary hospital in Pune state of India conducted in descriptive cross-sectional method shows that among 197 confirmed cases, the age of the majority of the patients was between 30-61 years and in terms of gender males were more affected. Among them, only one patient had a history of travel abroad. In contrast, 57% had a history of contact with a confirmed case, of which 47 % had contact with family members and 10% were infected in the work environment. Dyspnea was the most common symptom of the patients, followed by fever and cough. <sup>(12)</sup>

## **Objective**

The objective of this study is to find the clinical and epidemiologic characteristics of the COVID-19 cases admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020

## **Research Questions**

1. How is the distribution of the COVID-19 cases in terms of gender, age, residence and occupation

2. What are the symptoms and signs of COVID-19 cases in Afghan-Japan communicable diseases hospital
3. What are the laboratory findings of the COVID-19 cases into Afghan-Japan communicable diseases hospital

## **Materials and method**

This is a descriptive cross-sectional study. The data about the COVID-19 cases admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020 was collected using the patients' files and laboratory reports. These data were put in data collection sheets and the data was analyzed using the SPSS-20 and Microsoft Excel.

### **Sample Size**

All the patients admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020 constitute the study's sample size.

### **Sampling Method**

Sampling was convenient; information about all the patients admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020 was found.

### **Inclusion criteria**

- Those patients were admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020, and their PCR test of blood or respiratory samples was positive.

### **Exclusion criteria**

- Those patients whose information was lacking

### **Main variables**

Gender, age, residence, occupation, symptoms, signs, laboratory investigations

### **Research tools**

- Patients files
- Laboratory results
- Data collection sheet

**Results**


This research was conducted within two months in the descriptive cross-sectional retrospective method and included 124 COVID-19 patients admitted to Afghan-Japan communicable diseases hospital Kabul between 20 June to 20 August 2020. The results are presented as follows

**Table 1:** the percentage of COVID-19 prevalence in terms of sex

<b>Sex</b>				
	Frequency	Percent	Valid Percent	Cumulative Percent
Valid male	96	77.4	77.4	77.4
female	28	22.6	22.6	100.0
Total	124	100.0	100.0	

its seen in the table that 77.4% of the COVID-19 patients are males.

**Chart 1:** age distribution of the COVID-19 patients




According to the above chart, the majority of the COVID-19 cases are aged 65-76 years

**Table 2:** Number and percentage of the COVID-19 cases according to the place of residence

		Place			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Kabul	104	83.9	83.9	83.9
	Parwan	2	1.6	1.6	85.5
	Badakhshan	4	3.2	3.2	88.7
	Takhar	4	3.2	3.2	91.9
	Logar	4	3.2	3.2	95.2
	Ghazni	2	1.6	1.6	96.8
	Bamyan	2	1.6	1.6	98.4
	Nengarhar	2	1.6	1.6	100.0
	Total	124	100.0	100.0	

**Chart 2:** the place of residence of COVID-19 patients



As it is seen in the table majority of the COVID-19 cases are the residence of Kabul province


**Table 3:** prevalence of the COVID-19 in terms of occupation

		<b>Job</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Jobless	70	56.5	56.5	56.5
	Shopkeeper	2	1.6	1.6	58.1
	Farmer	5	3.2	3.2	61.3
	Employee	14	11.3	11.3	72.6
	Free worker	4	3.2	3.2	75.8
	Housewife	8	6.5	6.5	82.3
	worker	112	8.1	8.1	90.3
	Cook	9	6.5	6.5	96.8
Total		124	100.0	100.0	

As it can be seen in the above table, the majority of the COVID-19 cases occurred among jobless people (56.5%)

**Table 4:** COVID-19 cases in terms of severity of the disease

		<b>Severity</b>			
		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Severe	100	80.6	80.6	80.6
	Critical	18	14.5	14.5	95.2
	Moderate	6	4.8	4.8	100.0
	Total	124	100.0	100.0	



**Chart 3:** the severity of the COVID-19 patients

According to the above chart, the majority of the COVID-19 cases were in severe form (80.6%)

**Table 5:** symptoms and signs of covid-19 patients

<b>Variables</b>	<b>Total patients(n = 124)</b>	<b>Variables</b>	<b>Total patients(n = 124)</b>
Shortness of breathing	116(93.5%)	Tachycardia	6(4.8%)
Cough	98(79%)	Vomiting	6(4.8%)
Fever	70(56.5%)	Weakness	6(4.8%)
Body pain	52(41.9%)	Anorexia	4(3.2%)
Tachypnea	40(32.3%)	Chest tightness	4(3.2%)



Crepitation	38(30.6%)	Cyanosis	4(3.2%)
Respiratory distress	24(19.4%)	Epigastric pain	4(3.2%)
Sorethroat	22(17.7%)	Chest indrawing	2(1.6%)
Ronchi	12(9.7%)	Limb pain	2(1.6%)
Chest pain	12(9.7%)	Unconscious state	2(1.6%)
Loose motion	10(8.1%)	Hemoptysis	2(1.6%)
Lethargic	8(6.5%)	Hiccup	2(1.6%)
Myalgia	6(4.8%)	Fatigue	2(1.6%)
Dizziness	6(4.8%)	Sneezing	2(1.6%)
Headach	6(4.8%)		

According to the above table, shortness of breath, cough, and fever were the most common symptoms

**Table 6:** Laboratory Findings of Coronavirus Disease (COVID-19)

Laboratory Test	Normal	Reduced	Elevated
Leucocyte	36(29)	8(6.5)	80(64.5)
Lymphocyte	78(62.9)	42(33.9)	4(3.2)
Hemoglobin	86(69.4)	30(24.2)	8(6.5)
Platelet	76(61.3)	26(21)	22(17.7)
ALT	80(64.5)	0(0)	44(35.5)
AST	84(67.7)	0(0)	40(32.3)
ESR	64(51.6)	0(0)	60(48.4)
CRP	62(50)	4(3.2)	58(46.8)
SPO2	62(50)	62(50)	0(0)
Creatinine	108(87.1)	0(0)	16(12.9)

BUN	108(87.1)	0(0)	16(12.9)
-----	-----------	------	----------

Note—Values in parentheses are percentages.

## Discussion

This research work about the clinical and epidemiologic characteristics of the COVID-19 cases in Afghan-Japan communicable diseases hospital was conducted using descriptive-case series retrospective. The data about the COVID-19 cases that were admitted between 20 June to 20 August 2020 were collected using the files of the patients and their laboratory results.

Based on our review of the literature, the following issues were found: according to a study by Zheng et al. about analysis of the clinical and epidemiologic characteristics of the COVID-19 cases conducted in Wuhan city of China between 16 January to 4 February 2020 with 73 participants majority of the were aged 40-49 years. Common symptoms of the patients were fever and cough, respectively. In 30% of the patients, the level of WBC was low, and in 45% there was lymphopenia.<sup>(10)</sup>

The findings of our study are not compatible with this study in terms of age and sex, which may be because the majority of the cases in Afghanistan are among men compared to women, and this could be explained by the fact that men are mostly working outside the home in Afghanistan compared to women. Their risk of contracting the virus is high.

In terms of age, there is also a difference; our study shows the cases among older ages that are not compatible with other studies in the world. And that why the cases are less in Afghanistan may be due lack of predisposing factors. In laboratory investigations in this study, leukopenia was more common. In contrast, in our study, leukocytosis was more common. Still, the presence of lymphopenia is similar in both studies, and also the main symptoms, which are fever and cough, are similar in both studies.

Another study by Maria Khan et al. titled clinical and epidemiologic characteristics of COVID-19 cases in a tertiary hospital in Peshawar, Pakistan, conducted between 15 March to 21 April 2020, which shows that among 121 patients whose PCR test were positive, the mean age of the patients was 43 years and 70% of them were men. The common symptoms were fever, cough and shortness of breath, respectively. The findings of our study are slightly different from this study in the mean age; the remaining results are similar, and it might be the similarity between the culture and traditions and habits of life and shared geography<sup>(11)</sup>

Another study by Tambe et al. between 31 March to 24 April 2020 in a tertiary hospital in Pune state of India conducted in descriptive cross-sectional method shows that among 197 confirmed cases, the age of the majority of the patients was between 30-61 years and in terms of gender males were more affected. Among them, only one patient had a history of travel abroad. In contrast, 57% had a history of contact with a confirmed case, of which 47 % had contact with family members and 10% were infected in the

work environment. Dyspnea was the most common symptom of the patients, followed by fever and cough. Our study is slightly different in the mean age of the patients with this study, and it might be since the men are mostly working outside the home in Afghanistan and are at increased risk of contracting the virus. Also, there is a difference; our study shows higher ages that are compatible with other studies in the world and that why the cases are less in young aged people may be due to the absence of predisposing factors. Rest of the finding is similar and might be due to the shared culture, tradition and life habits and also shared geography<sup>(12)</sup>

Based on the other studies that were reviewed, it can be concluded that in different studies, there are different findings which are normal, but it should be mentioned that compared to other countries; in Afghanistan, there are limitations in different levels in research works which affect the result of study hence more studies should be conducted in this regard.

## Conclusion

The prevalence of the COVID-19 cases is most common among people aged 65-75 years and sex in males and among jobless people. The most common symptoms of the patients were shortness of breath, cough and fever. The most common laboratory changes were leukocytosis, lymphopenia and thrombocytopenia

## Recommendations

1. Establishment of a registration system by the authorities in the Afghan-Japan communicable diseases hospital
2. Establishment of equipped laboratory for bacteriology, virology, biochemistry and serology in the Afghan-Japan communicable diseases hospital
3. Establishment of a computerized database of the patients so that the researchers wouldn't face difficulties.
4. The sample size of this study is limited, so the results may not be representative of the community; hence it's better to conduct more studies
5. Increasing public awareness about the COVID-19 and the ways of transmission and prevention

## References

1. Suleyman G, Fadel RA, Malette KM, Hammond C, Abdulla H, Entz A, et al. Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. *JAMA*. 2020;3(6):1–12.
2. Pormohammad A, Ghorbani S, Baradaran B, Khatami A. Clinical characteristics , laboratory findings , radiographic signs and outcomes of 61 , 742 patients with confirmed

- COVID-19 infection : A systematic review and meta-analysis. *Microb Pathog.* 2020;147(January).
3. J. R-MA, Cardona-Ospina JA, Gutiérrez-ocampo E, Villamizar-peña R, Holguin-rivera Y, Franco-paredes C, et al. Clinical , laboratory and imaging features of COVID-19 : A systematic review and meta-analysis. *Travel Med Infect Dis.* 2020;34(January).
  4. Baj J, Karakuła-Juchnowicz H, ´GT ´nski, Buszewicz G. COVID-19 : Specific and Non-Specific Clinical Manifestations and Symptoms : The Current State of Knowledge. *J Clin Med.* 2020;9(1753):1–22.
  5. Liang W, Guan W, Li C, Li Y, Liang H, Zhao Y. Clinical characteristics and outcomes of hospitalised patients with COVID-19 treated in Hubei (epicentre) and outside. *Eur Respir Dis.* 2020;55(March).
  6. Yu C, Lei Q, Li W, Wang X, Li W, Liu W. Epidemiological and clinical characteristics of 1663 hospitalized patients infected with COVID-19 in Wuhan , China : a single-center experience. *J Infect Public Health.* 2020;13(9):1202–9.
  7. Hassan SA, Sheikh FN, Jamal S, Ezeh JK, Akhtar A. Coronavirus (COVID-19): A Review of Clinical Features , Diagnosis , and Treatment. *Cureus.* 2020;12(3).
  8. Argenziano MG, Bruce SL, Slater CL, Tiao JR, Baldwin MR, Barr RG, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York : retrospective case series. *BMJ.* 2020;10(March).
  9. Jin A, Yan B, Hua W, Feng D, Xu B, Liang L, et al. Biosafety and Health Clinical characteristics of patients diagnosed with COVID-19 in Beijing. *Biosaf Heal.* 2020;2(2):104–11.
  10. Zheng Y, Xiong C, Liu Y, Qian X, Tang Y, Liu L. Epidemiological and clinical characteristics analysis of COVID-19 in the surrounding areas of Wuhan , Hubei Province in 2020. *Pharmacol Res.* 2020;157(January).
  11. Khan M, Khan H, Khan S, Nawaz M. Epidemiological and clinical characteristics of coronavirus disease (COVID-19) cases at a screening clinic during the early outbreak period : a single-centre study. *J Med Microbiol.* 2020;69(March):1114–23.
  12. Tambe MP, Parande MA, Tapare VS, Borle PS, Lakde RN, Shelke SC, et al. An Epidemiological Study of Laboratory Confirmed COVID - 19 Cases Admitted in a Tertiary Care Hospital of Pune , Maharashtra. *Indian J Public Heal |.* 2020;103(42).