

Result and Discussion

Decline in BOD

The reduction in BOD was calculated using the formula from Dwinanto (2009). During the study, each treatment experienced a significant decrease in BOD. The highest reduction in BOD was obtained in the C1 treatment which was 2.72 mg / L while the lowest BOD reduction was obtained in the D1 treatment by 0.32 mg / L. The decrease in BOD during the study can be seen in the following graph.




Figure 1. Decrease of BOD

Treatment A had a small BOD reduction value compared to other treatments. That was due to the absence of organisms that can decompose organic compound in tofu liquid waste. According to Husin (2003), the presence of organic compounds causes the liquid waste of tofu industry to contain high BOD, COD and TSS.

Treatment B, C and D have a large rate of decline due to *L. minor* as a phytoremediator can decompose organic matter. According to Datko et al. (1985), *L. minor* can absorb organic compounds

directly and utilize them as amino acid ingredients that can be used in their growth and development.

Treatment	Average	Notation
A	-1,06	a
B	-2,11	b
C	-2,57	b
D	-0,86	a

Table 1. Average Decreased BOD

Result of statistical test showed that treatment A has significant difference with both treatment C and B but has insignificant difference with treatment D.

Fresh Weight Change

Fresh weight changes were made to determine the difference between fresh weight at the beginning and end of the research. Changes in fresh weight are calculated using the formula from Juswardi (2010). Percentage changes in measured fresh weight during the research ranged from 43.33% -60.00%.




Figure 2. Fresh Weight Changes

Treatment B with the addition of *L.*

minor biomass of aquatic plants as much as 3.0 grams occurred an increase in fresh weight on average by 50% from the beginning to the end of the study. *L. minor* used in treatment B did not experience a significant death but increased instead. An increase in fresh weight was caused by the presence of light which plays an important role in the process of photosynthesis (Sudjadi 2005). Treatment C and D experienced a decrease in fresh weight at the end of the study. The decrease in fresh weight in treatments C and D was due to a high death rate of *L. minor* and a dense population (Akbar *et. al* 2011).

Treatment	Average	Notation
B	50	a
C	50,60	a
D	46,25	a

Table 2. Average reduction in fresh weight

Statistical test results showed that each treatment was not significantly different from the other treatments. Treatment D is the best treatment with a low percentage value. This showed that *L. minor* in treatment D can absorb organic compounds in tofu liquid waste well without inhibiting its growth rate (Hasan 2009).

Daily Growth Rate

Daily Growth Rate is an illustration of the increase in fresh weight during the study. The measured daily growth rate during the study ranged from 9% - 25%.




Figure 3. Daily Growth Rate

Based on the results of the study, the treatment B, C and D experienced an increase in the daily growth rate on day 5. According to Landesman *et. al* (2005), the growth and development of *Lemna sp.* at optimal conditions it could be doubled in two days. All treatments experienced growth marked by an increase in fresh weight. The presence of nutrients and the resistance of *L. minor* to tofu liquid waste caused *L. minor* to experience growth and increase in fresh weight.

Day 10 on treatment B, C and D experienced a significant decrease in fresh weight. The color of *L. minor* leaves in all three treatments has started to turn yellowish and most of it had turned pale. In addition, the size of the plant gets smaller and the leaves fall out so that they accumulate underneath and settle. It was because the resistance of *L. minor* to pollutants decreases and inhibits its growth (Tanghu *et al.* 2011).

Treatment	Average	Notation
B	10,00	a
C	16,67	b
D	23,67	c

Tabel 3. Average Daily Growth Rate

Statistical test results showed that each treatment was significantly different from the other treatments. Treatments C and D are the best treatment with a high percentage value. It showed that *L. minor* in treatments C and D utilize organic compounds in the waste so that they experienced growth and increase in fresh weight (Landesman *et. al* 2005).

Temperature

Temperature is one of the parameters in determining the quality of a waters. During the research temperature varies between 21,6⁰C – 22,2⁰C. The highest temperature range was in the D2 treatment at 22,2⁰C while the lowest temperature was in the B3 treatment at 21,6⁰C.




Figure 4. Temperature

Based on the graph above, *L. minor* in all treatments could grow at temperatures ranging from 21,6⁰C – 22,2⁰C, which means that the temperature in the research media was still within the tolerance range. Treatment A and B experienced a significant temperature drop from the beginning to the end of the study. Treatments C and D increased on the 10th day and then decreased on the 15th day. A decrease in temperature occurred almost every time the measurement eventhough not too

significant. According to Leng *et. al* (1994), *L. minor* is able to grow well in temperatures of 6⁰C - 33⁰C. This statement is in accordance with the results of the study that in research media *L. minor* aquatic plants grow in a temperature range of 21,6⁰C - 22,2⁰C.

pH

The pH value of the measured tofu liquid waste varies from 4,86 – 7,58. The highest pH range was obtained in the C3 treatment while the lowest pH value was obtained in the A1 treatment.




Figure 5. pH

The measured pH range on day 5 and day 10 for treatments A, B and C decreased while treatment D was increased. It was due to the decomposition of organic compounds, namely protein into alkaline amonia. Decomposition of organic compounds was is carried out by aerobic bacteria to produce NH₃. According to Alaerts *et al.* (1987) in Dhahiyat (1990), the results of oxidation from ammonification will form CO₂, H₂O and NH₃. The measured pH range on the 5th and 10th days on average did not eppropriate the standards set for the liquid waste quality standard that is 6-9.

The measured pH of tofu liquid waste on the 15th day ranged from 7,54 –

7,58. The pH range in this observation was in accordance with the standards specified for the liquid waste quality standard in the Minister of Environment Regulation of the Republic of Indonesia No. 5 of 2014, namely 6-9.

DO

The dissolved oxygen (DO) concentration ranged from 5,49 – 6,27 mg / L. The highest DO concentration of 6,3 mg / L was in the A3 treatment while the lowest DO concentration was in the D2 treatment.




Figure 6. DO

Overall treatment A had the highest DO concentration compared to the other three treatments. That was caused by the presence of *L. minor* in treatments B, C and D. According to Wijayanti (2007), the concentration of dissolved oxygen can be influenced by the process of respiration of aquatic biota and the process of decomposition of organic matter by microbes.

The measured DO value has accordanced the standard according to the Regulation of the Minister of Environment of the Republic of Indonesia No. 5 of 2014 which is more than 3 mg / L. According to Jenie et al. in Yusuf (2008), in waters the dissolved oxygen content of 3 - 5 mg / L has been fulfilled for the life of aquatic

organisms because in anaerobic conditions in the waters can be prevented so that life in it can take place.

Nitrate

Nitrate is the main form of nitrogen in natural waters and is the main nutrient that is useful for the growth of phytoplankton and other plants. Based on the results of the study, the measured nitrate concentration ranged from 0,326 to 0,445 mg / L.




Figure 7. Nitrate

Overall, each treatment has a high nitrate concentration at the beginning of the research but there was a significant decrease in the research of the 10th day and the 15th day. That was caused by *L. minor* which played a role in the phytoremediation process as a biological liquid waste treatment plant (Lisnasari 1995).

According to Zimmo et. al (2005), *L. minor* is effective in fixing nitrogen in polluted waters. *L. minor* can utilize nitrates dissolved in the research media through nutrient uptake at its roots.

Phosphate

Based on the results of the study, the measured phosphate concentration

ranged from 0,662 – 1,341 mg / L.




Figure 8. Phosphate

The function of phosphate in plants for root growth, strengthening the stem and accelerating the growth of leaves. According to Thompson and Troeh (1978) in Aleel (2008), phosphate is needed by plants for the formation of cells in the root tissue and buds that are growing and strengthens the stem so that it does not easily fall into natural ecosystems.

Almost every week the concentration of phosphate in each treatment has decreased. This is because *L. minor* is a remediation plant that is able to assimilate organic compounds and inorganic compounds contained in tofu liquid waste. According to Romimohtarto *et. al.* (2005), the decrease in phosphate is also caused by high light intensity which influences the process of photosynthesis so that plants utilize phosphate directly in the form of phosphorus.

BOD

BOD is the oxygen requirements needed by microorganisms in the process of decomposition of organic matter. The measured BOD concentration during observation ranged from 8,92 – 15,67 mg / L.




Figure 9. BOD

Each treatment decreased on the 10th day except treatment A. This was due to the high organic compounds which had to be decomposed in the research media and the lack of bacterial population that could decompose the organic material. BOD concentrations on the 15th day in treatments B, C and D decreased but in treatment A experienced a significant increase. The occurrence of the decrease caused by bacteria available in the media can decompose organic matter. According to Datko *et al.* (1985), *L. minor* can absorb organic compounds directly and utilize them as amino acid ingredients that can be used in their growth and development. It showed that the BOD content in the tofu liquid waste already accorded the liquid waste quality standard requirements specified in the Regulation of the Minister of Environment of the Republic of Indonesia No. 5 of 2014 which is below 50 mg / L.

CONCLUSION

The results showed that treatment C, namely *L. minor* biomass of 5,0 gr / L of wastewater was the best treatment with a reduction in BOD of 2.56 mg / L.

SUGESTIONS

L. minor water plants can be used as phytoremediators in tofu liquid waste and applied to the Wastewater Treatment Plant (WWTP) to reduce excess BOD concentrations before being polluted into water bodies.

REFERENCES

- Agnes. 2013. Kajian Efektifitas Tanaman Air Lemna minor dan Hydrilla Verticillata dalam Mereduksi BOD dan COD sebagai Upaya Perbaikan Kualitas Limbah Cair Industri Tahu. Institut Teknologi Nasional Malang : Jurusan Teknik Lingkungan.
- Akbar, B., M. Muryono, F. Hendrayana. 2011. Pengaruh Kerapatan Terhadap Pertumbuhan dan Produktivitas Tanaman Tembakau (*Nicotiana tabacum*) Varietas Serumpung dan Semboja. Hal 9.
- Aleel, K. G. 2008. Phosphate Accumulation in Plant : Signaling, Plant Physiol. 148:3-5
- Anonim. 2011. *Laporan Final Penyusunan Feasibility Studi Sentra Industri Tahu Cibuntu, Studi di Kecamatan Babakan Ciparay dan Kecamatan Bandung Kulon, Kota Bandung*. Bandung : Dinas Tata Ruang dan Cipta Karya.
- Dhahiyat, Y. 2011. Buku Ajar : Ekologi Perairan. Universitas Padjadjaran. Bandung.
- Endro S., Sri S. dan Nurdiansyah. 2010. Pengaruh Tanaman Rumput Bebek (*Lemna minor*) Terhadap Penurunan BOD dan COD Limbah Cair Domestik. Jurnal Presipitasi 7 (1).
- Hasan, M. dan Chakrabarti, R. 2009. Use of algae and aquatic macrophytes as feed in small-scale aquaculture. FAO Fisheries and Aquaculture Technical Paper No. 531. FAO : Rome.
- Husin, A. 2003. Pengolahan Limbah Cair Industri Tahu Menggunakan Biji Kelor (*Moringa oleifera*) Sebagai Koagulan. Laporan Penelitian Dosen Muda. Fakultas Teknik Universitas Sumatra Utara : Medan.
- Juhaeti, Titi, N. Hidayati, F. Syarif dan S. Hidayat. 2009. Uji Potensi Tumbuhan Akumulator Merkuri untuk Fitoremediasi Lingkungan Tercemar Akibat Kegiatan Penambangan Emas Tanpa Izin (PETI) di Kampung Leuwi Bolang, Desa Bantar Karet, Kecamatan Nanggung, Bogor. Jurnal Biologi Indonesia 6 (1) : 1-11
- Kuncoro, Mudjarat. 2003. Metode Riset untuk Bisnis & Ekonomi. Erlangga. Jakarta.
- Leng, R. A, Stambolie, J. H, and Bell, R. 1994. Duckweed a potential high protein feed resource for domestic animal and fish. AAAP Animal Science ke-7. Bali.
- Lisnasari, S.F. 1995. Pemanfaatan Gulma Air (Aquatic Weeds) Sebagai Upaya Pengolahan Limbah Cair Industri Pembuatan Tahu [Thesis Master]. Program Pascasarjana USU : Medan.
- Mkandawire dan Dudel. 2007. Are Lemna sp. effective phytoremediation agents?. Bioremediation, Biodiversity and Bioavailability Journal (1) : 56-71.
- Romimohtarto, K dan Juawana, K. 2009. Biologi Laut. Djambatan : Jakarta.
- Sudjadi. 2005. Pengaturan cahaya lampu sebagai fotosintesis phytoplankton buatan dengan menggunakan mikrokontroler At89s52.

Tanghu, B. V., S. R. S. Abdullah, H. Basri, M. Idris, N. Anuar dan M. Mukhlisin. 2011. A Review on Heavy Metals (As, Pb and Hg) Uptake By Plants Through Phytoremediation. International Journal of Chemical Engineering.

Wijayanti, H. 2007. Kajian Kualitas Perairan di Pantai Kota Bandar Lampung Berdasarkan Komunitas Hewan Makrobenthos [Thesis].

Universitas Diponegoro : Manajemen Sumberdaya Pantai.

Yusuf, Guntur. 2008. Bioremediasi Limbah Rumah Tangga dengan Sistem Simulasi Tanaman Air. Universitas Islam Makassar:FMIPA. Jurnal Bumi Lestari 8 (2) : 136-144

© GSJ