

Tap water from public drinking water supply	49	49.0
Ground water	10	10.0
Nile river	28	28.0
Other	13	13.0
Total	100	100.0

Table (6) shows that the most (49%) common resource of drinking water according to the participants was tap water from public drinking water supply, whereas (28%) of the participants use the Nile river, while only (10%) of them use groundwater. Moreover, (13%) of the respondents use other resources for drinking water.

Table (7) shows that most (68%) of participants have heard about Fluoride, mostly from schools/college (48%) or social media (16%). Furthermore, (55%) of participants were not sure what the source of Fluoride is, while (34%) thought that the main source of Fluoride is water.

It is shown that (54%) of participants thought that Fluoride has positive effects on human health and the majority (56%) of them was aware that Fluoride can be found in drinking water.

Most (50%) of participants knew that drinking fluoridated water can affect human health. On the contrary, (28%) of them were not sure about that. In addition, (38%) of participants believed that Fluoride can be added to public drinking water supplies, while (40%) of them were not sure.

Table (8) shows that most of the participants did not have spinal restriction in their back (back stiffness) (54%), numbness in the limbs and the trunk (73%), joint pain (64%), nor feel pain or stiffness in your neck (76%). Therefore, there were no significant effects of fluoride on bones seemed in the study area.

Table (9) indicates that the majority (92%) of the participants revealed that they do not have problems with their Kidneys. Moreover, most (66%) of them have never heard about the Water Defluoridation process. Only (26%) of them believed that drinking water that contains a small amount of Fluoride can lead to Dental Caries, while (40%) of them were not sure. According to the results, (48%) of the participants have Dental Caries and most of the participants brushed their teeth with fluoride toothpaste (69%). In addition, the majority of the respondents eat a lot of sugars and starches and do not clean their teeth well afterward (59%). Based on the results, the most common dental problem was a painless brown stain in one pair of teeth or more (11%) compared to having a painless brown stain with a corroded appearance in one pair of teeth or more (6%). In contrast, the highest percentage was 68%, in which the participants revealed that they do not have those dental problems and 32 % of them had problems during childhood. The results showed 9% of the participants believed that the reason for getting those kinds of dental problems is drinking water. However, 36% of them had a different opinion. They thought that drinking tea or coffee can cause dental problems. Moreover, 55% of the participants were not sure. The majority of the participants (82%) have gone to the dental clinic and (18%) have not gone. Furthermore, (3%) of the participants who went to any dental clinic, have done a fluoride blood test, while (89%) have not done fluoride blood test and (8%) were not sure.

Table 7: Distribution of participants according to knowledge about Fluoride

		Count	Percentage
Have you ever heard about Fluoride?	Yes	68	68 %
	No	20	20.0%
	Not sure	12	12.0%
What was your source of information about Fluoride?	Schools or College	48	48.0%
	Social media	16	16.0%
	Other	15	15.0%
	None	21	21.0%
What is the main source of Fluoride?	From food	11	11.0%
	From water	34	34.0%
	Not sure	55	55.0%
Does Fluoride have positive effects on human health?	Yes	54	54.0%
	No	8	8.0%
	Not sure	38	38.0%
Do you know that Fluoride can be found in drinking water?	Yes	56	56.0%
	No	22	22.0%
	Not sure	22	22.0%
Do you know that drinking fluoridated water can affect human health?	Yes	50	50.0%
	No	22	22.0%
	Not sure	28	28.0%
Do you believe that Fluoride can be added to public drinking water supplies?	Yes	38	38.0%
	No	22	22.0%
	Not sure	40	40.0%

Table 8: Distribution of participants according to Fluoride’s effects on bones:

		Yes	No	Not sure
Do you have spinal restriction in your back (back stiffness)?	Frequency	28	54	18
	Percentage	28%	54%	18%
Do you have numbness in the limbs and the trunk?	Frequency	15	73	12
	Percentage	15%	73%	12%
Do you have joint pain?	Frequency	32	64	4

	Percentage	32%	64%	4%
If you try to touch your chest with your chin, do you feel pain or stiffness in your neck?	Frequency	17	76	7
	Percentage	17%	76%	7%

Table 9: Distribution of participants according to effects of fluoride on kidneys and teeth

		Count	Percentage
Do you have any problem with your Kidneys?	Yes	4	4.0%
	No	92	92.0%
	Not sure	4	4.0%
Have you ever heard about Water Defluoridation (Removing excess Fluoride from drinking water by filtering with a strong adsorbent like alumina or carbon)?	Yes	18	18.0%
	No	66	66.0%
	Not sure	16	16.0%
Do you believe that drinking water that contains a small amount of Fluoride can lead to Dental Caries?	Yes	26	26.0%
	No	34	34.0%
	Not sure	40	40.0%
Do you have Dental Caries?	Yes	48	48.0%
	No	49	49.0%
	Not sure	3	3.0%
Do you brush your teeth with fluoride toothpaste?	Yes	69	69.0%
	No	13	13.0%
	Not sure	18	18.0%
Do you eat a lot of sugars and starches and do not clean your teeth well afterwards?	Yes	59	59.0%
	No	41	41.0%
Do you have one of the following dental problems?	Painless white spots in one pair of teeth or more	8	8.0%
	Painless white areas in one pair of teeth or more	7	7.0%
	Painless brown stain in one pair of teeth or more	11	11.0%
	Painless brown stain with corroded appearance in one pair of teeth or more	6	6.0%
	None	68	68.0%

When did these dental problems occur?	During Childhood	32	32.0%
	I do not have these problems	68	68.0%
What do you think is the reason for these kinds of dental problems?	Drinking Water	9	9.0%
	Drinking Tea or Coffee	36	36.0%
	Not sure	55	55.0%
Have you ever gone to a dental clinic?	Yes	82	82.0%
	No	18	18.0%
If you went to any dental clinic, have you ever done a Fluoride blood test?	Yes	3	3.0%
	No	89	89.0%
	Not sure	8	8.0%

Discussion

As we know, observational studies and epidemiological studies habitually constitute a valuable provider that is used for pointing a weight of evidence for an accurate approach (A.K. Gupta, 2016). The study assessed the public awareness on the effect of drinking water that contains fluoride on human health. (Samuel Y Ganyaglo, 2019) .According to the study, a high percentage of participants knew about fluoride and this is because schools and colleges gave the way for students to gain information about fluoride. Furthermore, the study showed that participants are aware that fluoride can be found in drinking water but unfortunately they were not certain what the main source of fluoride is.

The majority of the participants thought that fluoride is beneficial to human health, and that was expected to be since it is well-established that fluoride plays a major role in increasing the strength of the tooth enamel by applying fluoride on the surface of the teeth, therefore, using fluoride in a wide range of dental products is considered as a significant mean to enhance dental health. On the contrary, the participants lack information about the standard amount of fluoride this is crucial to their body.

The chemical analysis of fluoride samples obtained from the 11 locations from Khartoum and Omdurman revealed that all values were less than the WHO standard value of fluoride that exists naturally in drinking water. In addition, the results showed that almost all of the values of fluoride

found in drinking water were less than the recommended concentration for inhibiting tooth decay which is 0.7 mg/L (J.Farwell, 2003). This is indicated in the results of the questionnaire as the majority of the participants enrolled showed that they are suffering from tooth decay. Therefore, there is a high possibility that the people in those areas are affected by lower levels of fluoride contained in their drinking water.

According to the results, a high percentage of participants said that they did not have spinal restriction in their back (back stiffness), numbness in the limbs and the trunk, joint pain, nor feel pain or stiffness in their neck. Therefore, the process of bone turnover is homeostasis controlled by osteoblasts and osteoclasts. Consuming huge amounts of fluoride can interfere with this balance, affecting osteoblasts and osteoclasts and ultimately leading to skeletal fluorosis (ChenYANG, 2017). Since the values of fluoride received from the chemical analysis were lower than the value that can cause skeletal fluorosis, there were no significant effects of fluoride on bones seemed in the study area.

The results obtained indicated that most of the participants do not suffer from kidney problems. The reason behind that is the low amount of fluoride found in the samples of drinking water. On the contrary, a previous literature review suggested that when getting a high amount of fluoride, there could be high risk of harming kidneys. A study was done in 2006 about the toxic effects of fluoride on kidney function and histological structure. The results showed that extreme fluoride exposure caused multiple renal histological changes as well as increased renal cell apoptosis in the study groups. Moreover, the study revealed that the enzymes of kidney tissues such as lactate dehydrogenase (LDH) activity were significantly raised, whereas alkaline phosphatase (AKP) activity was decreased. Furthermore, there was an increase in urea nitrogen (UN), an elevation in creatinine (Cre), and a reduction in Na⁺. These findings show that chronic excessive fluoride

exposure is toxic to kidney structure and its function. Therefore, the fluoride in drinking water should be in a moderate amount in order not to damage kidneys. (Xiu-An Zhan, 2006)

A high percentage of participants were not aware of defluoridation. This pattern was expected given the poor education and low level of knowledge of people about fluoride. Drinking water that contains an excessive amount of fluoride leads to an accumulation of fluoride in the body (toxicity of fluoride). Consequently, this causes a variety of adverse effects on human health involving kidney diseases, dental fluorosis, skeletal fluorosis, and influences on IQ (Shabiimam M. A, 2017). Various defluoridation techniques have been developed to eliminate such an extreme amount of fluoride. Fluoride removal can be done by the application of some techniques such as precipitation, adsorption, ion exchange, and deionization (Elhessin, 2004). So, to ensure drinking safe water, some measures such as popularizing defluoridation techniques and optimizing water supply strategies need to be implemented (Hui Jia, 2019)

The results showed that the majority of participants have no idea that fluoride can damage their teeth and they only thought that drinking tea and coffee is the main reason. Furthermore, the majority of the participants went to dental clinics and a few of them have done fluoride blood tests. Important findings are the fact that people's knowledge about fluoride is very poor. Since dentists and science teachers are well educated about fluoride, they need to emphasize the significance of fluoride for human health for people and students in schools and colleges by increasing awareness about fluoride. Also, during their study, they should concentrate on practical chemical analysis of fluoride to have accurate information about fluoride levels that are required to inhibit health problems. As a result, conducting the Fluoride tests in patients is very crucial. The reason behind that is that while fluoride is a naturally occurring compound, it can still possess hazardous effects when taken in large amounts or small amounts. Dentists should properly

counsel their patients, parents, and children on correct tooth brushing behaviours. (Ritu Bansal, 2012).

Conclusion

In conclusion, it is important to raise awareness about diverse chemical analyses of fluoride amongst students and people. Moreover, educating them about the positive and negative effects of fluoride in drinking water is required. Also, carrying out the fluoridation process (adding low concentrations of fluoride to drinking water) to raise the fluoride level in drinking water is considered as a solution to prevent diseases caused by the decline in fluoride levels in the human body. In case of having high levels of fluoride, using the defluoridation process can help to reduce it.

Acknowledgment

I would like to express my thanks and gratitude the teams of central Lab in Khartoum and to International College of Engineering and Management for their support and the facilities provided

References

- A.K. Gupta, S. A. (2016). *Fluoride in Drinking Water*. CRC Press.
- Bo, Z., Mei, H., Yongsheng, Z. et al. Distribution and Risk Assessment of Fluoride in Drinking Water in the West Plain Region of Jilin Province, China. *Environmental Geochemistry and Health* 25, 421–431 (2003). <https://doi.org/10.1023/B:EGAH.0000004560.47697.91>
- Cafasso, J. (2019). What Is Fluoride, and Is It Safe? *Healthline*.
- ChenYANG, Y. H. (2017). Treatment and Prevention of Skeletal Fluorosis. *Biomedical and Environmental Sciences*, 147-149.
- Elhessin, H. A. (2004). *Assessment of Fluoride Concentration in Drinking Water in Gedaref Town and its Uptake by Some Vegetable Plants*. Khartoum: University of Khartoum.
- F. M. Fordyce, K. V. (2007). A health risk assessment for fluoride in Central Europe. *Environmental Geochemistry and Health*, 83-102.
- Fluoridation, U. D. (2015). U.S. Public Health Service Recommendation for Fluoride Concentration in Drinking Water for the Prevention of Dental Caries. *Public Health Reports*, 318–331.
- Gupta, A.K. Ayoob S. (2016). Fluoride in the Environment and Its Toxicological Effects. Book chapter: Fluoride in Drinking Water, pp.76-103.

- Gupta, A.K. Ayoob S. (2016). Stress Effects of Fluoride on Humans. Book chapter: Fluoride in Drinking Water, pp.68-75.
- Ibrahim, E. M. (2021). *The Relation between Risk and Level of Chemical Components in Drinking Water*. Khartoum: Lulu Press.
- Iftekhhar Ahmed, A. A. (2020). Frequency of dental fluorosis in population drinking water with high fluoride level in Thar. *Journal of the Pakistan Dental Association*, 5.
- J. Fawell, K. B. (2006). *Fluoride in Drinking-water*. London: World Health Organization.
- J. Farwell, E. O. (2003). *Fluoride in Drinking-water, Background document for development of WHO Guidelines for Drinking-water Quality*. Retrieved from World Health Organization: https://www.who.int/water_sanitation_health/dwq/chemicals/fluoride.pdf?ua=1
- Kanduti, D. S. (2016). FLUORIDE: A REVIEW OF USE AND EFFECTS ON HEALTH. *Materia socio-medica*, 133–137.
- Majumdar KK. (2011). Health impact of supplying safe drinking water containing fluoride below permissible level on fluorosis patients in a fluoride-endemic rural area of West Bengal. *Indian J Public Health*. Oct-Dec;55(4):303-8. doi: 10.4103/0019-557X.92411. PMID: 22298140.
- Ritu Bansal, K. A. (2012). Knowledge, Attitude and use of Fluorides among Dentists in Texas. *The Journal of Contemporary Dental Practice*, 375-379.
- Samuel Y Ganyaglo, A. G.-G.-A. (2019). Groundwater fluoride contamination and probabilistic health risk assessment in fluoride endemic areas of the Upper East Region, Ghana. *Chemosphere*, 233, 862-872.
- Shabiimam M. A, S. S. (2017). Status of Defluoridation Techniques in India: A Review Study. *NICMAR 2nd International Conference On Construction, Real Estate, Infrastructure and Project (CRIP) Management* (pp. 11-21). Pune, India: ResearchGate.
- Xiu-An Zhan, a. M.-R.-X. (2006). *TOXIC EFFECTS OF FLUORIDE ON KIDNEY FUNCTION AND HISTOLOGICAL STRUCTURE IN YOUNG PIGS*. Hangzhou, China: International Society for Fluoride Research.
- Hui Jia, H. Q. (2019). Fluoride Occurrence and Human Health Risk in Drinking Water Wells from Southern Edge of Chinese Loess Plateau. *international journal of environmental research and public health*, 18.
- Rahman, M. M., Doza, M. B., Siddiqua, M. T., Zahid, A., Md, A., & Islam, T. (2020). *Spatiotemporal distribution of fluoride in drinking water and associated probabilistic human health risk appraisal in the coastal region, Bangladesh*,. Science of The Total Environment.