

Sharma and Sharma (2002) observed that the increase in organic carbon content in treatments with combination of both organic and inorganic sources may be attributed to higher biomass addition to soil through crop residues.

Priyanka *et al.* (2013) in a study reported that the dry matter accumulation was highest with the application of FYM @ 20 t/ha and was followed by FYM application @ 10 t/ha and lowest dry matter accumulation was observed when no FYM was applied which may be due to the FYM provides better growing condition to plants by continuous supply of nutrients and improvement of soil properties.

The crop having 50% recommended dose of fertilizer (RDF) + 50% recommended dose of nitrogen (RDN) through mustard oil cake (MOC) and 75% RDF + 25% RDN through MOC + biofertilizer significantly increased dry matter accumulation (DMA) at initial and vital period of grain growth over those of 25% RDF + 75% RDN through MOC and 100% RDN through MOC (Mondal *et al.*, 2015).

Geetha *et al.* (2020) is a study reveals that among the treatments tested, treatment T9 (125% RDF + FYM @ 6 t ha⁻¹ + biofertilizers @ 12.5 kg ha⁻¹ + foliar spray of 19:19:19 (1%) at 45, 60 & 75 DAS) recorded the maximum dry matter production at harvest stage (55.35 g/ plant).

Rao *et al.* (2020) reported that nutrient management practices significantly affected dry matter accumulation with the application of T10 (125% RDF+ 25% V.C.) which was closely followed with T9 (125% RDF+ 25% FYM) in dry matter accumulation.

iii. Tiller Number

On sandy loam soils at Ludhiana (India), incorporation of wheat straw and FYM @ 67 and 12 t / ha⁻¹, respectively it gave higher tiller per unit area (Maskina *et al.*, 1987).

Relatively more number of tillers per hill was observed with the application of poultry manure @ 15 t ha⁻¹ compared to FYM application @ 5 t ha⁻¹ (Budhar *et al.*, 1991).

Increase in number of tillers per hill was observed with increasing levels of nitrogen (Shashikumar *et al.*, 1995).

Sarawgi and Sarawgi (2004) found that higher level of nutrients (50:50:40 kg NPK ha⁻¹ + nitrogen blended with FYM) recorded significantly higher number of tillers plant⁻¹ when compared to lower levels of nutrients (25:40:30 kg NPK ha⁻¹ either with or without blending with FYM).

Satyanarayana *et al.* (2002) showed that application of farmyard manure at 10 t / ha⁻¹ increased tiller numbers subsequently.

Umar *et al.* (2007) revealed that NPK + GM + Zn (Soil application) (T7) gave the highest number of tillers m⁻² (315.0), which differed from all other treatments except NPK +FYM+ Zn (soil) (T6) NPK + FYM + Zn (R.D) (T9), and NPK + GM + Zn (R.D) (T10) . The lowest number of tiller m⁻² (181.7) was recorded for control (T1) that differed significantly from all other treatments.

Arif *et al.* (2014) concluded that maximum number of effective tillers per hill was recorded with application of poultry manure @ 10 t ha⁻¹ + 50% RDF (75:45:30 kg NPK ha⁻¹). It was statistically at par with application of FYM @ 10 t ha⁻¹ and compost @ 5 t ha⁻¹ along with 50% RDF and significantly superior over 100% RDF.

With the application of FYM @ 20 t ha⁻¹ the number of tillers m⁻² were significantly influenced and the least was seen with the application of FYM @ 10 t ha⁻¹ (Priyanka *et al.*, 2013). Application of RDF (80:60:40 kg NPK ha⁻¹) along with FYM @ 5 t ha⁻¹ showed relatively higher number of tillers than 50% RDF with FYM @ 10 t ha⁻¹ (Wahland *et al.*, 2015).

Shalini *et al.* (2017) observed that application of vermicompost 1.5 t ha⁻¹ + brown manuring of dhaincha @ 25 kg ha⁻¹ + RDF resulted in significantly higher number of effective tillers m⁻² over RDF (120:60:40 kg NPK ha⁻¹).

Siddaram *et al.* (2017) studied the effect of FYM on growth of aerobic rice and documented that application of FYM 12.5 t + bio-digester liquid manure equivalent to 150 kg N ha⁻¹ produced significantly more number of tillers over control.

Adhikari *et al.* (2018) observed that total number of tillers plant⁻¹ decreased progressively with decreasing levels of nitrogen and became the least when no nitrogen was applied. Application of higher level of N (180 kg ha⁻¹) produced more number of tillers as compared to application of 150 kg of N ha⁻¹.

Kipgen *et al.* (2018) concluded that varying the nitrogen levels influenced the growth characteristics of rice application of 140 kg N ha⁻¹ produced significantly higher number of tillers m⁻² which was statistically at par with 120 kg N ha⁻¹.

More number of tillers hill⁻¹ was obtained with integrated use of organic and inorganic sources i.e., 50% RDF + 50% N through FYM which being significantly superior over control (Singh *et al.*, 2017). A similar significant effect was reported by Harikesh *et al.* (2017). Significantly higher number of tillers m⁻² was recorded with application of 75% NPK + 25% FYM which was statistically at par with 100% NPK application (Tomar *et al.*, 2018).

B. Yield Attributes

i. Number of panicles per square meter

In 2008 maximum number of panicles per square meter area was noted in plants treated with 2 ton/ha organic fertilizer (329.58), it was followed by organic fertilizer in combination with NPK (326.27) and 2.5 ton/ha organic fertilizer (324.81) and minimum of that was for 0.5 ton/ha organic fertilizer (310.57), whereas control plants showed 306.17 as an average panicle number in said unit area, but in 2009 the utmost panicles per square meter area was noticed for organic fertilizer 1.5 ton/ha + NPK (343.28), afterward plot treated by organic fertilizer 2 ton/ha (341.04) then plot treated with organic fertilizer 2.5 ton/ha (335.69), and the minimum amount of panicles per square meter area was for 0.5 ton/ha organic fertilizer (327.63), although control plants showed 308.50 panicles (Siavoshi *et al.*, 2011).

Hasanuzzaman *et al.* (2010) reported that maximum panicle m⁻² was obtained when 50% NPK was applied along with poultry manure @ 4 t / ha⁻¹.

Gohain (2014) reported that with increasing levels of nitrogen the number of panicles/m² significantly increased at highest level of 120 kg N/ha which was registered with the maximum number of panicles/m² (444.19) that might be due to increased accumulation of photosynthetic from the source to the sink.

ii. Panicle Length

Application of FYM at 5 t ha⁻¹ resulted in greater panicle length in rice compared to each counterpart treatment having the same NPK levels [T1 = control (N0P0K0); T2 = nitrogen (N)–phosphorus pentoxide (P₂O₅)–dipotassium oxide (K₂O) at 30:15:15 kg ha⁻¹; T3 = N–P₂O₅–K₂O at 30:15:15 kg ha⁻¹ + farmyard manure (FYM) at 5 t ha⁻¹ (oven-dry weight basis); T4 = N–P₂O₅–K₂O at 60:30:30 kg ha⁻¹; T5 = N–P₂O₅–K₂O at 60:30:30 kg ha⁻¹ + FYM at 5 t ha⁻¹; T6 = N–P₂O₅–K₂O at 90:45:45 kg ha⁻¹; T7 = N–P₂O₅–K₂O at 90:45:45 kg ha⁻¹ + FYM at 5 t ha⁻¹] (Choudhary, Thakur, and Kumar 2007).

Hussain *et al.* (2012) reported that yield attributes viz., panicle length were significantly higher with application of RFD + poultry manure @ 20 t ha⁻¹, whereas grains panicle⁻¹ and panicles m⁻² were significantly higher with application of FYM @ 20 t ha⁻¹ + 75% recommended fertilizer dose.

Green manuring along with application of 50% RN (40 kg N ha⁻¹) resulted in maximum panicle length (Aulakh *et al.*, 2016).

The effect of fertilizer doses on the length of panicle was non-significant. However, it was recorded highest (28.93 cm) in F2 (100% RDF + 5 t ha⁻¹ FYM (Apon *et al.*, 2018).

iii. Number of grains per panicle

Satyanarayana *et al.* (2002) observed that application of farmyard manure significantly improves number of filled grains per panicle. The number of filled grains was increased by 9% due to application of farmyard manure (72.30) when compared to no farmyard manure at 68.50.

Combination of FYM along with inorganic fertilizers increases panicle number per hill, grain number per panicle (Naing Oo, 2010).

C. Yield

i. Grain Yield

Acharya *et al.* (1998) pointed out that the application of N, P, and K fertilizer along with FYM increased the growth attributes, yield components and grain yield of rice compared with that of N, P and K through fertilizer alone.

Singh *et al.* (1998) reported that in a field experiment in 1994-1995 in Uttar Pradesh rice was given 07 t/ha FYM and 75, 100 and 125% recommended NPK fertilizer rate (100 kg N + 50 kg P₂O₅ + 50 kg K₂O/ha), grain yield were highest with 7.5 t FYM (5.59 t/ha) and the highest NPK rate (5.53 t).

Mann and Ashraf (2000) found that the maximum paddy yield could be obtained by either applying recommended dose of inorganic N fertilizer (80 kg ha⁻¹) or lowering its level to 40 kg ha⁻¹, when supplemented by green manure, dhaincha (*Sesbania aculeata*).

Jayabala and Kuppuswamy (2001) concluded that application of 50% N through vermicompost + 50% N through chemical fertilizers and biofertilizers led to record higher grain yield of rice.

Kumar *et al.* (2002) obtained comparable yields with the application of either 10 t FYM/ha or 2.5 t/ha vermicompost than that of 100% RDF (100:50:50 kg N P₂O₅ K₂O/ha) alone in scented rice (cv. Pusa Basmati 1).

Barik *et al.* (2008) concluded that combined application of 40% recommended dose of N as vermicompost + 60% recommended dose of N as urea produced the higher grain yield (52.7 q/ha) as compared to 100% recommended dose of N alone through urea at Mohanpur (W.B.) in transplanted *khari* rice.

Raul and Sarawagi (2005) reported that grain yield, N content and recommended dose of N blended with FYM and 100% RDN +5 t/ha FYM were better than other treatment on pooled data basis.

Chaudhary and Thakur (2007) reported that the application of FYM in conjunction with chemical fertilizers had stimulatory effect on yield irrespective of crops and season and the highest rice grain yield was registered when 50% N was supplied through green manure in conjunction with 50% NPK through inorganic fertilizers.

In a study on the efficacy of different organic manure and inorganic fertilizer on the yield and yield attributes of Boro Rice. The maximum number of total grain plant⁻¹ (97.45 was recorded from application of 70% NPKS + 2.4 tonnes poultry manure ha⁻¹ (Hossain *et al.*, 2011).

Application of enriched compost @ 2.5 t/ha gave the highest grain yield (2.45 t/ha) and it also enhanced the other quality parameters of the grain (Bora *et al.*, 2013).

ii. Straw Yield

Ahmad *et al.* (2005) reported that the increase in straw yield was with higher nutrient levels.

Higher straw yields of rice was seen with the application of FYM + Neemcake equivalent to 90 Kg N/ha in both the years than that of other treatments including application of 100% RDF (90 Kg N + 40 Kg P₂O₅ + 40 Kg K₂O /ha) (Kumar *et al.*, 2005).

Gupta *et al.* (2006) reported that INM in maintaining the sustainable straw yield of rice-wheat system in rainy season without degradation of soil health under irrigated production system attained highest productivity of 33.12 kg ha⁻¹ compared to other fertilizer treatments i.e., 50%, 75% and 100% NPK with substitution of 50% N by green manure with sunhemp.

The increase in fertility level from 50 to 100 % of recommended dose increased the straw yield of the crop significantly (Mankotia, 2007).

Patnayak *et al.* (2007) revealed that 40 kg inorganic N (50% N dose) integrated with biofertilizers (Azotobacter, Azospirillum and Azolla) and 17.5 kg of P and 32 kg of K ha⁻¹ resulted in the highest straw yield (4.32 t ha⁻¹) of rice.

Virdia and Mehta (2010) studied the effect of integrated nutrient management in transplanted rice with treatments comprising various quantity of press mud, FYM and RDF. They found that straw yield was significantly higher with integrated nutrient management (press mud @ 20 t ha⁻¹ + RDF), which remained at par with press mud @ 15 t ha⁻¹ + RDF or FYM @ 10 t ha⁻¹ + RDF.

Mehdi *et al.* (2011) found that different combinations of organic manures with chemical fertilizers increased straw yield significantly over application of organic manures alone. Among different combinations, Sesbania at 20 ton ha⁻¹ + 75% recommended dose proved to be the best combination followed by Sesbania 20 t ha⁻¹ + 50% R.D.

Balasubramanian and Wahab (2012) observed that straw yields were favorably influenced by combined application of inorganic fertilizers and organic manures.

In a study conducted by Singh *et al.* (2012) reported that application of 100% RDF through inorganic fertilizers being on par with 50% RDF as inorganic fertilizers + 50% RDN as farm yard manure but produced significantly straw yield (2.23 t ha^{-1}) over rest of the fertility treatments.

Sharma and Subehia (2014) revealed that through continuous substitution of 50% N through green manure in rice produced maximum straw (7.37 t ha^{-1}) yield was 16.8 percent higher over 100% NPK added through chemical fertilizers.

D. Nutrient uptake

Murali and Setty (2001) observed that application of vermicompost at 5 t ha^{-1} combined with NPK at $150-75-75 \text{ kg ha}^{-1}$ recorded maximum total nitrogen uptake (168 kg ha^{-1}) as compared to no vermicompost treatment (1152 kg ha^{-1}).

Mhaskar and Thorat (2005) in a study on the effects of different nitrogen levels (0, 40, 80 and 120 kg N/ha) on the N, P and K uptake of scented rice cultivars reported that N, P and K uptake was significantly influenced by the cultivar. Indrayani registered the maximum uptake of N, P; Sugandha was significantly inferior to all the cultivars. The different levels of N had significant effect in augmenting the uptake of N, P and K nutrients. Application of 120 kg N/ha recorded significantly higher N, P and K uptake in rice compared to the rest of the N levels.

Maximum mean nitrogen uptake (94.9 kg ha^{-1}) was recorded under combined use of farm yard manure and poultry manures. Incorporation of organic manures caused improvement in organic carbon and available nitrogen content of soil after crop harvest as compared to control (Kumar *et al.*, 2006).

Banik and Sharma (2008) reported that nutrient uptake in rice under the integrated nutrient management system was greater due to greater biomass production and greater nutrient mineralization from organic sources. Increase in fertilizer levels significantly increased the N uptake in rice (Choudhary and Suri 2009). Similar trends were also observed for P and K uptake in rice crop through crop residue management and chemical fertilizers as well (Choudhary and Suri 2009).

Various integrated nutrient management affected significantly nutrient uptake by rice. The maximum NPK uptake was recorded by the application of 75% NPK + 25% FYM through inorganic and organic fertilizer which was at par with 100% NPK. Availability of nutrients might be sufficient & it led to higher nutrient uptake. Minimum nutrient uptake was recorded where 50% NPK + 50% FYM) was applied. It might be due to inadequate availability of nutrient. The results are in close proximity of Talathi *et al.* (2009).

Yadav *et al.* (2010) observed the efficacy of substituting fertilizer N at different proportions (25%, 50% and 75% of total N) with organic N sources (i.e., farm yard manure (FYM), green leaf manure (GLM), poultry manure and BGA) on nutrient uptake (NPK) of rice variety Sarju 52. In general, the maximum uptake of the nutrients was obtained with the application of 25% N through green manure + 75% through inorganic urea.

Siddaram *et al.* (2010) revealed that significantly higher nitrogen, phosphorus and potassium uptake ($124.2, 30.6$ and 93.9 kg ha^{-1} , respectively) registered with recommended dose of fertilizer ($100:50:50 \text{ kg N:P:K ha}^{-1}$) + 10 tonnes of FYM ha^{-1} .

Weijabhandara *et al.* (2011) reported that application of 75% RDF + biofertilizers resulted in significantly higher grain yield, uptake of N, P, K and Zn by grains and residual available N, P, Zn compared to other treatments.

Acharya *et al.* (2012) revealed that nutrient uptake of rice was highest due to integrated nutrient application than that of inorganic nutrients alone, whereas lowest value was observed with control plot where no nutrient was applied.

The higher uptake of N, P and K in crop (106.44, 16.32 and 111.86 kg/ha respectively) with post emergence application of bispyribac sodium @ 25 g a.i. kg/ha and this treatment was at par to pre-emergence application of bensulfuron methyl @60g a.i/ha + pretilachlor @600g a.i/ha (Manjunatha *et al.*, 2012).

Ranjitha and Reddy (2013) reported that higher nitrogen uptake by grain and straw (56.0 and 26.7 kg ha⁻¹ respectively) was observed with the application of FYM @ 10 t ha⁻¹ + 100 percent RDF but was comparable with the treatment of 100 percent RDF alone. Similarly, highest P and K uptake (16.6 kg ha⁻¹ and 10.3 kg ha⁻¹ P; 18.9 and 127.1 kg ha⁻¹ K) by grain and straw was obtained by FYM @10 t ha⁻¹ + 100 percent RDF, followed by 100 percent RDF and lowest was with FYM @ 10 t ha⁻¹.

The rice crop had higher NPK uptake by grain and straw being recorded with System of Rice Intensification (SRI) method as compared to the other treatment (Transplanting (S1), Drum Seeded (S3), Direct Seeded (S4) and three integrated management practices [100% NPK (F1), 75% NPK + 25% FYM (F2), 50% NPK + 50% FYM (F3)], while lower NPK uptake by direct seeded method by grain and straw, respectively. The higher nutrient uptake was attributed to the higher grain and straw yield (Tomar *et al.*, 2018).

E. Soil nutrient status

Vennila and Jayanthi (2007) observed that application of 75% RDFN + organic manure resulted in higher soil available organic carbon, nitrogen and phosphorus. Application of 75% RDFN along with 25% N as organic manure to preceding wet seeded rice had significant residual effect on yield and nutrient uptake of succeeding green gram.

Available N, P and K status of soil was also significantly affected by various treatments. Plots supplied with 5 t ha⁻¹ Gliricidia over and above farmers' practice had the highest amount of available N (295.7 kg ha⁻¹) which was significantly higher than all other treatments (Das *et al.*, 2009). Again, application of 2.5 t ha⁻¹ Gliricidia along with 50% RDF and bio-fertilizers, being next best treatment, significantly enhanced soils available N-status. This could be ascribed to higher N content of Gliricidia and its role in stimulating microbial activity, contribution of Azotobactor by fixing atmospheric N (Rao, 2007).

Ghosh *et al.* (2012) reported significant increase in soil nutrient availability with the application of farm yard manure (FYM @ 7.5t/ha), paddy straw (PS @ 10 t/ha) and green manure (GM @ 8 t/ha) along with inorganic fertilizer. Both microbial biomass C and mineralizable C as well as yield of *kharif* rice were increased with the addition of the organic inputs.

References:

- FAOSTAT. 2018. Crops/Regions/World list/Production Quantity (pick lists), Rice (paddy), <http://www.fao.org/economic/est/publications/rice-publications/rice-market-monitor-rmm/en/>
- FAOSTAT. 2016. Retrieved from. <http://faostat3.fao.org/download/Q/QC/E>.
- Agricultural Statistics at a Glance. 2018. Directorate of Economics and Statistics, Ministry of Agriculture and Farmers Welfare, Government of India.
- UNSD. 2015. UN projects world population to reach 8.5 billion by 2030, driven by growth in developing countries. <https://www.un.org/sustainabledevelopment/blog/2015/07/un-projects-world-population-to-reach-8-5-billion-by-2030-driven-by-growth-in-developing-countries/>
- Roy, R.N. 1995. FAO, Rome
- Sannathimmappa, H.G.; Gurumurthy, B.R.; Jayadeva, H.M.; Rajanna, D. and Shivanna, M.B. 2015. Effect of Paddy Straw Based Integrated Nutrient Management Practices for Sustainable Production of Rice. *IOSR Journal of Agriculture and Veterinary Science*, 8: 74-77.
- Dahiphale, A.V.; Giri, D.G.; Thakre, G.V. and Gin, M.D. 2003. Effect of integrated nutrient management on yield yield contributing parameters of the scented rice. *Annual Review of Plant Physiology*, 17(1): 24-26.
- Baradhan, G. and Kumar, S.M. 2018. Studies on the effect of integrated nutrient management in the yield of maize (*Zea mays*). *Plant Archives*, 18(2): 1795-1800.
- Rao, P.M., Prasad, P. R. K., Jayalakshmi, M. and Srihari Rao, B. 2017. Effect of organic and inorganic sources of nutrients on NPK uptake by rice crop at various growth periods. *Research Journal of Agriculture and Biological Science*, 8: 64-69.
- Upadhyay, V.B., Jain, V., Vishwakarma, S.K. and Kumhar, A.K. 2011. Production potential, soil health, water productivity and economics of rice (*Oryza sativa*)-based cropping systems under different nutrient sources. *Indian Journal of Agronomy*, 56(4): 311-16.
- Heffer, P. 2013. Assessment of Fertilizer Use by Crop at the Global Level, 2010 – 2010/11, International Fertilizer Industry Association (IFA), Paris, France.
- Ram, M.S., Shankar, T., Maitra, S., Adhikary, R. and Swamy, G.V.V.S.N. 2020. Productivity, nutrient uptake and nutrient use efficiency of summer rice (*Oryza sativa*) as influenced by integrated nutrient management practices. *Crop Research*, 55(3-4): 65-72.
- Khurshed, S., Arora, S. and Ali, T. 2013. Effect of Organic Sources of Nitrogen on Rice (*Oryza sativa* L.) and Soil Carbon Pools in Inceptisols of Jammu. Columbia International Publishing, *International Journal of Environmental Pollution and Solutions*, 1: 17-21.
- Mhaske NS, Borkar SL and Rajegire HJ. 1997. Effects of nitrogen levels on growth, yield and grain quality of rice. *Journal of Soils and Crops*, 7(1): 83-86.
- Maiti S, Saha M, Banerjee H, Pal S. 2006. Integrated nutrient management under hybrid rice (*Oryza sativa* L.) hybrid rice cropping sequence. *Indian Journal of Agronomy*, 51(3): 157-159.
- Singh, G., Singh, S. and Singh, R.K. 2012. Effect of fertility management on yield and economics of traditional scented rice varieties in lowlands. *Annals in Plant and Soil Research*, 14(1): 1-4.
- Singh RK, Kumar A and Kaleem MJS. 2013. Yield maximization of Hybrid Rice (*Oryza sativa* L.) through Integrated Nutrient Management. *Journal of Progressive Agriculture*, 4(1): 18-22.
- Mahmud AJ, Shamsuddoha ATM, Haque MN. 2016. Effect of Organic and Inorganic Fertilizer on the Growth and Yield of Rice (*Oryza sativa* L.). *Natural Science*, 14(2): 45-54.

- Shankar, T. A. N. M. O. Y., Maitra, S. A. G. A. R., Sairam, M., & Mahapatra, R. 2020. Influence of integrated nutrient management on growth and yield attributes of summer rice (*Oryza sativa* L.). *Crop Research*, 55(1): 1-5.
- Kumar, J., Yadav, M. P. and Kedar, P. 2010. Production potential of hybrid rice (*Oryza sativa* L.) as influenced by integrated nutrient management. *Crop Research*, 39: 20-23.
- Babar, S. and Dongale, J. H. 2011. Effect of integrated use of inorganic and organic manure on yield and monetary returns of mustard-cowpea-rice cropping sequence in lateritic soils of Konkan. *Journal of Soils and Crops*, 21: 225-33.
- Singh, N. P., Singh, M. K. S. S., Tyagi, S. and Singh, S. S. 2018. Effect of integrated nutrient management on growth and yield of rice (*Oryza sativa* L.) *International Journal of Current Microbiology and Applied Science*, 7: 3671-81.
- Ajay kumar, R., & Sivakumar, S. D. 2020. Effect of Integrated Nutrient Management on Growth, Physiological, Nutrient Uptake, Root and Yield Parameters of Transplanted Lowland Rice. *Current Journal of Applied Science and Technology*, pp. 35-43.
- Naveen Kumar, G., P. K. Singh, R. K. Naresh, M. Sharath Chandra and Akash Kumar. 2021. Effect of Planting Techniques with Organic and Inorganic Nutrient Management on Growth, Yield Attributes and Productivity of Rice (*Oryza sativa* L.) under Rice-Wheat Cropping System. *International Journal of Current Microbiology and Applied Science*, 10(01): doi: <https://doi.org/10.20546/ijcmas.2021.1001.xx>
- Ghosh, B. C., Ghosh, R., Mitra, B. N., Mitra, A and Jana, M. R. 1994. Effect of organic and inorganic fertilization on growth and nutrition of rice and fish in a dual culture system. *Journal of Agricultural Science*, 122 (12): 41-45.
- Mandal, S. S., Mandal, T. K., Dandapat, S and Sarkar, S. 1994. Effect of sulphur bearing fertilizers in conjunction with FYM on growth, productivity and nutrient uptake of rice. *Indian Journal of Agronomy*, 29: 116-118.
- Singh, Y.; Singh, B.; Khera, T.S. Meelu, O.P. 1994. Integrated Management of Green Manure, Farmyard Manure, and Nitrogen Fertilizer in a Rice-Wheat Rotation in Northeastern India. *Arid Soil Research and Rehabilitation*, 8 (2): 199–205.
- Chung, R.S.; Wang, C.H.; Wang, C.W.; Wang, Y.P. 2000. Influence of Organic Matter and Inorganic Fertilizer on the Growth and Nitrogen Accumulation of Corn Plants. *Journal of Plant Nutrition*, 23 (3): 297–311.
- Sharma SK and Sharma SN. 2002. Integrated nutrient management for sustainability of rice-wheat cropping system. *Indian Journal of Agricultural Science*, 72: 573-576.
- Priyanka, G., Sharma, G.D., Rana, R. and Lal. B. 2013. Effect of integrated nutrient management and spacing on growth parameters, nutrient content and productivity of rice under system of rice intensification. *International Journal of Research in Biosciences*, 2(3): 53-59.
- Mondal, S., Mallikarjun, M., Ghosh, M., Ghosh, D. C., & Timsina, J. 2015. Effect of Integrated Nutrient Management on Growth and Productivity of Hybrid Rice. *Journal of Agricultural Science and Technology*, 297-308.
- Geetha, B., Balanagoudar, S. R., Gaddi, A. K., Manjunatha, B., & Ramesh, Y. M. 2020. Effect of integrated plant nutrient supply system on growth, yield and economics of dry-direct seeded rice. *International Journal of Communication Systems*, 8(3): 2241-2244.
- Rao, A., Singh, N. B., Pandey, D., & Singh, M. P. 2020. Effect of integrated nutrient management on productivity and profitability of rice under SRI. *Journal of Pharmacognosy and Phytochemistry*, 9(5): 665-667.
- Maskina, M. S., Yadvinder Singh and Bijay Singh. 1987. Wheat straw management for rice on a coarse textured soil. *International Rice Research*, 12 (2): 40.
- Budhar, M. N., Palaniappan, S. P and Rangaswamy, A. 1991. Effect of farm wastes and green manures on low land rice. *Indian Journal of Agronomy*, 36 (2): 251-252.
- Shashi Kumar, G. S., Reddy, S. N and Mohd. Ikramullah. 1995. Effect of age of seedling and N level on performance of rice under late planting. *Indian Journal of Agricultural Sciences*, 65 (5): 354-355.

Sarawgi, S.K. and Sarawgi, A.K. 2004. Effect of blending of N with or without FYM on semi-dwarf, medium to long slender scented rice varieties in lowland alfisols of Chhattisgarh. In: International symposium on Rainfed Rice Ecosystems: Perspective and Potential. IGKV, Raipur, India. 11-13th October, 2004: 159-160.

V. Satyanarayana , P. V. Vara Prasad , V. R.K. Murthy & K. J. Boote. 2002. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice, *Journal of Plant Nutrition*, 25:10, 2081-2090.

M. Umar Khan, M. Qasim and Israr Ullah Khan. 2007. Effect of integrated nutrient management on crop yields in rice-wheat cropping system. *Sarhad Journal of Agriculture*, 23, (4).

Arif, M., Tasneem, M., Bashir, F., Yaseen, G. and Iqbal, R.M. 2014. Effect of integrated use of organic manures and inorganic fertilizers on yield and yield components of rice. *Journal of Agricultural Research*, 52(2): 197-206.

Priyanka, G., Sharma, G. D., Rachana, R., & Lal, B. 2013. Effect of integrated nutrient management and spacing on growth parameters, nutrient content and productivity of rice under system of rice intensification. *International Journal of Biological Sciences*, 2: 53-59.

Wahlang, B., Das, A., Layek, J., Munda, G.C., Ramkrushna, G.I. and Panwar, A.S. 2015. Effect of establishment methods and nutrient management on physiological attributes and water-use efficiency of rice. (*Oryza sativa*) in a sub-tropical climate. *Indian Journal of Agronomy*, 60(4): 157-163.

Shalini, Singh, V.P. and Jangid, B. 2017. Yield and economics in direct seeded rice using organic manures and micronutrients. *International Journal of Chemical Studies*, 5(3): 105-109.

Siddaram., Reddy, V.C. and Yogananda, S.B. 2017. Effect of Farmyard manure and Bio-digester liquid manure on Growth and Yield of Aerobic rice (*Oryza sativa* L.). *International Journal of Pure and Applied Bioscience*, 5(1): 832-839.

Adhikari, J., Sarkar, M. A. R., Uddin, M. R., Sarker, U.K., Hossen, K. and Rosemila, U. 2018. Effect of nitrogen fertilizer and weed management on the yield of transplanted aman rice. *Journal of Bangladesh Agricultural University*, 16(1): 12-16.

Kipgen, N., Irungbam, P., Pal, S., Gogoi, M. and Sanatombi, Y. 2018. Effect of varying levels of nitrogen and planting geometry on high yielding boro rice in new alluvial zone of west bengal. *International Journal of Current Microbiology and Applied Sciences*, 7(7): 2090-2098.

Harikesh., Ali, A., Shivam., ., Yadav, R.P., Kumar, S., Kumar, A. and Yadav, A. 2017. Effect of integrated nutrient management and plant geometry on growth and quality of rice (*Oryza sativa* L.) varieties under SRI technique. *Indian Journal of Current Microbiology and Applied Sciences*, 6(10): 2503-2515.

Tomar, R., Singh, N.B., Singh, V. and Kumar, D. 2018. Effect of planting methods and integrated nutrient management on growth parameters, yield and economics of rice. *Journal of Pharmacognosy and Phytochemistry*, 7(2): 520- 527.

Singh, S., Bohra, J.S., Singh, Y.V., Upadhyay, A.K., Verma, S.S., Mishra, P.K. and Raghuvver, M. 2017. Effect of integrated nutrient management on growth and development stages of rice under rice-wheat ecosystem. *Indian Journal of Current Microbiology and Applied Sciences*, 6(7): 2032-2042.

Siavoshi, M., Nasiri, A., & Laware, S. L. 2011. Effect of organic fertilizer on growth and yield components in rice (*Oryza sativa* L.). *Journal of Agricultural science*, 3(3): 217.

Hasanuzzaman, M., Ahamed, K.U., Rahmatullah, N.M., Akhter, N., Nahar, K. and Rahman, M.L. 2010. Plant growth characters and productivity of wetland rice (*Oryza sativa* L.) as affected by application of different manures. *Emirates Journal of Food and Agriculture*, 22(1): 46-58.

Gohain, T., 2014. Performance of local rice cultivars under aerobic ecosystem of Nagaland. *Annals of Plant and Soil Research*, 16 (4): 342-345.

Choudhary, A. K., R. C. Thakur, and N. Kumar. 2007. Effect of organic manures and chemical fertilizers on productivity and profitability in wheat–rice crop sequence. *Oryza*, 44 (3): 239–242.

- Hussain A, Sheraz S, Bhat RA, Rasool FA and Raihana HK. 2012. Integrated nutrient management of Rice (*Oryza sativa* L.) under temperate conditions of Kashmir. *Agricultural Science Digest*, 32(1): 18-22.
- Aulakh, C.S., Kaur, P., Walia, S.S., Gill, R.S., Sharma, S. and Buttar, G.S. 2016. Productivity and quality of basmati rice (*Oryza sativa*) in relation to nitrogen management. *Indian Journal of Agronomy*, 61(4): 467-473.
- Apon, M., Gohain, T., Apon, R., Banik, M., & Mandal, A. K. (2018). Effect of integrated nutrient management on growth and yield of local rice (*Oryza sativa* L.) under rainfed upland condition of Nagaland. *The Pharma Innovation J*, 7: 426-29.
- V. Satyanarayana , P. V. Vara Prasad , V. R.K. Murthy & K. J. Boote. 2002. Influence of integrated use of farmyard manure and inorganic fertilizers on yield and yield components of irrigated lowland rice. *Journal of Plant Nutrition*, 25:10, 2081-2090.
- Naing Oo A, Banterng P, Polthane A, Trelo-Ges V. 2010. The effect of different fertilizers management strategies on growth and yield of upland black glutinous rice and soil property. *Asian Journal of Plant Science*, 9: 414-22.
- Acharya CL, Bisnoi SK and Yaduvanshi H S.1998. Effect of long term application of fertilizer and amendment under continuous cropping and soil physical properties in Alfisols. *Indian Journal of Agricultural Science*, 58: 509-516.
- Singh AK, Kumar A and Mishra SS. 1998. Performance of rice varieties at different nitrogen level under irrigation and ecosystem of north Bihar. *Indian Journal of Agronomy*, 43 (2): 273-277.
- Mann, R. A., Ashraf, M. 2000. Reduction of chemical fertilizers through organic matter supplement for rice production. *Pakistan Journal of Agricultural Research*, 16(1): pp.20-23.
- Jayabala A and Kuppaswamy G. 2001. Recycling of organic wastes for the production of vermicompost and its response in rice – legume cropping system and soil fertility. *European Journal of Agronomy*, 15(3): 153-170.
- Kumar HMP, Meli SS and Angadi VV. 2002. Response of scented rice to integrated nutrient management under upland sown drill condition. *Research on Crops*, 3(3): 481-487.
- K. Barik, A. Raj and R. K. Saha, "Yield Performance, Economics and Soil Fertility through Organic Sources (Vermi-Compost) of Nitrogen as Substitute to Chemical Fertilizers in Wet Season Rice," *Crop Research*, 36(1) pp. 4-7.
- Raul PK and Sarawagi SK. 2005. Effect of integrated nitrogen nutrition techniques on yield, nitrogen content, uptake and efficiency of rice. *Indian Journal of Agronomy*, 50(2): 129-131.
- Choudhary, A. K., R. C. Thakur, and N. Kumar. 2007. Effect of organic manures and chemical fertilizers on productivity and profitability in wheat–rice crop sequence. *Oryza*, 44 (3): 239–242.
- Hossain MA, Shamsuddoha ATM, Paul AK, Bhuiyan MSI and Zobaer ASM. 2011. Efficacy of different organic manures and inorganic fertilizer on the yield and yield attributes of boro rice. *The Agriculturists*, 9(1-2): 117-125.
- Bora D, Goswami J, kumar R and Begum M. 2013. Effect of organic inputs on production and quality of scented rice variety keteki joha in Assam and its economic aspect. *Agriculture Science Digest*, 34 (2): 115-118.
- Ahmad S, Hussain A, Ali H and Ahmad A. 2005. Grain yield of transplanted rice (*Oryza sativa* L.) as influenced by plant density and nitrogen fertilization. *Journal of Agriculture and Social Sciences*, 1: 212-215.
- Kumar S, Saravanan S, Ramesh K, Natarajan SK and Mani S. 2005. Organic farming: Impact on rice productivity and soil health. *Asian Journal of Plant Sciences*, 4(5): 510-512.
- Gupta V, Sharma RS and Vishvakarma SK. 2006. Long-term effect of integrated nutrient management on yield sustainability and soil fertility of rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system. *Indian Journal of Agronomy*, 51: 160-164.
- Mankotia BS. 2007. Effect of fertilizer application with farm yard manure in standing rice (*Oryza sativa*) wheat (*Triticum aestivum*) cropping system. *Indian Journal of Agricultural Science*, 77(8): 39–43.

Pattanayak, S.K., Rao, D.L.N. and Mishra, K.N. 2007. Effect of biofertilizers on yield, nutrient uptake and nitrogen economy of rice-peanut cropping sequence. *Journal of the Indian Society of Soil Science*, 55(2): 184-189.

Virdia, H.M. and Mehta, H.D. 2010. Integrated nutrient management in transplanted rice (*Oryza sativa* L.). *International Journal of Agricultural Sciences*, 6(1): 295-299.

Mehdi, S.M., Sarfraz, M., Abbas, S.T., Shabbir, G. and Akhtar, J. 2011. Integrated nutrient management for rice-wheat cropping system in a recently reclaimed soil. *Soil Environment*, 30(1): 36-44.

Balasubramanian, A. and Wahab, K. 2012. Integrated nutrient management in rice for Cauvery Deltaic zone of Tamil nadu, India. *Plant Archives*, 12(1): 95-97.

Singh, G., Singh, S. and Singh, R.K. 2012. Effect of fertility management on yield and economics of traditional scented rice varieties in lowlands. *Annals Plant Soil Research*, 14(1): 1-4.

Sharma U and Subehia SK. 2014. Effect of Long-Term Integrated Nutrient Management on Rice (*Oryza sativa* L.) - Wheat (*Triticum aestivum* L.) Productivity and Soil Properties in North-Western Himalaya. *Journal of the Indian Society of Soil Science*, 62(3): 248-254.

Murali, M.K. And Setty, R.A., 2001, Growth, yield and nutrient uptake of scented rice (*Oryza sativa* L.) as influenced by level of NPK, vermicompost and triacontanol. *Mysore Journal of Agricultural Sciences*, 35(1): 1-4.

Mhaskar NV and Thorat ST. 2005. Effect of nitrogen levels on NPK uptake and grain yield of scented rice varieties under Konkan condition. *Journal of Soils and Crops*, 15(1): 206-209.

Kumar, V., Singh, O.P. and Kumar, V. 2006. Integrated nutrient management in rice (*Oryza sativa*)- wheat (*Triticum aestivum*) cropping system. *Bhartiya vaigyanikevam audyogik anusandhan patrika*, 15(1): 34-43.

Banik, P., and R. C. Sharma. 2008. Effects of integrated nutrient management with mulching on rice-based system in rainfed eastern plateau area. *Indian Journal of Agricultural Sciences*, 78: 240-243.

Choudhary, A. K., and V. K. Suri. 2009. Effect of organic manures and inorganic fertilizers on productivity, nutrient uptake, and soil fertility in wheat (*Triticum aestivum*)-paddy (*Oryza sativa*) crop sequence in western Himalayas. *Current Advances in Agricultural Sciences*, 1(2): 65-69.

Talathi MS, Pinjari SS, Ranshur NJ, Bhondave TS, Suryawansi JS. 2009. Response of hybrid rice to green leaf manure, FYM and Chemical fertilizer. *International Journal of Agriculture Science*, 5(2): 501-509.

Kumar, J., Yadav, M. P. and Kedar, P. 2010. Production potential of hybrid rice (*Oryza sativa* L.) as influenced by integrated nutrient management. *Crop Research*, 39: 20-23.

Siddaram, K., Murali, M.K., Basavaraja, Krishna Murthy, N. and Manjunatha, B. N., 2010. Effect of nitrogen levels through organic sources on yield attributes, yield and economics of irrigated Aerobic rice. *Mysore Journal of Agricultural Sciences*, 44(3): 485-489.

Weijabhandara, D.M.D.I., Dasog, G.S., Patil, P.L. and Hebbar, M. 2011. Effect of nutrient levels on rice (*Oryza sativa* L) under system of rice intensification (SRI) and traditional methods of cultivation. *Journal of the Indian Society of Soil Science*, 59(1): 67-73.

Acharya, R., Dash, A.K and Senapati, H.K. 2012. Effect of integrated nutrient management on microbial activity influencing grain yield under rice-rice cropping system in an acid soil. *Asian Journal of Microbiology, Biotechnology & Environmental Sciences Paper*, 14(3): 365-368.

Manjunatha B, Hanumanthappa M, Nahesha L, Kalyanamurthy KN, Kamath SKV. 2012. Effect of new herbicides on nutrient uptake in transplanted rice in Western Karnataka. *Journal of Agricultural Science*, 46(4): 928-930.

Ranjitha, P.S. and Reddy, K.I. 2013. Effect of different nutrient management options on rice under SRI method of cultivation-a review. *International Journal of Plant, Animal and Environmental Sciences*, 4(1): 201-204.

Tomar, R., Singh, N. B., Singh, V., & Kumar, D. 2018. Effect of planting methods and integrated nutrient management on growth parameters, yield and economics of rice. *Journal of Pharmacognosy Phytochemistry*, 7(2): 520-527.

Vennila C and Jayanthi C. 2007. Growth and yield of *coleus forskohlii* as influenced by source of nitrogen. Tamil Nadu Agri. Univ. Coimbatore.

Dass, A., Sudhishri, S., & Lenka, N. K. 2009. Integrated nutrient management for upland rice in Eastern Ghats of Orissa. *Oryza*, 46(3): 220-226.

Rao NSS 2007. Soil Microbiology. Oxford and IBH Publishing Co. Pvt. Ltd. 405p

Ghosh S, Wilson B, Ghoshal S, Senapati N and Mandal B. 2012. Organic amendments influence on soil quality and carbon sequestration in the indo- Gangetic plains of India. *Agriculture, Ecosystem and Environment*, 156(3): 134-141.

