

- Clubfoot in Uganda*. (August), 341–344.
- Ministry of Finance and Economic Planning Rwanda. (2019). *Sustainable Development Goals .2019 Rwanda Voluntary National Review (VNR) Report*.
- MINISTRY OF HEALTH FOURTH HEALTH SECTOR July 2018 – June 2024**. (2018). (July).
- Mkandawire, N. ., & Kaunda, E. (2004). Incidence and Patterns of Congenital Talipes Equinovarus (Clubfoot) Deformity At Queen Elizabeth Central Hospital, Banter, Malawi. *East and Central African Journal of Surgery*, 9(2), 28–31.
- Munambah, N., Chiwaridzo, M., & Mapingure, T. (2016). A cross-sectional study investigating impressions and opinions of medical rehabilitation professionals on the effectiveness of the Ponseti method for treatment of clubfoot in Harare, Zimbabwe. *Archives of Physiotherapy*, 6(1), 1–9.
<https://doi.org/10.1186/s40945-016-0021-5>
- Naddumba, E. . (2010). Preventing neglected club feet in Uganda: A challenge to the health workers with limited resources. *East African Orthopaedic Journal*, 3(1), 23–28.
<https://doi.org/10.4314/eaoj.v3i1.62560>
- Owen, R. M., & Kembhavi, G. (2012). A critical review of interventions for clubfoot in low and middle-income countries : effectiveness and contextual influences. *Journal of Pediatric Orthopaedics B*, 21, 59–67.
<https://doi.org/10.1097/BPB.0b013e3283499264>
- Paton, R. W., Fox, A. E., Foster, A., & Fehily, M. (2010). Incidence and aetiology of talipes equino-varus with recent population changes. *Acta Orthopaedica Belgica*, 76(1), 86–89.
- Rasheed, N., Zaidi, I. H., Rasheed, N., & Hussain, G. (2017). Awareness Regarding Clubfoot in Parents. *JPOA*, 90–94.
- Rwanda Clubfoot Program. (2019). *Annual report*.
- Wallander, H., Hovelius, L., & Michaelsson, K. (2006). Incidence of congenital clubfoot in Sweden. *Acta Orthopaedica*, 77(6), 847–852.
<https://doi.org/10.1080/17453670610013123>