

2 PROBLEM STATEMENT

One in every 59 child has been diagnosed with ASD. Fever is the main presenting symptoms for these infections, and such fever during pregnancy is supposedly linked to ASD, are infections during pregnancy linked to ASD?

3 OBJECTIVES

3.1 General Objective

To study maternal infection and other characteristics during pregnancy and its link to autism spectrum disorders in Khartoum locality, Sudan.

Specifically we wanted to assess the link between history of gestational infection and ASD, and to study some of the maternal characteristics.

4 METHODS

4.1 Ethics

Ethical clearance was obtained from the Ministry of Health Research Ethical committee.

The protocol, aims and benefits of the study were explained for all the participants in the study, and a written voluntary informed consent was obtained from each.

4.2 Population

The study enrolled parents and guardians of children who were 5 years or younger and suffering from ASD. According to Intitute administration there were (180) Child with ASD enrolled at the time of the study. Using the modified Cochran Formula for small populations sample calculation: $n = n_0 / 1 + ((n_0 - 1) / N)$. Where $n_0 =$ Recommended Sample size (384)* and $N =$ Population. $n_0 = 384 / (1 + (384 / 180)) = 123$. The sampling procedure was systematic sampling, with sampling interval of 3. Where a participant was selected after every 2 names were encountered

4.3 Procedure and Analysis

The data was collected during the study period between March and July 2019 from the institution by the researcher, the data collection. Data was collected via an interview by the researcher with a semi-structured questionnaire that was based on previous studies then was modified to meet the objectives of this study.

The analysis was carried out using the Statistical Package for Social Sciences (SPSS) version 25.0, SPSS Inc. Chicago, IL. The frequency distributions for independent variable and dependent variable was generated.

5 RESULTS

The study included 123 child diagnosed with ASD, 31.7% were males and 68.3% were females, Of all the participants' mothers 64.9% had a significant maternal condition (Table 1) 43.0% had diabetes, 13.0% had Placenta praevia (PP), 6.5% had Asthma, 2.4% had Pregnancy hypertension, and 2.4% had

more than one condition.

TABLE 1
MATERNAL CONDITIONS

Maternal conditions	F	%
Maternal diabetes	53	43.0
Pregnancy hypertension	3	2.4
Asthma	8	6.5
Placenta praevia	16	13.0
Placenta Abruptio	0	0
More than one condition	3	2.4
Total	80	67.3

Most of the participants mothers did not have an infection during pregnancy, neither Bacterial 82.9% nor Viral 95.1%, 12.2% had a bacterial infection the third trimester and 4.9% had one in the second trimester, the viral infection rate in each of the first and second trimesters was 2.4% (Figure 1).





FIGURE 1
INFECTIONS IN PREGNANCY

As for the Hospitalization and antibiotic use (Figure 2); 73.2% of the participants' mothers was not hospitalized during pregnancy, 12.2% reported hospitalization during second trimester, 7.3% reported for both first and third trimester. Antibiotic during pregnancy was used by 17.1% during second trimester, 9.8% in third trimester and 7.3% in first trimester, and was not used by 65.9% at all during pregnancy (Figure 2).



As for the condition of delivery (Table 2); elective Caesarean section (CS) compromised 41.4% of all labour and delivery conditions, followed by Preterm delivery at 21.1%, then emergency CS 18.6%, and 4.8% had Forceps use to assist in delivery.

TABLE 2
DELIVERY CONDITIONS

Labour and Delivery	F	%
Preterm	26	21.1
PROM/Spontaneous	6	4.8
Forceps use	6	4.8
Vacuum use	0	0
Elective CS	51	41.4
Emergency CS	23	18.6
Breech	0	0
Total	112	90.7%

6 DISCUSSION

The aim of this study was to describe some of the maternal infection and other characteristics during pregnancy and labour. The aim of this study was to describe some of the maternal infection and other characteristics during pregnancy and labour, maternal diabetes was most abundant, in the study participants, almost half of the study participants mothers suffered from diabetes, and although the degree of control during pregnancy was not explored in this study, studies suggested the connection between maternal diabetes and ASD [46]. It was also suggested that exposure to maternal GDM was associated with risk of ASD for the child [47].

Preeclampsia has been examined as a risk factor for ASD in multiple investigations with mixed results for example a meta-analysis [48] of 17 studies suggested substantial unexplained het-

erogeneity of effect estimates and also population-based, case-control studies [49],[50],[51] reported statistically significant increased adjusted odds of ASD after pregnancies complicated by preeclampsia. However in this study the percentage of maternal hypertension was low which is more likely due to coincidence. Findings also suggest that other maternal conditions in pregnancy such as asthma is not likely associated with increased risk of ASD agreeing with another study [52].

In this study infections; both bacterial and viral were not prevalent in pregnancy nor was hospital admission, in a previous study also no overall association between diagnoses of any maternal infection during pregnancy whether viral or bacterial was associated with ASD. However, women with infections diagnosed during a hospital admission, mainly bacterial infections were at increased risk of delivering a child with ASD. And multiple infections during pregnancy were associated with ASD [53], our results do not suggest that use of antibiotics during pregnancy is strong risk factors for ASD similar to another study [54].

Delivery conditions were not momentous except for CS, which risk is agreed upon with previous findings that stated children born by CS (elective or emergency) are approximately 20% more likely to be diagnosed as having ASD, considering the reason behind choosing this mode of delivery is probably the actual risk factor [55],[56].

7 CONCLUSION

Our results suggests that DM and CS are risk factors for ASD, also it suggests that bacterial infections, viral infections, or use of antibiotics during pregnancy are strong risk factors for ASD. The few positive findings are possible coincidence findings.

8 RECOMMENDATIONS

The recommendations from this study is to carry out educational campaigns for the community about the risk of developing ASD. And to conduct further research exploring these factors in a larger population in various regions of the country, and comparing the risk factors in pregnant ladies who give birth to healthy children.

END SECTIONS

ACKNOWLEDGMENT

The authors wish to thank A, B, C. This work was supported in part by a grant from XYZ.

REFERENCES

- [1] American Psychiatric Association. Diagnostic Criteria from dsm-iv-tr. American Psychiatric Pub; 2000.
- [2] Baio J, Wiggins L, Christensen DL, Maenner MJ, Daniels J, Warren Z, Kurzius-Spencer M, Zahorodny W, Rosenberg CR, White T, Durkin MS. Prevalence of autism spectrum disorder among children aged 8 years—autism and developmental disabilities monitoring network, 11 sites, United States, 2014. *MMWR Surveillance Summaries*. 2018 Apr 27;67(6):1.
- [3] Gillberg C, Wing L. Autism: not an extremely rare disorder. *Acta Psychiatrica Scandinavica*. 1999 Jun;99(6):399-406.
- [4] American Psychiatric Association. Diagnostic and statistical manual of mental disorders (DSM-5®). American Psychiatric Pub; 2013 May 22.
- [5] Abrahams BS, Geschwind DH. Advances in autism genetics: on the threshold of a new neurobiology. *Nature reviews genetics*. 2008 May;9(5):341.
- [6] De Rubeis S, He X, Goldberg AP, Poultnery CS, Samocha K, Cicek AE et al. Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature* 2014; 515: 209-215.
- [7] Geschwind DH, State MW. Gene hunting in autism spectrum disorder: on the path to precision medicine. *Lancet Neurol* 2015; 14: 1109-1120.
- [8] Iossifov I, O'Roak BJ, Sanders SJ, Ronemus M, Krumm N, Levy D et al. The contribution of de novo coding mutations to autism spectrum disorder. *Nature* 2014; 515: 216-221.
- [9] Krumm N, Turner TN, Baker C, Vives L, Mohajeri K, Witherspoon K et al. Excess of rare, inherited truncating mutations in autism. *Nat Genet* 2015; 47: 582-588.
- [10] Casanova EL, Sharp JL, Chakraborty H, Sumi NS, Casanova MF. Genes with high penetrance for syndromic and non-syndromic autism typically function within the nucleus and regulate gene expression. *Mol Autism* 2016; 7: 18.
- [11] Sanders SJ, Murtha MT, Gupta AR, Murdoch JD, Raubeson MJ, Willsey AJ et al. De novo mutations revealed by whole-exome sequencing are strongly associated with autism. *Nature* 2012; 485: 237-241.
- [12] Atladottir HO, Thorsen P, Ostergaard L, Schendel DE, Lemcke S, Abdallah M et al. Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord* 2010; 40: 1423-1430.
- [13] Zerbo O, Qian Y, Yoshida C, Grether JK, Van de Water J, Croen LA. Maternal infection during pregnancy and autism spectrum disorders. *J Autism Dev Disord* 2015; 45: 4015-4025.
- [14] Lee BK, Magnusson C, Gardner RM, Blomstrom A, Newschaffer CJ, Burstyn I et al. Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun* 2015; 44: 100-105.
- [15] Fang SY, Wang S, Huang N, Yeh HH, Chen CY. Prenatal infection and autism spectrum disorders in childhood: a population-based case-control study in Taiwan. *Paediatr Perinat Epidemiol* 2015; 29: 307-316.
- [16] Patterson PH. Maternal infection and immune involvement in autism. *Trends Mol Med* 2011; 17: 389-394.
- [17] Abdallah MW, Larsen N, Grove J, Norgaard-Pedersen B, Thorsen P, Mortensen EL et al. Amniotic fluid inflammatory cytokines: potential markers of immunologic dysfunction in autism spectrum disorders. *World J Biol Psychiatry* 2013; 14: 528-538.
- [18] Goines PE, Croen LA, Braunschweig D, Yoshida CK, Grether J, Hansen R et al. Increased midgestational IFN-gamma, IL-4 and IL-5 in women bearing a child with autism: a case-control study. *Mol Autism* 2011; 2: 13.
- [19] Cordeiro CN, Tsimis M, Burd I. Infections and brain development. *Obstet Gynecol Surv* 2015; 70: 644-655.
- [20] Dreier JW, Andersen AM, Berg-Beckhoff G. Systematic review and meta-analyses: fever in pregnancy and health impacts in the offspring. *Pediatrics* 2014; 133: e674-e688.
- [21] Atladottir HO, Thorsen P, Ostergaard L, Schendel DE, Lemcke S, Abdallah M et al. Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord* 2010; 40: 1423-1430.
- [22] Jiang HY, Xu LL, Shao L, Xia RM, Yu ZH, Ling ZX et al. Maternal infection during pregnancy and risk of autism spectrum disorders: a systematic review and meta-analysis. *Brain Behav Immun* 2016; 58: 165-172.
- [23] Lee BK, Magnusson C, Gardner RM, Blomstrom A, Newschaffer CJ, Burstyn I et al. Maternal hospitalization with infection during pregnancy and risk of autism spectrum disorders. *Brain Behav Immun* 2015; 44: 100-105.
- [24] Baron-Cohen S, Auyeung B, Norgaard-Pedersen B, Hougaard DM, Abdallah MW, Melgaard L et al. Elevated fetal steroidogenic activity in autism. *Mol Psychiatry* 2015; 20: 369-376.
- [25] Barker DJ. In utero programming of chronic disease. *Clin Sci (Lond)* 1998; 95: 115-128.
- [26] Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med* 2008; 359: 61-73.
- [27] Brown AS, Sourander A, Hinkka-Yli-Salomaki S, McKeague IW, Sundvall J, Surcel HM. Elevated maternal C-reactive protein and autism in a national birth cohort. *Mol Psychiatry* 2014; 19: 259-264.
- [28] Chess S. Autism in children with congenital rubella. *Journal of autism and childhood schizophrania*. 1971 Jan 1;1(1):33-47.
- [29] DEYKIN EY, MACMAHON B. Viral exposure and autism. *American Journal of Epidemiology*. 1979 Jun 1;109(6):628-38.
- [30] Ghaziuddin M, Tsai LY, Eilers L, Ghaziuddin N. Brief report: autism and herpes simplex encephalitis. *Journal of autism and developmental disorders*. 1992 Mar 1;22(1):107-13.
- [31] Yamashita Y, Fujimoto C, Nakajima E, Isagai T, Matsuishi T. Possible association between congenital cytomegalovirus infection and autistic disorder. *Journal of autism and developmental disorders*. 2003 Aug 1;33(4):455-9.
- [32] Mason-Brothers A, Ritvo ER, Pingree C, Petersen PB, Jenson WR, McMahan WM, Freeman BJ, Jorde LB, Spencer MJ, Mo A, Ritvo A. The UCLA-University of Utah epidemiologic survey of autism: prenatal, perinatal, and postnatal factors. *Pediatrics*. 1990 Oct 1;86(4):514-9.
- [33] Zerbo O, Iosif AM, Walker C, Ozonoff S, Hansen RL, Hertz-Picciotto I. Is maternal influenza or fever during pregnancy associated with autism or developmental delays? Results from the CHARGE (Childhood Autism Risks from Genetics and Environment) study. *Journal of autism and developmental disorders*. 2013 Jan 1;43(1):25-33.
- [34] Gardener H, Spiegelman D, Buka SL (2009) Prenatal risk factors for autism: comprehensive meta-analysis. *The British Journal of Psychiatry* 195: 7-14.
- [35] Burstyn I, Sithole F, Zwaigenbaum L (2010) Autism spectrum disorders, maternal characteristics and obstetric complications among singletons born in Alberta, Canada. *Chronic Dis Can* 30: 125-134.
- [36] Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparen P, et al. (2009) Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics* 124: e817-825 Epub 2009 Oct 2019.
- [37] Mann JR, McDermott S, Bao H, Hardin J, Gregg A (2010) Pre-eclampsia, birth weight, and autism spectrum disorders. *J Autism Dev Disord* 40: 548-554.
- [38] Larsson HJ, Eaton WW, Madsen KM, Vestergaard M, Olesen AV, et al. (2005) Risk Factors for Autism: Perinatal Factors, Parental Psychiatric History, and Socioeconomic Status. *American Journal of Epidemiology* 161: 916-925.
- [39] Gillberg C, Gillberg IC (1983) Infantile autism: A total population study of reduced optimality in the pre-, peri-, and neonatal period. *Journal of Autism and Developmental Disorders* 13: 153-166.

- [40] Limperopoulos C, Bassan H, Sullivan NR, Soul JS, Robertson RL, et al. (2008) Positive Screening for Autism in Ex-preterm Infants: Prevalence and Risk Factors. *Pediatrics* 121: 758-765.
- [41] Williams K, Helmer M, Duncan GW, Peat JK, Mellis CM (2008) Perinatal and maternal risk factors for autism spectrum disorders in New South Wales, Australia. *Child: Care, Health and Development* 34: 249-256.
- [42] Juul-Dam N, Townsend J, Courchesne E (2001) Prenatal, Perinatal, and Neonatal Factors in Autism, Pervasive Developmental Disorder-Not Otherwise Specified, and the General Population. *Pediatrics* 107: e63
- [43] Croen LA, Grether JK, Selvin S (2002) Descriptive Epidemiology of Autism in a California Population: Who Is at Risk? *Journal of Autism and Developmental Disorders* 32: 217-224.
- [44] Hultman CM, Sparén P, Cnattingius S (2002) Perinatal Risk Factors for Infantile Autism. *Epidemiology* 13: 417-423.
- [45] Gardener H, Spiegelman D, Buka SL (2011) Perinatal and Neonatal Risk Factors for Autism: A Comprehensive Meta-Analysis. *Pediatrics*
- [46] Xu G, Jing J, Bowers K, Liu B, Bao W. Maternal diabetes and the risk of autism spectrum disorders in the offspring: a systematic review and meta-analysis. *Journal of autism and developmental disorders*. 2014 Apr 1;44(4):766-75.
- [47] Xiang AH, Wang X, Martinez MP, Walthall JC, Curry ES, Page K, Buchanan TA, Coleman KJ, Getahun D. Association of maternal diabetes with autism in offspring. *Jama*. 2015 Apr 14;313(14):1425-34.
- [48] Gardener H, Spiegelman D, Buka SL. Prenatal risk factors for autism: comprehensive meta-analysis. *Br J Psychiatry*. 2009;195(1):7-14.
- [49] Mann JR, McDermott S, Bao H, Hardin J, Gregg A. Pre-eclampsia, birth weight, and autism spectrum disorders. *J Autism Dev Disord*. 2010;40(5):548-554.
- [50] Dodds L, Fell DB, Shea S, Armson BA, Allen AC, Bryson S. The role of prenatal, obstetric and neonatal factors in the development of autism. *J Autism Dev Disord*. 2011;41(7):891-902.
- [51] Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparén P, Cnattingius S. Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics*. 2009;124(5):e817-e825.
- [52] Croen LA, Grether JK, Yoshida CK, Odouli R, Van de Water J. Maternal autoimmune diseases, asthma and allergies, and childhood autism spectrum disorders: a case-control study. *Archives of pediatrics & adolescent medicine*. 2005 Feb 1;159(2):151-7.
- [53] Zerbo O, Qian Y, Yoshida C, Grether JK, Van de Water J, Croen LA. Maternal infection during pregnancy and autism spectrum disorders. *Journal of autism and developmental disorders*. 2015 Dec 1;45(12):4015-25.
- [54] Atladóttir HÓ, Henriksen TB, Schendel DE, Parner ET. Autism after infection, febrile episodes, and antibiotic use during pregnancy: an exploratory study. *Pediatrics*. 2012 Dec 1;130(6):e1447-54.
- [55] Curran E, O'Neill S, Cryan J, et al. Birth by caesarean section and development of autism spectrum disorder and attention-deficit/hyperactivity disorder: a systematic review and meta-analysis. *J Child Psychol Psychiatry*. 2015;56(5):500-508.
- [56] Lavender T, Hofmeyr GJ, Neilson JP, Kingdon C, Gyte GML. Caesarean section for non-medical reasons at term. *Cochrane Database Syst Rev*. 2012;3:CD004660.