

Herbal medicine is now one of the most significant areas of folk medicine, and herbal medicines are becoming increasingly popular for a variety of ailments. The comparison of phytochemicals extracted from plants and their pharmacological effects is becoming increasingly popular (George & Joseph, 2009). The therapeutic characteristics of the plant have been assessed on the basis of current systematic developments in the world due to its low toxicity, beneficial pharmacological activities, and economic viability. Flavonoids, phenols, sterols, and alkaloids are just a few of the bioactive molecules found in plants that have antioxidant properties.

Free radicals

Free radicals are a necessary component of any biochemical process and are an important feature of aerobic life and metabolism. A free radical is an atom or molecule with a lone pair of electrons that is chemically reactive and capable of rapid chain reactions that destabilize other molecules and generate a large number of free radicals. Most radicals are highly reactive and unstable, allowing them to give or absorb an electron from a nearby molecule, allowing them to operate as oxidants or reductants. Free radicals have half-lives measured in milliseconds, microseconds, or nanoseconds in general (Cheeseman & Slater, 1993). They can be reactive sulfur species, reactive oxygen species, which are formed from oxygen, or reactive nitrogen species, which are derived from nitrogen. RO (alkoxyl), O₂ (superoxide), HO₂ (hydroperoxyl), HO (hydroxyl), ROO (peroxyl), NO₂ (nitrogen dioxide), N₂O₃ (dinitrogen trioxide) and NO (nitric oxide) are the most common nitrogen-derived free radicals (Devasagayam & Kesavan, 1996). The majority of diseases are caused by oxidative stress caused by free radicals, antioxidant deficiency or reactive sulfur species (Velavan et al., 2007).

Antioxidant

Antioxidant protective tools in the form of biological enzymatic and non-enzymatic functions have been discovered in the human body to neutralize the damaging consequence of free radicals, reactive sulfur species and other oxidants (Alam et al., 2013). Neural disorders, cardiovascular disease, tissue damage, cancer, atherosclerosis, central nervous system injury, inflammation, obesity, arthritis, and ageing are all linked to free radicals (Köse et al., 2015; Gülçin et al., 2010).

Antioxidants in plants have recently been shown to reduce the damage caused by free radicals by inhibiting electron transport (Liochev, 2013) and have pharmacological benefits, including apoptosis, gene expression anticarcinogenic, free radical scavenging, antiallergic, antimutagenic, antibacterial, ion transportation, and antidiabetic characteristics (Puertollano et al., 2011; Lü et al., 2009; Pinchuk et al., 2012). To counteract oxidative stress, plants and humans have complex systems of redundant antioxidants, such as glutathione. Physiological functioning may deteriorate when the antioxidant defense is weakened owing to a range of factors, resulting in many diseases and aging. To avoid the harm effected by oxidation, however, a supply of antioxidants and antioxidant-containing foods is required (Halliwell & Gutteridge, 2003).

Type of antioxidants:

Antioxidants can be categorized into three types:

1. Primary antioxidants: Antioxidants that participate in the chain breaking system and react with lipid radicals to transform them into safer particles. The majority of the antioxidants in

this group are phenolic in nature. Because these antioxidants are cofactors of antioxidant enzymes, their absence may have an impact on macromolecule metabolism, particularly glucose metabolism.

2. Secondary antioxidants: These antioxidants are phenolic chemicals that play a role in stopping chain reactions and binding free radicals. Butylated hydroxytoluene (BHT), metal chelating agent (EDTA), glutathione-s-transferase, dehydrogenase, propyl gallate (PG), ubiquinone, butylated hydroxyl anisole (BHA), nordihydro guaretic acid, glucose-6-phosphate and tertiary butyl hydroquinone (TBHQ) are all examples of this group.

3. Tertiary antioxidants: These antioxidants, such as methionine sulfoxide, reductase, and DNA repair enzymes, help to repair biomolecules that have been damaged by free radicals (Nagar et al., 2017).

Antioxidants can also be classified into two groups. These are

1. Enzymatic antioxidants: These are antioxidants that have a role in the body's defense against free radicals and reactive oxygen species, either directly or indirectly. Enzymatic antioxidants comprise glutathione, superoxide dismutase (SOD), peroxidase, catalase, and others (Mehta & Gowder, 2015).

2. Non-enzymatic antioxidants: Dietary sources of antioxidants contain carotenoids, polyphenols, organosulfural, vitamins, compounds, and minerals. The largest class of antioxidants is polyphenols, which includes flavonoids, organosulfural, and phenolic acids (Bunaciu et al., 2016).

Antioxidants are divided into three categories based on their line of defense mechanism:

a. Minerals like selenium, copper, and zinc, as well as superoxide catalase (CAT), dismutase (SOD), and glutathione reductase (GR), may be found in first-line antioxidants.

b. Second-line antioxidants include albumin, glutathione (GSH), flavonoids, vitamin C, carotenoids, vitamin E and others.

c. Damaged DNA rehabilitates enzymes such as transferases, reductases, methionine sulphoxide and others, as well as deteriorated proteins, peroxides, and oxidized lipids, which are all part of the third line of antioxidants (Irshad & Chaudhuri, 2002).

Any antioxidant preparation used in food must be safe, be heat stable, simple to include, productive at intense concentrations, have no flavor, unpleasant odor, or color, and provide a financial advantage.

Means of antioxidants activity

Antioxidants are hypothesized to work in two different ways. To begin with, in a chain-breaking mechanism, the main antioxidant transfers one electron to the system's free radical. The process also involves the use of a quenching chain-initiating catalyst to eliminate reactive nitrogen species initiators (secondary antioxidants)/ROS. Other antioxidant methods, such as co-antioxidants, electron donation, metal ion chelation, or gene expression modulation, are effective in biological systems (Lobo et al., 2010).

Function of Antioxidants in human body

To protect the body's cells and organs from free radicals (ROS), a complex system has evolved in the human body, involving a mixture of endogenous and exogenous components that interact and synergize to minimize the effects of free radicals. These are the ones.

1. Vitamin E, carotenoids, vitamin C, vitamin B and other low molecular weight compounds including lipid acid and glutathione are all sources of antioxidants.
2. Glutathione peroxidase, glutathione reductase and superoxide dismutase are enzyme antioxidants that catalyze free radical quenching processes.
3. Ferritin, lactoferrin, ceruloplasmin and albumin are proteins that bind metals and capture free iron and copper ions, allowing them to catalyze oxidative processes.
4. A number of other antioxidant phytonutrients exist in a large diversity of plant foods (Sunil, 2014).

Antioxidants from dietary food

The consumption of fruits and vegetables on a regular basis has been shown to reduce the risk of chronic illnesses. People obtain antioxidant supplements directly from fresh vegetables and fruits, which contain a high concentration of antioxidant supplements and flavonoids that can aid in the defense mechanisms against a variety of cardiovascular disorders, as well as malignancies and other health problems (Hamid et al., 2010).

Studies show that a diet high in antioxidants has a large-scale positive impact on health. Antioxidants have been found in a wide range of plants and plant parts, including spinach, strawberries, grapes, plums, blueberries, broccoli blossoms, alfalfa sprouts, and many more, and they are also found in many dietary ingredients.

Citrus fruits, such as lemons and oranges, are also strong in natural antioxidants, particularly vitamin C. (Cao et al., 1998).

In addition, certain new and distinct antioxidants, such as flavonoid derivatives and p-coumaric acid, have been identified in spinach. The natural antioxidant p-coumaric acid and flavonoid derivatives derived from spinach have been shown to have a biological effect on the prevention of prostate cancer. Fruits like araticudomato, jack fruit and pindo palm have recently been revised as good sources of vitamin A, vitamin C, vitamin E, hydroxycinnamic acid and hydroxybenzoic acid. Research is being conducted on these fruits to create biological, chemical, or genetic variations in order to improve the antioxidant ability of the similar (Shebis et al., 2013).

Antioxidant from plants

Antioxidants, both natural and synthetic, are frequently utilized in foods and medicine, mostly those comprising fats and oils, to preserve the food against deterioration. Butylated hydroxyanisole (BHA), natural and synthetic food antioxidants, and butylated hydroxytoluene (BHT) have been mostly utilized in the therapeutic cosmetic and food industries. Nevertheless, customers' preferences led to a shift in producers' or manufacturers' selection of natural antioxidants over man-made antioxidants due to their instability at high temperatures, high volatility, and carcinogenic behavior (Papas, 1999).

In light of the rising risk of humans contracting numerous terminal diseases, there has been a worldwide direction toward using natural compounds found in dietary and medicinal

plants as therapeutic antioxidants. Antioxidant activity has been observed in a range of therapeutic herbs, including *Allium sativum*, according to Zingiber authorities. Numerous antioxidants, also known as active oxygen scavengers or free radicals, that are obtained naturally from plant sources and are used for food, cosmetics, and remedial purposes, have proven to be excellent alternatives to man-made antioxidants due to their absence of toxic effects on the human body and are inexpensive. Plants have a well-developed enzymatic and non-enzymatic scavenging strategy to combat the negative impacts of reactive oxygen species. Glutathione reductase, Catalase, superoxide dismutase, carotenoids, peroxidases, dehydroascorbate reductase, glutathione peroxidase. Glutathione, tocopherols, and ascorbate peroxidase are non-enzymatic molecules. In distinct organelles of plant cells, there are unmistakable, well-synchronized scavenging systems and ROS production. Lesser degrees of ROS comparatively act as signalling essences that arouse abiotic stress tolerance by altering the manifestation of resistant genes. In plants, elevated degrees of antioxidants have been accounted for to demonstrate better resistance to various types of environmental stresses (Hossain et al., 2012). Lower amounts of ROS, on the other hand, operate as signaling essences, modifying the expression of resistance genes to elicit abiotic stress tolerance. Increased antioxidant levels in plants have been linked to increased resilience to several forms of environmental stressors (Hasanuzzaman et al., 2012).

conclusion

The essential means of underlying a number of human neurological illnesses, autoimmune pathologies, inflammation, viral infections, digestive system and diabetes problems appears to be cellular damage caused by reactive nitrogen species (RNA), reactive oxygen species (ROS) and reactive sulfur species (RSS) production, or free radicals. Reactive sulfur species (RSS), Reactive oxygen species and free radicals are involved in various illnesses, according to a significant number of experimental studies. To combat these disorders, several man-made antioxidants are utilized in processed diets, but they have adverse effects. Several natural antioxidants may be useful in reducing the negative consequences of oxidative stress. As a result, estimating the antioxidant capacity of meals, botanicals, and other nutritional antioxidant supplements has received a lot of interest. Some medicinal plants, such as *Dodonaea viscosa*, *Barleria noctiflora*, and *Allium sativum*, have been shown to have high antioxidant capabilities because they comprise a huge amount of bioactive components such as carotenes, flavonoids carotenoids and phenolics. These chemicals have anti-diabetic, antibacterial, antifungal, anti-spasmodic, anticarcinogenic and anti-inflammatory, properties in addition to antioxidant action.

References

- A Puertollano, M., Puertollano, E., Alvarez de Cienfuegos, G., & A de Pablo, M. (2011). Dietary antioxidants: immunity and host defense. *Current topics in medicinal chemistry*, 11(14), 1752-1766.
- Akrout, A., Gonzalez, L. A., El Jani, H., & Madrid, P. C. (2011). Antioxidant and antitumor activities of *Artemisia campestris* and *Thymelaea hirsuta* from southern Tunisia. *Food and Chemical Toxicology*, 49(2), 342-347.
- Alam, M. N., Bristi, N. J., & Rafiquzzaman, M. (2013). Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*, 21(2), 143-152.

- Arumugam, S., Natesan, S., Ganesan, S., & Kanagarajan, S. (2015). In vitro screening of various extract of *Barleria noctiflora* for their antioxidant and free radical scavenging activity. *International Journal of Pharmaceutical and Phytopharmacological Research*, 5(2), 41-49.
- Benzie, I. F., & Wachtel-Galor, S. (Eds.). (2011). *Herbal medicine: biomolecular and clinical aspects*. CRC Press.
- Bunaciu, A. A., Danet, A. F., Fleschin, Ş., & Aboul-Enein, H. Y. (2016). Recent applications for in vitro antioxidant activity assay. *Critical Reviews in Analytical Chemistry*, 46(5), 389-399.
- Cao, G., Russell, R. M., Lischner, N., & Prior, R. L. (1998). Serum antioxidant capacity is increased by consumption of strawberries, spinach, red wine or vitamin C in elderly women. *The Journal of nutrition*, 128(12), 2383-2390.
- Chan, E. W. C., Tan, Y. P., Chin, S. J., Gan, L. Y., Kang, K. X., Fong, C. H., ... & How, Y. C. (2014). Antioxidant properties of selected fresh and processed herbs and vegetables. *Free Radicals and Antioxidants*, 4(1), 39.
- Cheeseman KH, Slater TF 1993. An introduction to free radicals chemistry. *Br Med Bull*.49:481–93. [PubMed]
- Devasagayam, T. P., & Kesavan, P. C. (1996). Radioprotective and antioxidant action of caffeine: mechanistic considerations. *Indian journal of experimental biology*, 34(4), 291-297.
- Dhiman, A., La, R., Bhan, M., Dhiman, B., & Hooda, A. (2012). Plebeian assessment of anti microbial and in vitro antioxidant zest of *D. fastuosa* L. seeds. *Journal of Pharmaceutical and Scientific Innovation*, 1(4), 49-53.
- Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P., & Vidal, N. (2006). Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food chemistry*, 97(4), 654-660.
- Ekor, M. (2013). The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in pharmacology*, 4.
- Elosta, A., Slevin, M., Rahman, K., & Ahmed, N. (2017). Aged garlic has more potent antiglycation and antioxidant properties compared to fresh garlic extract in vitro. *Scientific reports*, 7, 39613.
- Ganie, S. A., Dar, T. A., Hamid, R., Zargar, O., Abeer, S. U., Masood, A., ... & Zargar, M. A. (2014). In vitro antioxidant and cytotoxic activities of *Arnebia benthamii* (Wall ex. G. Don): a critically endangered medicinal plant of Kashmir valley. *Oxidative medicine and cellular longevity*, 2014.
- George, M., & Joseph, L. (2009). Anti-allergic, anti-pruritic, and anti-inflammatory activities of *Centella asiatica* extracts. *African Journal of Traditional, Complementary and Alternative Medicines*, 6(4), 554-559.

- Goveas, S. W., & Abraham, A. S. H. A. (2013). Evaluation of antimicrobial and antioxidant activity of stem and leaf extracts of *Coscinium fenestratum*. *Asian Journal of Pharmaceutical and Clinical Research*, 6(3), 218-221.
- Grace-Lynn, C., Darah, I., Chen, Y., Latha, L. Y., Jothy, S. L., & Sasidharan, S. (2012). In vitro antioxidant activity potential of lantadene A, a pentacyclic triterpenoid of *Lantana* plants. *Molecules*, 17(9), 11185-11198.
- Gülçin, I., Bursal, E., Şehitoğlu, M. H., Bilsel, M., & Gören, A. C. (2010). Polyphenol contents and antioxidant activity of lyophilized aqueous extract of propolis from Erzurum, Turkey. *Food and Chemical Toxicology*, 48(8), 2227-2238.
- Gupta, M., Mazumder, U. K., Kumar, R. S., Sivakumar, T., & Vamsi, M. L. M. (2004). Antitumor activity and antioxidant status of *Caesalpinia bonducella* against Ehrlich ascites carcinoma in Swiss albino mice. *Journal of Pharmacological Sciences*, 94(2), 177-184.
- Halliwell, B., & Gutteridge, J. M. (2015). *Free radicals in biology and medicine*. Oxford University Press, USA.
- Hamid, A. A., Aiyelaagbe, O. O., Usman, L. A., Ameen, O. M., & Lawal, A. (2010). Antioxidants: Its medicinal and pharmacological applications. *African Journal of pure and applied Chemistry*, 4(8), 142-151.
- Hasanuzzaman, M., Hossain, M. A., da Silva, J. A. T., & Fujita, M. (2012). Plant response and tolerance to abiotic oxidative stress: antioxidant defense is a key factor. In *Crop stress and its management: Perspectives and strategies* (pp. 261-315). Springer Netherlands.
- Irshad, M., & Chaudhuri, P. S. (2002). Oxidant-antioxidant system: role and significance in human body. *Indian J Exp Biol*, ; 40:1233- 1239.
- Jain, S., Jain, D. K., & Balekar, N. (2012). In-Vivo Antioxidant activity of ethanolic extract of *Mentha pulegium* leaf against CCl₄ induced toxicity in rats. *Asian Pacific Journal of Tropical Biomedicine*, 2(2), S737-S740.
- Kakouri, E., Daferera, D., Paramithiotis, S., Astraka, K., Drosinos, E. H., & Polissiou, M. G. (2017). *Crocus sativus* L. tepals: The natural source of antioxidant and antimicrobial factors. *Journal of Applied Research on Medicinal and Aromatic Plants*, 4, 66-74.
- Karagöz, A., Artun, F. T., Özcan, G., Melikoğlu, G., Anıl, S., Kültür, Ş., & Sütlüpinar, N. (2015). In vitro evaluation of antioxidant activity of some plant methanol extracts. *Biotechnology & Biotechnological Equipment*, 29(6), 1184-1189.
- Karker, M., Falleh, H., Msaada, K., Smaoui, A., Abdelly, C., Legault, J., & Ksouri, R. (2016). Antioxidant, anti-inflammatory and anticancer activities of the medicinal halophyte *Reaumuria vermiculata*. *EXCLI journal*, 15, 297.
- Kaur, S., & Mondal, P. (2014). Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. *J Microbiol Exp*, 1(1), 1-6.

- Köse, L. P., Gülçin, İ., Gören, A. C., Namiesnik, J., Martinez-Ayala, A. L., & Gorinstein, S. (2015). LC–MS/MS analysis, antioxidant and anticholinergic properties of galanga (*Alpinia officinarum* Hance) rhizomes. *Industrial Crops and Products*, 74, 712-721.
- Lee, J. Y., Hwang, W. I., & Lim, S. T. (2004). Antioxidant and anticancer activities of organic extracts from *Platycodon grandiflorum* A. De Candolle roots. *Journal of Ethnopharmacology*, 93(2), 409-415.
- Lobo, V., Patil, A., Phatak, A., & Chandra, N. (2010). Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy reviews*, 4(8), 118.
- Lobo, V., Patil, A., Phatak, A., & Chandra, N. (2010). Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacognosy reviews*, 4(8), 118.
- López, Z., Núñez-Jinez, G., Avalos-Navarro, G., Rivera, G., Salazar-Flores, J., Ramírez, J. A., & Knauth, P. (2017). Antioxidant and Cytotoxicological Effects of Aloe vera Food Supplements. *Journal of Food Quality*, 2017.
- Maizura, M., Aminah, A., & Wan Aida, W. M. (2011). Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract. *International Food Research Journal*, 18(2).
- Mehta, S. K., & Gowder, S. J. T. (2015). Members of Antioxidant Machinery and Their Functions. In *Basic Principles and Clinical Significance of Oxidative Stress*. InTech.
- Nagar, A., Sharma, V., & Chhipa, A. S. (2017). Role of antioxidants in biological system. *Mintage Journal of Pharmaceutical and Medical Sciences*, 7-12.
- Narayanaswamy, N., & Balakrishnan, K. P. (2011). Evaluation of some medicinal plants for their antioxidant properties. *International Journal of Pharm Tech Research*, 3(1), 381-385.
- Niki, E.; Shimaski, H.; M. Mino. (1994) Antioxidantism -- Free Radical and Biological Defence. Gakkai Syuppan Center, Tokyo.
- Papas, A. M. (1999). Diet and antioxidant status. *Food and Chemical Toxicology*, 37(9), 999-1007.
- Pinchuk, I., Shoval, H., Dotan, Y., & Lichtenberg, D. (2012). Evaluation of antioxidants: scope, limitations and relevance of assays. *Chemistry and physics of lipids*, 165(6), 638-647.
- Prejeena, V., Suresh, S. N., & Varsha, V. (2017). phytochemical screening, antioxidant analysis and antiproliferative effect of *costus pictus* d. don leaf extracts. *Cell*, 2, 0-974.
- Ravishankar, B., & Shukla, V. J. (2007). Indian systems of medicine: a brief profile. *African Journal of Traditional, Complementary and Alternative Medicines*, 4(3), 319-337.
- Said, O., Khalil, K., Fulder, S., & Azaizeh, H. (2002). Ethnopharmacological survey of medicinal herbs in Israel, the Golan Heights and the West Bank region. *Journal of ethnopharmacology*, 83(3), 251-265.

- Saranya, B., Sulfikarali, T., Chindhu, S., Muneeb, A. M., Leela, N. K., & Zachariah, T. J. (2017). Turmeric and cinnamon dominate in antioxidant potential among four major spices. *Journal of Spices and Aromatic Crops*, 26(1).
- Shafek, R. E., Shafik, N. H., Michael, H. N., El-Hagrassi, A. M., & Osman, A. F. (2015). Phytochemical studies and biological activity of *Dodonaea Viscosa* flowers extract. *Journal of Chemical and Pharmaceutical Research*, 7(5), 109-116.
- Sharma, H. M., Hanna, A. N., Kauffman, E. M., & Newman, H. A. (1992). Inhibition of human low-density lipoprotein oxidation in vitro by Maharishi Ayur-Veda herbal mixtures. *Pharmacology Biochemistry and Behavior*, 43(4), 1175-1182.
- Shebis, Y., Iluz, D., Kinel-Tahan, Y., Dubinsky, Z., & Yehoshua, Y. (2013). Natural antioxidants: function and sources. *Food and Nutrition Sciences*, 4(06), 643.
- Shori, A. B. (2015). Screening of antidiabetic and antioxidant activities of medicinal plants. *Journal of integrative medicine*, 13(5), 297-305.
- Shukla, S., Mehta, A., Mehta, P., & Bajpai, V. K. (2012). Antioxidant ability and total phenolic content of aqueous leaf extract of *Stevia rebaudiana* Bert. *Experimental and Toxicologic Pathology*, 64(7), 807-811.
- Sunil, K. (2014). The Importance of Antioxidant and their role in Pharmaceutical science. *Asian journal of research in chemistry and pharmaceutical sciences*, 1(1).
- Tag, H., Kalita, P., Dwivedi, P., Das, A. K., & Namsa, N. D. (2012). Herbal medicines used in the treatment of diabetes mellitus in Arunachal Himalaya, northeast, India. *Journal of Ethnopharmacology*, 141(3), 786-795.
- Tailor, C. S., & Goyal, A. (2014). Antioxidant activity by DPPH radical scavenging method of *Ageratum conyzoides* Linn. leaves. *American Journal of Ethnomedicine*, 1 (4), 244-249.
- Tohma, H., Gülçin, İ., Bursal, E., Gören, A. C., Alwasel, S. H., & Köksal, E. (2017). Antioxidant activity and phenolic compounds of ginger (*Zingiber officinale* Rosc.) determined by HPLC-MS/MS. *Journal of Food Measurement and Characterization*, 11(2), 556-566.
- Varsha, V., Suresh, S. N., & Prejeena, V. (2016). phytochemical screening, antioxidant property and anticancer potential against mcf-7 cell lines of *vernonia cinerea* leaves.
- Velavan, S., Nagulendran, K., Mahesh, R., & Begum, V. H. (2007). In vitro antioxidant activity of *Asparagus racemosus* root. *Pharmacognosy magazine*, 3(9), 26.
- Verma, M., & Kumar, A. (2017). Antimicrobial and antioxidant activity of whole plant extracts of *bacopa monnieri* (L.) pennell.
- Yan, X., Tang, J., dos Santos Passos, C., Nurisso, A., Simões-Pires, C. A., Ji, M., ... & Fan, P. (2015). Characterization of lignanamides from hemp (*Cannabis sativa* L.) seed and their antioxidant and acetylcholinesterase inhibitory activities. *Journal of agricultural and food chemistry*, 63(49), 10611-10619.

Yen, G. C., Chen, C. S., Chang, W. T., Wu, M. F., Cheng, F. T., Shiau, D. K., & Hsu, C. L. (2017). Antioxidant activity and anticancer effect of ethanolic and aqueous extracts of the roots of *Ficus beecheyana* and their phenolic components. *Journal of Food and Drug Analysis*.

Zahin, M., Aqil, F., & Ahmad, I. (2009). The in vitro antioxidant activity and total phenolic content of four Indian medicinal plants. *International Journal of pharmacy and pharmaceutical Sciences*, 1(1), 88-95.

© GSJ