

Acknowledgments

We acknowledge the support of USAID-ICARDA chickpea and TL III project for providing financial and technical support in conducting chickpea demonstration. We also thank farmers, development agents and wereda agricultural experts in the study without their involvement this trial would have not been realized. Finally, we gratefully acknowledge Holetta agricultural research center for material support.

Challenges

- **Ascochyta blight** - Though it was not appeared in the first year, this disease has occurred in the second year due to heavy rain and inflicted heavy yield loss on farmers that grew Natoli variety.

Reference

- CSA (2017). Agricultural sample survey report on area and production of crops in private peasant holdings, Meher season. Central Statistical Agency, Addis Ababa, Ethiopia.
- Geletu Bejiga and Yadeta Anbessa. (1994). Genetics and breeding research in chickpea. In: Asfaw *et al* (eds) *Cool-season food legumes of Ethiopia. proceedings of the first*
- Geletu Bejiga and Million Eshete (1996). Chickpea in Ethiopia. In: (Saxena N.P., Saxena M.C., Johansen C., Virmani S.M. and Harris H. (Eds.). *Adaptation of Chickpea in the West Asia and North Africa Region*. pp. 137-153, ICRISAT/ACARDA.
- Goa Y, Ashamo M (2016). Yield performance and Adaptation of Desi chickpea varieties in selected districts of Wolayta and Hadiya zone of south Ethiopia. *International journal of research-Grantha alayah* 4: 33-41
- Tufa B. and Abdala A. (2018). Pre-extension Demonstration of improved chickpea varieties in Konta special woreda of southern Nation Nationalities and peoples Regional State, Ethiopia *EPH-International journal of Agriculture and Environmental Research* 4: 1-6