[20] Hottenrott, K et Neumann, G. (2008). Methodik des Ausdauertrainings. Hofmann-Verlag, Schorndorf, p. 107-109.
[21] Imai, K., Sato, H., Hori, M., Kusuoka, H., Ozaki, H., Yokoyama, H etKamada, T. (1994). Vagally mediated heart rate recovery after exercise is accelerated in athletes but blunted in patients with chronic heart failure. Journal of the American College of Cardiology, 24(6), 1529-1535.
[22] Jouven, X. (2006). Réussir à ralentir son rythme cardiaque, Paris, INSERM, Paris.
[23] Kalliokoski, K., Nuutila., Laine, H et coll. (2002). Myocardial perfusion and perfusion reserve in endurance-trained men. Med Sci Sports Exerc; 34: 948-53.
[24] Kasmi.A., Mokrani, F. (2013). Effet de l'entrainement d'endurance sur la fréquence cardiaque, le poids corporel et les habitudes alimentaires. Sciences Humaines, n°39 pp 107- 116.
[25] Kraemer, R.R., Kraemer, G.R., Acevedo, E.O., Hebert, E.P., Temple, E., Bates, M, et coll. (1999). Effects of aerobic exercise on serum leptin levels in obese women. Eur J ApplPhysiol; 80:154–8.
[26] Marci nick, E., Potts, J., Schlabach, G., Will, S., Dawson, P. and Hurley, B. (1991) 'Effects ofStrength training on lactate threshold and endurance performance. Medicine andsciencesportexercise, pp. 739–743.
[27] Marquet, D., Hildes, J. et Schaefer, O. (1981). Le climat et la santé. In la santé et environnement: l'encyclopédie pour mieux vivre. OMS; 4: 168-183.
[28] McArdle, W.D., Katch, F.I and Katch, V.L. (2001). Exercise physiology energy, nutrition, andhuman performance. Lippincott Williams & Wilkins. USA. 5eedition.
[29] Mcardle., Katch et Katch. (2001). Physiologie de l'activité physique: Energie, Nutrition, Performance. Vigot, Paris; 558p.
[30] Milanovic, Z., Sporiš, G. and Weston, M. (2015) 'Effectiveness of High-IntensityIntervalTraining (HIT) and Continuous Endurance Training for VO2max Improvements: A SystematicReview and Meta-Analysis of Controlled Trials.', Sport. Med., 45, pp. 1469–1481
[31] Monod, H et Flandrois, R. (2000). Physiologie du sport, Masson, Paris; 378p.
[32] Pedro Leal et Autonne. (2017). Stage Exploration Fonctionnelle. Master En Sciences Du Sport. Université de Lausanne.
[33] Perini, R., Orizio, C., Comandè, A., Castellano, M., Beschi, M etVeicsteinas, A. (1989). Plasma norepinephrine and heart rate dynamics during recovery from sub maximal exercise in man. European journal of appliedphysiology and occupationalphysiology, 58(8), 879-883.
[34] Reiss, D et Prévost, P. (2013). La bible de la préparation physique. Paris: Amphora.