

<p>2</p>	<p>Weightmanagement supplements</p>	 <p>Source: deepbluehealtz.co.nz</p>
<p>3</p>	<p>Plant Organic Fertilizer</p>	 <p>Source: https://shopee.co.id/chifarm.html</p>
<p>4</p>	<p>Cosmetics</p>	 <p>Source : http://aquatonale.rs/en/chitosan-cellulaire/.html</p>

CONCLUSION

Based on the literacy study above, it can be concluded that the isolation of chitin from crab shells can be done by chemical methods, fermentation methods, and enzymatic methods. Of the three methods actually have the same steps as demineralization and deproteinization, the demineralization process aims to remove minerals in the crab shell, while deproteinization aims to remove the protein content in chitin.

REFERENCES

- Agustina, S., Swantara, IMD, & Suartha, IN (2015). Chitin isolation, characterization, and synthesis of chitosan from shrimp shells. *Journal of Chemistry*, 9(2), 271-278.
- ALFRIANTO, E. and E. LIVIAWATY 1992. *Crab Maintenance*, Kanisius Publishers, Yogyakarta: 74 pages.
- FUJAYA, Y., HY ZAINUDDIN, AZIZ, and ANWAR 200 I. Effect of Multivitamin Enrichment in Live Feed on Survival of Mangrove Crab Larvae. *Journal of Life*, 8(2): 50-52
- H. El Knidri, R. Belaabed, A. Addaou, A. Laajeb, and A. Lahsini. 2018. Extraction, chemical modification and characterization of chitin and chitosan, *Int. J. Biol. Macromol.*, vol. 120, pp. 1181– 1189.
- Hayes, M. 2012. *Marine Bioactive Compounds: Sources, Characterization and Applications*. United States: Springer US
- Hendri, J. 2008. *Technique of Enzymatically Deproteination of Crab Skin (Portunus pelagious) Using Pseudomonas aeruginosa Bacteria for Making Chitin Polymers and Their Deacetylation*. Seminar on Research Results and Community Service. Lampung University.
- Josua, VO, Edison, E., & Karnila, R. 2015. *The Effect of the Hydrochloric Acid Concentration to the Characteristics of Black Sea Cucumber (Holothuria Edulis) Chitin* (Doctoral dissertation, Riau University).
- Josua, VO, Edison, E., & Karnila, R. 2016. *The Effect of the Hydrochloric Acid Concentration to the Characteristics of Black Sea Cucumber (Holothuria Edulis) Chitin* (Doctoral dissertation, Riau University)
- Kaho, AR 2006. Chitosan "magic of Nature". <http://totalwellness.blogsome.com/2006/02/10/formalin-chitosan->
- Khoushab, F. and M. Yamabhai. 2010. Chitin Research Revised. *Marine Drugs*. Vol. 8: 1988-2012
- KT Oh, YJ Kim, VN Nguyen, WJ Jung, and RD Park. 2007. Demineralization of crab shell waste by *Pseudomonas aeruginosa* F722, *Process Biochem.*, vol. 42, no. 7, pp. 1069–1074,
- Kusumawati, N, 2009. Utilization of Shrimp Shell Waste as Raw Material for Making Ultrafiltration Membranes. *Inotek*. 13(2): 113-120
- NYBAKKEN, JW 1992. *Marine biology an ecological approach*, Translation. PT Gramedia. Jakarta: 459 p.
- Prashanth, KVH, and Tharanathan, RN., 2007, Chitin/Kitosan Modifications and Their Unlimited Application Potentia I-An Overview, *FoodSci. Tech*. 18:117-13
- Pratiwi, R. (2014). The benefits of chitin and chitosan for human life. *Oceana*, 39(1), 35-43.
- Purwatiningsih, S., Wukirsari, T. Sjahrizza, A., & Wahyono, D. 2009. *Chitosan is a Future Source of Biomaterials*. IPB Press. Bogor

- Rochima, E., Suhartono, MT, Syah, D., and Sugiyono., 2004, Characterization of Chitosan from Enzymatic Deacetylation by Chitin Deacetylase Isolate of *Bacillus Papandayan K29-14*, Padjadjaran University
- S. Hajji, O. Ghorbel-Bellaaj, I. Younes, K. Jellouli, and M. Nasri. 2015. Chitin extraction from crab shells by *Bacillus* bacteria. Biological activities of fermented crab supernatants, *Int. J. Biol. Macromol.*, vol. 79, pp. 167–173, .
- Sanjaya, I. & L. Yuanita, 2007. Adsorption of Pb (II) by Chitosan Isolation of Chitin from Mangrove Crab Shell (*Scylla* sp.) *Jurnal Basic Science*. 8(1): 30-36
- Sismaraini, D. (2015). Strategy for the development of chitin and chitosan industries in Indonesia (Doctoral dissertation, Bogor Agricultural University (IPB)).
- SLAHAINENIA, L. 2008. Biology of Mangrove Crabs (*Scylla* spp.) in Mangrove Ecosystems. Subang Regency, West Java. Graduate School. Bogor Agricultural University: 246 pages.
- Stephenson, W, Campbell B 1959, The Australian Portunids (Crustacea: Portunidae) III, The genus *Portunus*¶, *Aust J. mar. Freshwat. res.* 10:84 ± 124.
- Taufan, M. R S. & Zulfahmi, 2010. Utilization of Shrimp Shell Waste as Anti-Termite Material (Bio-termiticide) in Wooden Buildings, Thesis. Diponegoro University, Semarang, 44 pages
- Tong, Y., H. Guan, S. Wang, J. Xu, and C. He. 2011. Syntheses of Chitin-based Imprinting Polymers and Their Binding properties for Cholesterol. *Carbohydrate Research*. Vol. 346: 495-500
- Vargas M, Martinez CG. 2010. Recent Patents on Food Applications of Chitosan, *Recent Patents on Food, Nutrition & Agriculture*: 121-128
- Wei Zhang and Wenshui Xia. 2015. Effect of Media Milling on Lipid-lowering and Antioxidant Activities of Chitosan. *International Journal of Biological Macromolecules*. Vol. 73:1402-1405
- Wiyarsi, A. and E. Priyambodo. 2009. Effect of Chitosan Concentration from Shrimp Shell on Heavy Metal Absorption Efficiency. Department of Chemistry Education, Faculty of Mathematics and Natural Sciences, UNY. Yogyakarta.
- Xie, W., P. Xu, W. Wang, and Q. Liu. 2002. Preparation and antibacterial activity of a water-soluble chitosan derivative. *Carbohydrate Polymer* 50: 35-40