

its confirmation (Banda, 2002). Chickens less than 3 weeks of age present no clinical signs of disease but chickens greater than 3 weeks of age present clinical signs (Kegne and Chanie, 2014).

Gross and histopathological examinations of the bursa are used to diagnose IBD in young chickens or in those having maternal antibodies (Lukert and Saif, 2003). However, other methods used in diagnosis include isolation and detection of IBDV using embryonated chicken eggs, cell culture, reverse transcription polymerase chain reaction (RT-PCR) and serology, such as virus neutralisation, indirect ELISA and agar gel immune diffusion test (Brandt *et al*, 2001).

1.4.1 Virus Isolation

IBDV can be isolated (grown) on chicken embryo and primary cell culture especially chicken fibroblast cell. Isolation and identification of the agent provide the most certain diagnosis of IBD, but are not usually attempted for routine diagnostic purposes as the virus may prove difficult to isolate (Yousif, 2005).

I. Isolation of virus in embryos

The inoculation of bursal homogenates from IBDV infected chickens per the chorioallantoic membrane of 9-10 days old embryonated SPF (Specific-pathogen-free) chicken eggs is the most sensitive diagnostic method for virus isolation. The most sensitive route of inoculation is the Chorio allantoic membrane; the yolk sac route is also practicable(OIE, 2012). It is important especially for Wild-type IBDV, usually not replicating in conventional cell culture, can also be regenerated by the reverse genetics approach, but can grow in embryonated chicken eggs (Brandt *et al.*, 2001). Some strains grow well in embryos but are not readily adapted to grow in Chicken embryo fibroblasts or Chicken embryo kidney. Variant viruses however, do not kill the embryos but cause embryo stunting, discoloration, splenomegaly and hepatic necrosis (Lukert and Saif, 2003).

II. Isolation of virus in cell culture

IBDV grows in chicken embryo fibroblast and produces CPE characterized with an appearance of round retractile cells in about 3-5days (Orakpoghenor *et al.*, 2020).IBDV isolation in cell culture is not a routine use as a diagnostic test because the virus is difficult to culture. Some field strains failed to grow on cell cultures (Bumstead *et al.*, 1993). Wild-type Infectious bursal disease virus strains particularly very virulent strain do not grow in tissue culture. Comparison of genome sequence of wild-type and tissue culture adapted IBDV strains pointed to several mutations that might be responsible for invitro growth of IBDV in tissue culture (Islam, 2002).

1.4.2 Serological identification

For serological investigations, usually blood can be collected from the wing vein, allowed to clot and serum separated by centrifugation and stored at -20 °C until tested. Serological tests generally used for the detection of IBDV are Agar Gel Immuno diffusion test, ELISA and VN (Viral neutralization test) (Shaima *et al.*, 2014).

A. Agar gel immunodiffusion test(AGID)

The AGID is the most useful of serological tests for detection of specific antibodies in serum or for detecting viral antigen in bursal tissue (OIE, 2004). The test is specific because it cannot give false positive results, but it can give a false negative result. The presence of IBDV antigen can be detected in the bursal tissue by AGID for 5- 6 days post infection (Murphy *et al.*, 1999).

Blood samples should be taken early in the course of the disease, and repeat samples should be taken 3 weeks later. Because the virus spreads rapidly, only a small proportion of the flock needs to be sampled. Usually 20 blood samples are enough. For detection of antigen in the bursa of Fabricius, the bursae should be removed aseptically from about

ten chickens at the acute stage of infection. The bursae are minced using two scalpels in a scissor movement, then small pieces are placed in the wells of the AGID plate against known positive serum. Freeze– thaw cycles of the minced tissue may improve the release of IBDV antigens from the infected bursal tissue(OIE,2008).

B. Virus neutralization tests

VN tests are carried out in cell culture. The test is more laborious and expensive than the AGID test, but is more sensitive for detecting antibody (OIE,2004). This sensitivity is not required for routine diagnostic purposes, but may be useful for evaluating vaccine responses or for differentiating between IBDV 1 and 2 serotypes. To reduce test-to-test and operator-to-operator variation, a standard reference antiserum may be included with each batch of tests and the titer of the virus suspension must be reassessed in each new experiment using a sufficient number of repeats (wells) per virus dilution (OIE,2008).

C. Enzyme-linked immunosorbent assay

The enzyme linked immune sorbent assay (ELISA) is the most commonly used test for the detection and quantification of IBDV antibodies to check response to vaccination, natural field exposure and decay of maternal antibody titer. It is economical, simple, and quick tests a large number of samples at the same time and is adaptive to automation to computer software (Lukert and Saif, 2003).

The test sera are diluted according to the established protocol or kit instructions and each is dispensed into the requisite number of wells. After incubation under the appropriate conditions, the sera are discarded from the plates, and the wells are washed thoroughly. Anti-chicken immunoglobulins conjugated to an enzyme are dispensed into the wells, and the plates are again incubated as appropriate. The plates are emptied and rewashed before substrate containing a chromogen that gives a color change in the presence of the enzyme used is added to the plate. After a final incubation step, the substrate/chromogen reaction is stopped by addition of a suitable stopping solution and the colour reactions are

quantified by measuring the optical density of each well. The Sample to Positive (S/P) ratio for each test sample is calculated (OIE,2008)

1.4.3 Identification by molecular method

Molecular detection and characterisation, involving sequencing, and phenotypic and genotypic analyses have been utilised in the diagnosis of IBD. This method can detect the genome of IBDV, which is unable to grow in cell culture or embryonated eggs because it is unnecessary to grow the virus before amplification even when the virus is present in a very minute quantity and has lost its infectivity (Mittal *et al.*, 2005).The classical methods for molecular characterization and differentiation of IBDV field isolates include reverse transcriptase polymerase chain reaction (RT-PCR) and restriction fragment length polymorphism (RFLP), nucleotide sequence analysis, and quantitative real time RT-PCR (qRT-PCR) (OIE,2008).

Reverse Transcriptase polymerase chain reaction offers a rapid, highly sensitive and specific test for the confirmative diagnosis of the disease which would help in controlling the disease, thereby reducing the economic losses significantly. RT-PCR in combination with restriction enzyme analysis allows the rapid identification of vvIBDV. Nucleotide sequencing of RT-PCR products is widely used for further characterization of IBDV strains (Zierenberg *et al.*, 2004).

The VP2 gene of IBDV contains variable region which suggests the potential of this region for differentiation of IBDV strains. RT-PCR followed by digestion with multiple restriction enzymes or RFLP and nucleotide sequencing of VP2 gene have been used for differentiation of IBDV strains. The molecular differentiation of IBDV strains using VP2 has been improved by use of labeled probes in real-time RT-PCR (Jackwood, 2005).

1.4.4 Post mortem findings

Pathological change observed at the bursa of fabricius is characteristic and histopathological investigations combined with the demonstration of viral antigen by

immunohistochemistry confirm an IBDV infection (Zelege *et al*, 2005).Diagnostic lesion includes muscle hemorrhages and bursal enlargement . Pathognomonic gross lesions observed in the bursa of fabricius which show doubling in size with a yellowish gelatinous film that may surround it and sometimes hemorrhages may be seen on the surface of it (Liew *et al.*,2016).



Figure 2: gross lesions observed in the IBDV affected bursa of broiler chicken.

1.4.5 Histopathology examination

The lymphoid structures primarily affected by IBDV are BF, spleen, thymus, Harderian glands, caecal tonsils, gut-associated lymphoid tissue (GALT) and head-associated lymphoid tissues (HALT). Lymphocytic degeneration and necrosis in the medullary region of the BF at 1 day post infection are the first signs (Getachew and Fesseha, 2020).

Microscopic examination of tissues shows moderate hemorrhages in the muscles and kidneys and the spleen shows moderate lymphoid depletion in the lymphoid nodules. There is marked interfollicular oedema and depletion of 13 lymphocytes from the lymphoid nodules in the BFs. Other lymphoid nodules of the BF show degeneration and necrosis of lymphocytes and cystic cavitations with heterophil infiltrates.(Etteradossi and Saif, 2008).

1.5 Vaccine and Vaccination against IBD

Vaccination of chickens with high quality vaccines is the primary method of control of many poultry infectious diseases including IBD (Gumboro) disease (Dacic, 2008).

It becomes possible to safeguard chicken with a proper vaccination schedule. Rational vaccination schedules and strict biosecurity measures were indicated in many reports as essential tools for the control of IBD (Farooq, 2003). Even though there are different types of IBD vaccine being developed two of them are commonly used for the control of IBD. These are live attenuated vaccines, or inactivated oil-emulsion adjuvanted vaccines (Aregitu, 2018). Currently plant based and a live recombinant vaccine expressing IBDV antigens has also been licensed (Wagari, 2021)

1.5.1 *Live-attenuated vaccines*

Live viral vaccines can trigger immune system in the target host. They can replicate and induce both cellular and humoral immunity. They do not require an adjuvant to be effective and are suitable for mass administration to the chicken, but they may also have undesirable side effects. These include horizontal and vertical transmission (although the latter not in the case of IBD vaccines), reversion to virulence and vaccine reactions that may result in disease or production loss. In general, the live IBDV vaccines in use by the poultry industry have been attenuated by serial passage in tissue culture, eggs or embryo-derived tissues, with the aim of maintaining the immune response induced by the parent virus whilst attenuating the ability of the vaccine virus to cause clinical disease or significant immunosuppression (Schijns *et al.*, 2008).

1.5.2 *Inactivated vaccines*

Inactivated IBD vaccines are mostly formulated as water-in-oil emulsions, usually combining several antigens and have to be injected into each bird. It has been observed that inactivated IBD vaccine were able to induce IBDV-specific T-cell and inflammatory

responses in chickens (Rautenschlein *et al.*, 2005). It has been reported that inactivated IBD vaccines must have either a high or an optimized antigenic content in order to induce in breeders an immunity that helps protect the progeny from infection by variant IBDV strains (Muller *et al.*, 2012).

Killed-virus vaccines in an oil adjuvant are often used to boost levels of maternal antibodies and Confer longer lasting immunity in breeder hens. The duration and uniformity of this immunity may be influenced by the concentration and antigenic specificity of the vaccine strain. These vaccines are not ideal for stimulating a primary antibody response; therefore, they tend to be most effective in chicks that have been “primed” with a live virus vaccine or naturally infected through field exposure to IBDV (Etteradossi and Saif, 2008). Currently, many oiladjuvant vaccines contain both classic and variant IBDV strains. Killed-virus vaccines are administered by subcutaneous or intramuscular injection at sixteen to twenty weeks of age (Van den Berg , 2000).

1.5.3 New generation or Genetically-Engineered IBD vaccines.

Genetically engineered IBD vaccines have also been developed as a result of improved understanding on the molecular structure and immunology of IBDV. The viral capsid protein VP2, encoded by genomic segment A and derived from a large precursor protein VP0 by a series of proteolytic processes, carries immune determinants that control antibody-dependent neutralization and protection. Generally, these could be divided into two main categories, reflective of their replicative nature upon delivery into the chicken (Delmas, 2008).

1. Non-replicative IBD vaccines

Immunisation by DNA or subunit vaccines involves the use of non-replicating IBDV for induction of immune response in birds. DNA vaccination is based on direct inoculation of plasmid DNA encoding a target immunogen gene into subjects of study (Oshop, Elankumaran and Heckert, 2002). Under the influence of a mammalian promoter, the

target genes were expressed to produce proteins in vivo that are able to induce immune responses in the injected host. Repeated injections of DNA vaccines carrying the IBDV genes, either the polyprotein genes or gene of VP2 alone were shown to protect the chickens from challenge virus (Chen *et al.*, 2011).

However, the presence of MDA could affect the efficacy of DNA vaccines and a high dose of DNA vaccines was required to overcome the interference of MDA and induce immune response in chickens. It was shown that a booster vaccination with inactivated IBD vaccine after priming with DNA vaccine provided better and higher protection to the chickens compared to injection with DNA vaccines alone (Hsieh *et al.*, 2010).

2. Replication-competent IBD vaccines

Replication-competent viral vectors have been utilized to express and deliver immunogens of interest to chickens. In contrast to DNA and subunit vaccines, vaccination by live recombinant virus vectors employed the use of live and replicating virus to produce IBDV antigen upon in vivo infection. They have been shown to elicit both humoral and cell-mediated immune response in the chickens. As they could persistently infect the chickens, the potential of having a long-term protective immunity is high (Tsukamoto *et al.*, 2002).

Besides, the recombinant viral vectors are less sensitive to MDA and could therefore evade the neutralisation by the maternal anti-IBDV antibody (van den Berg, 2000). Several viruses have been experimented to express the VP2 protein of IBDV. This includes fowlpox virus, fowl adenovirus, Marek's disease virus, Newcastle disease virus, and avian adeno-associated virus among others (Tsukamoto *et al.*, 2002). The VP2 protein expressed in vivo from these various studies have been shown to confer from partial to full protection to vaccinated chickens from mortality, although they do not prevent the damage to the bursa (Zhou *et al.*, 2010).

1.5.4 Plant-Produced IBD vaccines

The plant-based expression system is a growing alternative platform for production and development of animal vaccines (Liew and Hair-Bejo, 2015). Being one of the pathogenic agents of importance in poultry, plant-based expression system using the stable (Wu *et al.*, 2004), transient (Gómez *et al.*, 2013), or chimeric viral particles (Chen *et al.*, 2012) approach has been used to produce IBD vaccine containing VP2 capsid protein. Transgenic rice expressing the VP2 protein was shown to protect the chickens from challenge following oral immunisation (Wu *et al.*, 2007).

Recently, the VP2 protein of IBDV has been transiently expressed in *Nicotiana benthamiana* leaves and extracted for subunit vaccination in chicken (Gómez *et al.*, 2013). The recombinant VP2 protein emulsified in oil adjuvant, injected intramuscularly to chicks at 18 days of age and followed by booster doses after 22 and 35 days, were shown to induce the production of anti-IBDV antibody with neutralising ability (Chen *et al.*, 2012).

1.5.5 In ovo Vaccination and post hatch vaccination

Recently, technology has been developed to deliver live vaccine into eggs during the incubation period. Live vaccine virus is blended with IBD antibody and the complex is injected in ovo at 18 days of incubation. The eggs go on to hatch and the vaccine virus is released when the chicks are about 7 days of age. In this way, the problem of maternally derived IBD antibody is overcome and the chicks are effectively immunized (OIE, 2012).

Compared to Post-hatch vaccination, In ovo injection of a live intermediate vaccine allowed faster recovery from bursa lesions although both methods exhibited similar protection against challenge (Rautenschlein and Haase, 2005). Although in ovo delivery of vaccines is an attractive alternative to post-hatch vaccination, various factors including the dosage, virulence, and efficacy, among others must be properly optimized before chasing large scale vaccinations (Corley and Giambrone, 2002).

2 CONCLUSIONS AND RECOMMENDATIONS

Infectious bursal disease is caused by IBD Virus that affects immune cells of chickens. It is mainly disease of young chickens between 3-6 weeks old and cause secondary problems due to the effect of the virus on the bursa of Fabricius. Diagnosis of IBD is depending on clinical signs, differential diagnosis, gross lesions, histopathological lesions, virus isolation, serological and molecular diagnosis. Isolation and identification of the agent can deliver the most confident diagnosis of Infectious bursal disease.

From the recommended serological tests for IBD virus, AGID is the simplest but least sensitive whereas ELISA is a rapid and sensitive method but cannot differentiate serotypes. Virus neutralization test is the golden standard and the only serologic test that differentiates antibodies of two serotypes and sensitive but it is more laborious and expensive than AGID. Molecular Identification Reverse Transcription-Polymerase Chain Reaction is used to detect IBDV without considering the viability of the virus by working on VP2 found on segment A of the viral Capsid.

Vaccination is the principal control measure of IBDV infection in chickens. From the available vaccine, live vaccine is more protective and most widely used IBD vaccine. Vaccination strategies in ovo, at-hatch or on-farm vaccinations, determines the choice of vaccines used in the farm. Therefore based on the above conclusion the following recommendations are forwarded:

- Virus neutralization test is the most sensitive but laborious and time consuming
- Molecular diagnostic technique reverse transcriptase polymerase chain reaction is relatively the easiest and sensitive one. .
- More safe and effective IBD vaccines that are affordable and readily available must be identified with further cost benefit analysis.
- In Ivo vaccination will be the promising best vaccination strategy against Infectious bursal disease.

3 REFERENCES

- Aregitu, M. (2018): Review on infectious Bursal disease: Threat for Ethiopian Poultry industry *.International Journal of Advanced Life Sciences.***11**:52-55.
- Banda, A.(2002): Characterization of Field Strains of Infectious Bursal Disease Virus (IBDV) Using Molecular Techniques *Tropical Animal Health and Production.***30**:70-80.
- Brandt, M., Yao, M., Liu, R., Heckert, A., Vakharia, N. (2001): Molecular determinants virulence, cell tropism, and pathogenic phenotype of IBDV. *J Virology.* **75**: 11974-11982.
- Bumstead, N., Reece, R. L., and Cook, J. K. A. (1993): Genetic differences in susceptibility of chickens lines to infection with infectious bursal disease virus. *Poult. Sci.* **72**: 403-410.
- Chen, T. H., Chen, T. H., Hu, C. C., Liao, J. T., Lee, C. W., Liao, J. W., Lin, M. Y., Liu, H. J., Wang, M. Y., Lin, N. S., & Hsu, Y. H. (2012): Induction of protective immunity in chickens immunized with plant-made chimeric bamboo mosaic virus particles expressing very virulent infectious bursal disease virus antigen. *Virus Research.* **166**:109-115.
- Chen, Y. Y., Hsieh, M. K., Tung, C. Y., Wu, C. C., & Lin, T. L. (2011): Infectious bursal disease DNA vaccination conferring protection by delayed appearance and rapid clearance of invading viruses. *Archives of Virology.***156**: 2241-2250.
- Corley, M. M., Giambrone, J. J., & Dormitorio, T. V. (2002): Evaluation of the immune response and detection of infectious bursal disease viruses by reverse transcriptase-polymerase chain reaction and enzyme-linked immunosorbent assay after in ovo vaccination of commercial broilers. *Avian Diseases.***4**:803-809.
- Dacic, M., Zugic, G., Petkovic, J .(2008): The effect of a gumboro disease control program on the reduction of economic losses. *Acta. Vet. (Beograd).* **58**: 53-62.

- Delmas, B. (2008): Birnaviruses. In B.W.J. Mahy & M.H.V. van Regenmortel (Eds.). *Encyclopedia of Virology 3rd edn.* Vol. **1**, (pp. 321-328).
- Eterradossi, N., & Saif, Y. M. (2008): Infectious bursal disease. *Diseases of Poultry* (pp. 185- 208).
- Farooq ,M., Durrani, F., Imran, N., Durrani, Z., Chand, N. (2003): Prevalence and economic losses due to infectious bursal disease in broilers in Mirpur and Kolti districts of Kashmir. *Int J Poultry Sci* **2**: 267-270.
- Getachew, Y and Fesseha, H. (2020): Infectious Bursal Disease in Poultry Production. *Global Veterinaria*. **4**: 185-196.
- Gómez, E., Lucero, M. S., Zoth, S. C., Carballeda, J. M., Gravisaco, M. J., & Berinstein, A. (2013): Transient expression of VP2 in *Nicotiana benthamiana* and its use as a plant-based vaccine against infectious bursal disease virus. *Vaccine*. **23**:2623-2627.
- Hsieh, M. K., Wu, C. C., & Lin, T. L. (2010): DNA-mediated vaccination conferring protection against infectious bursal disease in broiler chickens in the presence of maternal antibody. *Vaccine*, **28(23)**:3936-3943.
- Islam, M. R. (2002): Developing vaccines. Georg Forster.
- Jackwood D, Sommer S .(2005): Molecular studies on suspect very virulent infectious bursal disease virus genomic RNA samples. *Avian Dis*. **49**: 246- 251.
- Jackwood D, Sommer-Wagner.(2018): Genetic characteristics of infectious bursal disease viruses from four continents. *Virology*. **365**: 369- 375.
- Kegne T, Chanie M. (2014): Review on the incidence and pathology of infectious bursal disease. *British Journal of Poultry Sciences*. **3**: 68-77.
- Liew, P. S., & Hair-Bejo, M. (2015): Farming of plant-based veterinary vaccines and their applications for disease prevention in animals. *Advances in Virology*. <http://dx.doi.org/10.1155/2015/936940>.

- Liew, P. S., Nurulfiz , Abdul, Omar & Hair-Bejo, M. (2016): Vaccines and Vaccination against Infectious Bursal Disease of Chickens: Prospects and Challenges. *Pertanika Journal of Scholarly Research Reviews, PJSRR*. **2(2)**: 23-39.
- Lukert, P. D. and Saif, Y. M. (2003): Infectious bursal disease. In: Diseases of Poultry. 11th ed. Pages 161-180. Iowa State University Press, Ames.
- Mahgoub. H., Bailey. M. and Kaiser. P. (2012): An overview of infectious Bursal disease. *Arch. Virol.***157(11)**: 2047 - 2057.
- Mathivanan, K., Kumanan, K. and Nainar, A.M. (2004). *Vet. Res. Commun.*, **28**: 171-177.
- Mittal, N. D., Jindal, S. L., Gupta, Kataria, R. S. and Tiwari, A. K. (2005): Detection of Infectious Bursal Disease Virus in Field Outbreaks in Broiler Chickens by Reverse Transcription-Polymerase Chain Reaction. *International Journal of Poultry Science*. **4 (4)**: 239-243.
- Muller, H., Mundt, E., Eterradosi, N. and Islam, M.(2012): Current status of vaccines against infectious Bursal disease. *Avian. Pathol.* **41**: 133 - 39.
- Murphy, F. A., Gibbs, E. P. J., Horzinek, M. C., Studdert, M. J. (1999): Infectious bursal disease at Veterinary Virology. Third edition. *Academic press in United States of America*. 405-409.
- Mutinda .W.U. (2016).:Isolation, pathological and immunological characterization of Kenyan infectious bursal disease virus strains for vaccine development. *PhD thesis* University of Nairobi, Kenya.
- OIE. (2004)Infectious bursal disease (Gumboro disease) : Manual of diagnostic test and vaccine for terrestrial animals.
- OIE. (2008): Office International des Epizootics. Infectious Bursal Disease (Gumboro disease). In Manual of diagnostic tests and vaccines for terrestrial animals.*OIE,4 th edition, Paris (in press)*.

- OIE. (2012) :Office of International des Epizooties. Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. Infectious Bursal Disease. **3**: 549-565.
- Orakpoghenor,O., Sunday,B., Oladele & Paul. A., Abdu (2020): Infectious Bursal Disease: Transmission, Pathogenesis, Pathology and Control - An Overview, *World's Poultry Science Journal*, **76**:292-303.
- Oshop, G. L., Elankumaran, S., & Heckert, R. A. (2002): DNA vaccination in the avian. *Veterinary Immunology and Immunopathology*. **89**: 1-12.
- Rautenschlein S, Samson-Himmelstjerna G, Haase C,. (2007): A comparison of immune response to infection with virulent infectious bursal disease virus (IBDV) between specific-pathogen-free chickens infected at 12 and 28 days of age. *Vet. Immunol Immunopathol*. **115**: 251-260.
- Rautenschlein, S., & Haase, C. (2005): Differences in the immunopathogenesis of infectious bursal disease virus (IBDV) following in ovo and post-hatch vaccination of chickens. *Veterinary Immunology and Immunopathology*. **106**:139-150.
- Rautenschlein, S., Kraemer, C., Vanmarcke, J., & Montiel, E. (2005): Protective efficacy of intermediate and intermediate plus infectious bursal disease virus (IBDV) vaccines against very virulent IBDV in commercial broilers. *Avian Diseases*. **49**:231-237.
- Saidi.R., Shirima.G. , Buzal.J., and Kasanga.C.(2020):Factors associated with infectious bursal disease vaccination failure in Dar es salaam, Tanzania . *Journal of Veterinary Medicine and Animal Health* .PP:21-22.
- Schijns, V.E.J.C., Sharma, J. & Tarpy, I. (2008): Practical aspects of poultry vaccination. *Avian Immunology* 1st edn (pp. 373-393). London: Academic Press.
- Shaimaa I, Mohamed MH, Mohga, FB.(2014): Effect of infectious bursal disease field vaccines on avian influenza vaccination immunity. *Alexandria J of Vet Sci*.**43**: 33-36.

- Tsukamoto, K., Saito, S., Saeki, S., Sato, T., Tanimura, N., Isobe, T., Mase, M., Imada, T., Yuasa, N., & Yamaguchi, S. (2002): Complete, long-lasting protection against lethal infectious bursal disease virus challenge by a single vaccination with an avian herpesvirus vector expressing VP2 antigens. *Journal of Virology*. **76(11)**: 5637-5645.
- Van den Berg, T. P. (2000): Acute infectious bursal disease in poultry: a review. *Avian Pathology*. **29(3)**: 175-194.
- Wagari, A. (2021): A Review on Infectious Bursal Disease in Poultry. *Health Econ Outcome Res Open Access*. **7(2)**: 167 (018- 023).
- Wu, C. C., Lin, T. L., Zhang, H. G., Davis, V. S. and Boyle, J. A. (1992): Molecular detection of infectious bursal disease virus by polymerase chain reaction. *Avian Dis*. **36(2)**: 221-6.
- Wu, H., Singh, N. K., Locy, R. D., Scisum-Gunn, K., & Giambone, J. J. (2004): Expression of immunogenic VP2 protein of infectious bursal disease virus in *Arabidopsis thaliana*. *Biotechnology Letters*. **26(10)**: 787-792.
- Wu, J., Yu, L., Li, L., Hu, J., Zhou, J., & Zhou, X. (2007): Oral immunization with transgenic rice seeds expressing VP2 protein of infectious bursal disease virus induces protective immune responses in chickens. *Plant Biotechnology Journal*. **5(5)**: 570-578.
- Yip W, Hon C, Zeng UF, Leung, C. (2012): Cell culture-adapted IBDV uses endocytosis for entry in DF-1 chicken embryonic fibroblasts. *Virus Res*. **165**: 9-16.
- Yousif, MGO. (2005): Evaluation of various techniques used for the diagnosis of infectious bursal disease and its prevalence in Khartoum State. *MVSc thesis*. 99p. DOI 10.1008/s11250-012-0280-6.

- Zahid,B., Aslam,A., Tipu .Y, Yaqub,T,. and Butt.,T .(2016): Conventional and Molecular Detection of Infectious Bursal Disease Virus in Broiler Chicken .Pakistan J. Zool., vol. **48(2)**: pp. 601-603.
- Zelege A, Gelaye E, Sori T, Ayelet G, Sirak A, Zekarias B (2005b): Investigation on Infectious Bursal Disease outbreak in Debre Zeit, Ethiopia, *Int. J. Poult. Sci.* **4(7)**:504-506.
- Zhou, X., Wang, D., Xiong, J., Zhang, P., Li, Y., & She, R. (2010): Protection of chickens, with or without maternal antibodies, against IBDV infection by a recombinant IBDV-VP2 protein. *Vaccine.* **28(23)**:3990-3996.
- Zierenberg, K., R. Raue, H. Nieper, M.R. Islam, N. Eterradosi, D. Toquin and H. Müller. (2004): Generation of serotype 1/serotype 2 reassortant viruses of the infectious bursal disease virus and their investigation in vitro and in vivo. *Virus Research.* **105(1)**: 23-34.

