

Table 1: Distribution of Age by Gender

Age Group	GENDER		TOTAL
	Male (%)	Females (%)	
10-19	0 (00.0)	1 (1.6)	1 (0.9)
20-29	6 (13.3)	5 (8.1)	11 (10.3)
30-39	5 (11.1)	8 (12.9)	14 (12.1)
40-49	11 (24.4)	17 (27.4)	28 (26.2)
50-59	15 (33.3)	17 (27.4)	32 (29.9)
60-69	5 (11.1)	9 (14.5)	14 (13.1)
70-80	3 (6.7)	5 (8.1)	8 (7.5)
TOTAL	45 (100.0)	62 (100.0)	107 (100.0)

Table 2: Clinical Features at Presentation

Clinical Feature	Frequency	% Frequency
Throbbing Headache	98	91.6
Altered Consciousness	54	50.0
Vomiting	30	28.0
Meningism*	27	25.2
Motor weakness	26	24.3
Dysphasia	17	15.9
Anisocoria	7	6.5
Seizures	6	5.6

**Signs of meningeal irritation*

Table 3: Sites of aneurysm rupture by gender

VESSEL	GENDER		TOTAL (%)
	Male (%)	Female (%)	
Anterior Circulation	41 (89.1)	58 (95.1)	99 (92.5)
Anterior cerebral artery	4 (8.7)	5 (8.2)	9 (8.4)
Anterior communicating artery	20 (43.5)	15 (24.6)	35 (32.7)
Middle cerebral artery	11 (23.9)	21 (34.4)	32 (29.9)
Posterior communicating artery	4 (8.7)	9 (14.8)	13 (11.2)
ICA** -related branches	4 (8.7)	9 (14.8)	13 (11.2)
Posterior Circulation	5 (10.9)	3 (4.9)	8 (7.5)
Posterior cerebral artery	0 (0.0)	1 (1.6)	1 (0.9)
Basilar artery	2 (4.3)	0 (0.0)	2 (1.9)
Vertebral artery	1 (2.2)	0 (0.0)	2 (1.9)
TOTAL	46 (43.0)	61 (57.0)	107 (100.0)

**Internal carotid artery

Table 4: Factors affecting Outcome at discharge among Patients

	Good outcome*	Poor outcome**	Total	Significant
WFNS Grade at Admission				
1 -2	58(54.2%)	16(15.0%)	74(69.2%)	X ² =4.895, p-value 0.027
3-5	19(17.8%)	14(13.0%)	33(30.8%)	
Total	77(72%)	30(28%)	107(100%)	
Sex of Patients				
Female	42(44.6%)	20(17.4%)	62(57.9%)	X ² =1.3016, p-value0.254
Male	35(32.4%)	10(12.6%)	45(42.1%)	
Total	77(72%)	30(28%)	107(100%)	

*GOS 4&5, ** GOS 1 - 3