

- Okutucu, B., Dincer, A., Habib, O. and Zihnioglu, F. (2007) Comparison of five methods for determination of total plasma protein concentration, *J. Biochem. Bioph. Meth.*, 70, 709–711,
- Pereira MD, Eleutherio ECA, Panek AD. 2001. Acquisition of tolerance against oxidative damage in *Saccharomyces cerevisiae*. *BMC Microbiology* 1:11–20.
- Saka, W., Akigbe, R.E., Popoola, O.T, and Oyekunle, O.S. (2012) Changes in Serum Electrolytes, Urea, and Creatinine in *Aloe Vera*-treated Rats *J Young Pharm*4(2); 78- 81
- Scandalios J.G. 1993. *Oxygen stress and superoxide dismutases*. *Plant Physiol* 101:7–12.
- Schalm, O. W., Jain, N.C and Carrol, E.J. (1975). *Veterinary Haematology*, 3rd Edn., Lea and Febiger, Philadelphia, pp. 197 - 199
- Sehrawat A, Sultana S. (2006) *Tamarix gallica* ameliorates thioacetamide-induced hepatic oxidative stress and hyperproliferative response in Wistar rats. *J Enzyme Inhib Med Chem*. 21(2):215-23.
- Sims, G.K. (2006) Using the Berthelot Method for Nitrite and Nitrate Analysis. *Soil Sci. Soc. Am. J.* 70 (3): 1038.
- Vasudha K .C, Nirmal Kumar A, Venkatesh T (2006). Studies on the age dependent changes in serum adenosine deaminase activity and its changes in hepatitis. *Indian J Clin Biochem.*, 21: 116 - 120.
- Yakubu M.T., Akanji MA., and Oladiji AT., (2007). Haematological parameters of male albino rats following chronic administration of aqueous extracts of *fedogia agrestis* stem. *Pharmacognosy magazine* 3: 34